Document Identifier: **PMC8771116**

SciScore Report

Below you will find your SciScore report containing three tables. Your score is calculated based on adherence to scientific rigor criteria (Table 1) and identification of key biological resources (Table 2). Table 3 contains statistical tests and oligonucleotides but is not scored. If SciScore makes any mistakes, please contact us to help us learn and improve.

Table 1: Rigor Adherence Table

Ethics
IACUC: Surgery was approved by the Animal Ethics Committee of Nantong University , China (approval No. 20190226-001) on February 26 , 2019 .
Euthanasia Agents: , Darmstadt , Germany) , including 42 mgkg magnesium sulfate , 85 mgkg trichloroacetaldehyde monohydrate , and 17 mgkg sodium pentobarbital , rat sciatic nerves were crushed using forceps (Beaver-Visitec International , Inc . , Waltham , MA , USA) (three times , 10 secondstime) .
Inclusion and Exclusion Criteria
not detected.
Attrition
not detected.
Sex as a biological variable
Materials and Methods Bioinformatic analysis High-throughput data of RNA in male Sprague-Dawley rat sciatic nerve stumps (0, 1, 4, 7, and 14 days post-injury) and L4-5 dorsal root ganglia (0, 3, 9 hours, 1, 4, and 7 days post-injury) were preserved in the National Center for Biotechnology Information with accession numbers SRP113121 (Zhao and Yi, 2019) and SRP200823 (Shen et al., 2020),
Subject Demographics
Weight: Animal surgery and tissue collection A total of 15 clean rats (male , around 8 weeks , 180-220 g) were obtained from Nantong University Experimental Animal Center (animal license Nos .
Randomization
The 15 rats were randomly divided into five groups (designated as 0 , 1 , 4 , 7 , and 14 days n = 3 rats in each group) .
Blinding
not detected.
Power Analysis
not detected.

Replication	
Number: Data were summarized from three independent experiments and are presented as the mean +- standard error of the mean (SEM).	
Data Information	
Identifiers: Materials and Methods Bioinformatic analysis High-throughput data of RNA in male Sprague-Dawley rat sciatic nerve stumps (0, 1, 4, 7, and 14 days post-injury) and L4-5 dorsal root ganglia (0, 3, 9 hours, 1, 4, and 7 days post-injury) were preserved in the National Center for Biotechnology Information with accession numbers SRP113121 (Zhao and Yi, 2019) and SRP200823 (Shen et al., 2020),	SRP113121 SRP200823

Table 2: Key Resources Table

Your Sentences	REAGENT or RESOURCE	SOURCE	IDENTIFIER
Antibodies			
Table 1 List of antibodies used in this study Antibody Source Catalog number Species RRID Dilution Incubation time Temperature Ki67 Abcam , Cambridge , MA , USA ab16667 Rabbit AB_302459 1:500 Overnight 4degC p16 Signalway Antibody , College Park , MD , USA 32050 Rabbit 1:200 Overnight 4degC CD68 Abcam , Cambridge , MA , USA	Ki67	Abcam	(Abcam Cat# ab16667, RRID:AB_302459)(link)
	p16		
	CD68		
ab31630 Mouse AB_1141557 1:100 Overnight 4degC P4HB Abcam , Cambridge , MA , USA			(Abcam Cat# ab31630, RRID:AB_1141557)(link)
ab2792 Mouse AB_303304 1:100 Overnight 4degC S100b MilliporeSigma , St Louis , MO , USA S2532 Mouse AB_477499 1:400 Overnight 4degC Cy3 goat anti-rabbit IgG (H+L) Proteintech Group , Rosemont , IL , USA SA00009-2 Goat AB_2890957 1:400 2 h Room temperature Alexa Fluor 488 goat anti-mouse IgG (H+L) Abcam , Cambridge , MA , USA ab150117 Goat AB_2688012 1:400 2 h Room temperature P4HB: Prolyl 4-hydroxylase subunit beta .			(Abcam Cat# ab2792, RRID:AB_303304)(link)
	USA Mouse	Sigma-Aldrich	(Sigma-Aldrich Cat# S2532, RRID:AB_477499)(link)
	Cy3 goat anti-rabbit IgG	Proteintech	(Proteintech Cat# SA00009-2, RRID:AB_2890957)(link)
	Alexa Fluor goat anti-mouse	Abcam	(Abcam Cat# ab150117, RRID:AB_2688012)(link)
Collected tissue sections were exposed to rabbit anti-S100b (primary antibody , 1:400 Cat ab52642 , RRID AB_882426 , Abcam , Cambridge , MA , USA) overnight at 4degC and Cy3 goat anti-rabbit IgG (H+L) (secondary antibody , 1:400 Cat SA00009-2 , RRID AB_2890957 , Proteintech Group , Rosemont , IL , USA) for 2 hours at room temperature.	anti-S100b (	Abcam	(Abcam Cat# ab52642, RRID:AB_882426)(link)
	anti-rabbit IgG		
Experimental Models: Organisms/Strains			
Materials and Methods Bioinformatic analysis High-throughput data of RNA in male Sprague-Dawley rat sciatic nerve stumps (0, 1, 4, 7, and 14 days post-injury) and L4-5 dorsal root ganglia (0, 3, 9 hours, 1, 4, and 7 days post-injury) were preserved in the National Center for Biotechnology Information with accession numbers SRP113121 (Zhao	Sprague-Dawley		Suggestion: RRID:RGD_70508(link)

and Yi, 2019) and SRP200823 (Shen et al., 2020),			
Software and Algorithms			
KEGG) database (map04218	KEGG		
The b-galactosidase-positive staining area was measured using ImageJ software (version 1.48 National Institutes of Health , Bethesda , MD , USA)	ImageJ		Suggestion: ImageJ, RRID:SCR_003070(link)
One-way analysis of variance with Dunnetts multiple comparisons post hoc test was performed to compare the differences among injured nerve stumps and the uninjured control (day 0) group using GraphPad Prism 6.0 (GraphPad Software, San Diego, CA, USA).	GraphPad		Suggestion: GraphPad Prism, RRID:SCR_002798(link)

Other Entities Detected

Your Sentences	Recognized Entity
Statistical Tests	
One-way analysis of variance with Dunnetts multiple comparisons post hoc test was performed to compare the differences among injured nerve stumps and the uninjured control (day 0) group using GraphPad Prism 6.0 (GraphPad Software, San Diego, CA, USA).	analysis of variance
	Dunnetts multiple comparisons post hoc test
Oligonucleotides	
Table 2 List of primer sequences used in this study Full name Abbreviation Forward Revers Cyclin dependent kinase inhibitor 1A Cdkn1a 5-GGT GAT GTC CGA CCT GTT CC-3 5-ACG CTC CCA GAC GTA GTT GC-3 Cyclin dependent kinase inhibitor 2A Cdkn2a 5-AAC ACT TTC GGT CGT ACC CC-3 5-CCC AGC GGA GGA GAG TAG AT-3 ATM serinethreonine kinase Atm 5-AGG TGT GTA ATG AAG CAG GTC T-3 5-GAA CCG CGC TAA TGA GAG GA-3 Cyclin dependent kinase 1 Cdk1 5-AGG ACT CCA GGC TGT ATC TCA T-3 5-TAT CGG TAT TCC AAA CGC TCT-3 Minichromosome maintenance complex component 5 Mcm5 5-TGA GAC AAA AGG GGA GGA CA-3 5-GCT CGG CAG TAA GCA ATG AA-3 Cyclin F Ccnf 5-CAC TGG GTT CTC CTA CAG CG-3 5-TCT CTC GCT TCC GTT TGC TT-3 Dihydrofolate reductase Dhfr 5-GCA AGA ACG GAG ACT TAC CCT-3 5-GTT TTC CTA CCC ATA ATC ACC AG-3 TIMP metallopeptidase inhibitor 1 Timp1 5-CCT CTG GCA TCC TCT TGT TG-3 5-CGT	5-GGT GAT GTC CGA CCT GTT CC-3 5-ACG CTC CCA GAC GTA GTT GC-3
	5-ACG
	5-AAC ACT TTC GGT CGT ACC CC-3 5-CCC AGC GGA GGA GAG TAG AT-3
	5-CCC
	5-AGG TGT GTA ATG AAG CAG GTC T-3 5-GAA CCG CGC TAA TGA GAG GA-3
	5-GAA
	5-AGG ACT CCA GGC TGT ATC TCA T-3 5-TAT CGG TAT TCC AAA CGC TCT-3
	5-TAT
	5-TGA GAC AAA AGG GGA GGA CA-3 5-GCT CGG CAG TAA GCA ATG AA-3
	5-GCT
	5-CAC TGG GTT CTC CTA CAG CG-3 5-TCT CTC GCT TCC GTT TGC TT-3
	5-TCT
	5-GCA AGA ACG GAG ACT TAC CCT-3 5-GTT TTC CTA CCC ATA ATC ACC AG-3
	5-GTT
	5-CCT CTG GCA TCC TCT TGT TG-3
Glyceraldehyde-3-phosphate dehydrogenase Gapdh 5-ACA GCA ACA GGG TGG TGG AC-3	5-ACA GCA ACA GGG TGG TGG AC-3
Unresolved RRIDs	
Table 1 List of antibodies used in this study Antibody Source Catalog number Species RRID Dilution Incubation time Temperature Ki67 Abcam , Cambridge , MA , USA ab16667 Rabbit AB_302459 1:500 Overnight 4degC p16 Signalway Antibody , College Park , MD , USA 32050 Rabbit 1:200 Overnight 4degC CD68 Abcam , Cambridge , MA , USA	RRID Dilution

SciScore is an automated tool that is designed to assist expert reviewers by finding and presenting formulaic information scattered throughout a paper in a standard, easy to digest format. ***SciScore is not a substitute for expert review.*** SciScore also checks for the presence and correctness of several unique identifiers, including RRIDs (research resource identifiers) in the manuscript, detects sentences that appear to be missing RRIDs, and can even suggest RRIDs under certain circumstances. **All RRID suggestions should be verified;** only the author can know whether the suggestions are correct.

For a full description of scored criteria and tips for improving your score, please see https://www.scicrunch.com/sciscorereport-faq