Document Identifier: **PMC8771815**

SciScore Report

Below you will find your SciScore report containing three tables. Your score is calculated based on adherence to scientific rigor criteria (Table 1) and identification of key biological resources (Table 2). Table 3 contains statistical tests and oligonucleotides but is not scored. If SciScore makes any mistakes, please contact us to help us learn and improve.

Table 1: Rigor Adherence Table

Ethics
IRB: Study approval was done by the Ethics Committee of the Iran University of Medical Sciences (Code: IR.IUMS.FMD.REC.1399.119).
Consent: Written informed consent was obtained from all patients after a detailed explanation of the study to the patients.
Inclusion and Exclusion Criteria
Study Outcomes The primary outcomes of our study were the effects of the two methods on pain score and the total analgesic requirement in the first 24 hours after surgery, and the secondary outcomes were the duration of hospital stay and the incidence of nausea and vomiting in the first 24 hours after surgery. 3.3. Eligibility Criteria The inclusion criteria included age between 18 and 50 years, transgender confirmation, being single, the need for mastectomy, and ASA I-II.
Attrition
not detected.
Sex as a biological variable
not detected.
Subject Demographics
Age: Study Outcomes The primary outcomes of our study were the effects of the two methods on pain score and the total analgesic requirement in the first 24 hours after surgery, and the secondary outcomes were the duration of hospital stay and the incidence of nausea and vomiting in the first 24 hours after surgery. 3.3. Eligibility Criteria The inclusion criteria included age between 18 and 50 years, transgender confirmation, being single, the need for mastectomy, and ASA I-II.
Randomization
Study Design, Patients, and Ethics This was a single-blind randomized clinical trial.
Blinding
Follow-up and Data Collection The senior resident, who performed all assessments, data collection, and follow-ups, was unaware of patients' grouping (blinding).
Power Analysis

Considering the alpha cut-off of 0.05 and a power of 80%, the sample size was calculated to be 23 for each group, but because of the possibility of sample dropout, we mustered 50. 3.2.
Replication
not required.

Table 2: Key Resources Table

Your Sentences	REAGENT or RESOURCE	SOURCE	IDENTIFIER
Software and Algorithms			
Pain intensity at 3, 6, 12, and 24 hours after surgery, upper limb paresthesia, frequency of nausea and vomiting, shortness of breath, hematoma, and the length of hospital stay were also assessed. 3.9. Statistical Analysis Data were analyzed using SPSS version 23 software (SPSS Inc. Chicago, The USA).	SPSS		Suggestion: SPSS, RRID:SCR_002865(link)

Other Entities Detected

Your Sentences	Recognized Entity
Statistical Tests	
A chi-square test was used to compare qualitative variables.	chi-square test
Besides, quantitative variables were compared by the t test if they followed a normal distribution or the Mann-Whitney test if they did not have a normal distribution.	t test
	Mann-Whitney test
Pearson's or Spearman’s correlation test was used to examining the correlation between variables.	Pearson's
	Spearman’s correlation test

SciScore is an automated tool that is designed to assist expert reviewers by finding and presenting formulaic information scattered throughout a paper in a standard, easy to digest format. ***SciScore is not a substitute for expert review.*** SciScore also checks for the presence and correctness of several unique identifiers, including RRIDs (research resource identifiers) in the manuscript, detects sentences that appear to be missing RRIDs, and can even suggest RRIDs under certain circumstances. **All RRID suggestions should be verified;** only the author can know whether the suggestions are correct.

For a full description of scored criteria and tips for improving your score, please see https://www.scicrunch.com/sciscorereport-faq