Document Identifier: **PMC8789530**

SciScore Report

Below you will find your SciScore report containing three tables. Your score is calculated based on adherence to scientific rigor criteria (Table 1) and identification of key biological resources (Table 2). Table 3 contains statistical tests and oligonucleotides but is not scored. If SciScore makes any mistakes, please contact us to help us learn and improve.

Table 1: Rigor Adherence Table

Ethics
IRB: Xenograft model Animal experiments were performed according to protocols approved by the ethical committee of Guangzhou University of Chinese Medicine Institutional Animal Care and Use Committee.
IACUC: Xenograft model Animal experiments were performed according to protocols approved by the ethical committee of Guangzhou University of Chinese Medicine Institutional Animal Care and Use Committee.
Euthanasia Agents: After treatment for 21 days, the mice were sacrificed by cervical dislocation method, and tumor tissues were collected for immunohistochemical staining (IHC) and Western blot.
Inclusion and Exclusion Criteria
not detected.
Attrition
not detected.
Sex as a biological variable
not detected.
Subject Demographics
Age: Five-week-old BALBC nude mice were used for in vivo anti-SCLC effect of Sg.
Randomization
not detected.
Blinding
not detected.
Power Analysis
not detected.
Replication
not required.

Cell Line Authentication	
not detected.	
Data Information	
Identifiers: Gene expression analysis from the public database Data from the Gene Expression Omnibus (GEO, www.ncbi.nlm.nih.govgeo) was used to analyze CDKN1A mRNA expression in SCLC, GSE6044 20 and GSE40275 21, a total of 19 normal lung and 17 SCLC samples were included in the present study.	GSE6044 GSE40275

Table 2: Key Resources Table

Your Sentences	REAGENT or RESOURCE	SOURCE	IDENTIFIER
Antibodies			
Antibodies for CDKN1A (A19094), CDKN1B (A19095), CDKN1C (A2060) and p53 (A0263) were purchased from ABclonal Technology.	CDKN1A		
	CDKN1B		
	CDKN1C		
	p53		
Next day, the membranes were incubated for 1 h at room temperature with an HRP-conjugated anti-rabbit or anti-mouse secondary antibody, The immunoreactive proteins were visualized with SuperSignal West Dura Chemiluminescent (Thermo Fisher).	anti-rabbit		
	anti-mouse		
Experimental Models: Cell Lines			
Materials and methods Cell culture and reagents Human SCLC cell line NCI-H1688 and human lung bronchial epithelial cell line BEAS-2B were obtained and authenticated from the Shanghai Institute of Cell Biology, Chinese Academy of Sciences (Shanghai, China).	BEAS-2B		Suggestion: RRID:CVCL_0168(link)
NCI-H82, NCI-H526 and 16HBE were purchased and authenticated from the ATCC.	NCI-H82		Suggestion: (RRID:CVCL_1591)(link)
	NCI-H526		
For first round screeding, NCI-H1688 cells were treated with each compound at a concentration of 30 uM.	NCI-H1688		Suggestion: (RRID:CVCL_1487)(link)
To establish stably transfected cell lines, lentivirus was produced by HEK-293T cells with the second-generation packaging system pMD2.G and psPAX2, and harvested at 48 h after transfection.	HEK-293T		Suggestion: RRID:CVCL_0063(link)
Experimental Models: Organisms/Strains			
Five-week-old BALBC nude mice were used for in vivo anti-SCLC effect of Sg.	BALBC nude		
2 x 106 NCI-H1688 cells were injected subcutaneously into athymic nude mice.	athymic nude		
Recombinant DNA			

Plasmids and cell transfections Short hairpin RNA (shRNA) sequences were cloned into psiF-copGFP vectors (System Biosciences, Mountain View, CA).	psiF-copGFP		
To establish stably transfected cell lines, lentivirus was produced by HEK-293T cells with the second-generation packaging system pMD2.G and psPAX2, and harvested at 48 h after transfection.	pMD2.G		
	psPAX2		
Software and Algorithms			
b-actin antibody (H11459) was purchased from Sigma as a loading control.	Sigma		Suggestion: sigma-tau (RRID:SCR_000488)(link)
Images were acquired using an Olympus IX51 microscope and processed using cellSens Dimension software.	cellSens		
Ingenuity Pathway Analysis (IPA) was used to identify enriched molecular pathways, gene networks, canonical pathways, and cellular biological consequences.	Ingenuity Pathway Analysis		Suggestion: Ingenuity Pathway Analysis, RRID:SCR_008653(link)
Protein-protein interaction (PPI) analysis of genes was processed by STRING (https:string-db.org) and visualized by Cytoscape.	STRING		
	Cytoscape		Suggestion: Cytoscape (RRID:SCR_003032)(link)
Gene expression analysis from the public database Data from the Gene Expression Omnibus (GEO, www.ncbi.nlm.nih.govgeo) was used to analyze CDKN1A mRNA expression in SCLC, GSE6044 20 and GSE40275 21, a total of 19 normal lung and 17 SCLC samples were included in the present study.	Gene Expression Omnibus		Suggestion: Gene Expression Omnibus (GEO) (RRID:SCR_005012)(link)
All statistical analyses were carried out using SPSS software (version 22).	SPSS		Suggestion: SPSS, RRID:SCR_002865(link)

Other Entities Detected

Your Sentences	Recognized Entity
Statistical Tests	
An unpaired two-tailed Students t-test was performed for two-group comparisons and one-way ANOVA analysis (Post hoc: LSD) was performed for multiple group comparisons.	Students t-test
	ANOVA
For the survival analysis, Kaplan-Meier curves were analyzed by log rank test.	log rank test

SciScore is an automated tool that is designed to assist expert reviewers by finding and presenting formulaic information scattered throughout a paper in a standard, easy to digest format. ***SciScore is not a substitute for expert review.*** SciScore also checks for the presence and correctness of several unique identifiers, including RRIDs (research resource identifiers) in the manuscript, detects sentences that appear to be missing RRIDs, and can even suggest RRIDs under certain circumstances. **All RRID suggestions should be verified;** only the author can know whether the suggestions are correct.

For a full description of scored criteria and tips for improving your score, please see https://www.scicrunch.com/sciscorereport-faq