Document Identifier: **PMC8799151**

SciScore Report

Below you will find your SciScore report containing three tables. Your score is calculated based on adherence to scientific rigor criteria (Table 1) and identification of key biological resources (Table 2). Table 3 contains statistical tests and oligonucleotides but is not scored. If SciScore makes any mistakes, please contact us to help us learn and improve.

Table 1: Rigor Adherence Table

Ethics
IACUC: The Ethical Review Committee of the Fourth Hospital of Hebei Medical University granted approval for this study.
Consent: Each patient signed an informed consent form.
Inclusion and Exclusion Criteria
not detected.
Attrition
The study participants had an average age of 61.6+-8.0 (range, 47-73) years 25 patients were male and 15 were female.
Sex as a biological variable
The study participants had an average age of 61.6+-8.0 (range, 47-73) years 25 patients were male and 15 were female.
Subject Demographics
Age: not detected.
Weight: not detected.
Randomization
not detected.
Blinding
not detected.
Power Analysis
not detected.
Replication
Each reaction was performed in triplicate.

Number: Each reaction was performed in triplicate.
Cell Line Authentication
not detected.

Table 2: Key Resources Table

Your Sentences	REAGENT or RESOURCE	SOURCE	IDENTIFIER
Antibodies			
The membrane was incubated with primary antibodies for Bcl-2, Bcl-2-associated X protein (BAX), and glyceraldehyde 3-phosphate dehydrogenase (GAPDH 1:1,000 diluted in TBST).	Bcl-2-associated		
	glyceraldehyde 3-phosphate dehydrogenase (GAPDH 1:1,000 diluted in TBST)		
	GAPDH		
Experimental Models: Cell Lines			
Western blot After 48 h of transfection with microRNA oligos, Eca-109 cells were harvested.	Eca-109		Suggestion: RRID:CVCL_6898(link)
Recombinant DNA			
Finally, the DNA fragment was inserted into the pmiRGLO vector (Promega) with T4 ligase (Promega) to construct the recombinant luciferase vector (pmiRGLO-3-UTR).	pmiRGLO		Suggestion: RRID:Addgene_117339(link)
	pmiRGLO-3-UTR		
Software and Algorithms			
TargetScan (http:www.targetscan.org)	TargetScan		
), miRanda (http:www.targetscan.org), and PicTar (http:pictar.mdc-berlin.de) were employed for the prediction of miR-449a-5ps target gene.	PicTar		
Statistical analysis The SPSS 13.0 statistical software package was used to perform the data analyses in this study.	SPSS		Suggestion: SPSS, RRID:SCR_002865(link)

Other Entities Detected

Your Sentences	Recognized Entity
Statistical Tests	
Differences between two groups were analyzed using the non-parametric Mann-Whitney U test.	Mann-Whitney U test
Statistically significant differences between four groups were determined through one-way analysis of variance (ANOVA).	analysis of variance
	ANOVA

SciScore is an automated tool that is designed to assist expert reviewers by finding and presenting formulaic information scattered throughout a paper in a standard, easy to digest format. ***SciScore is not a substitute for expert review.*** SciScore also checks for the presence and correctness of several unique identifiers, including RRIDs (research resource identifiers) in the manuscript, detects sentences that appear to be missing RRIDs, and can even suggest RRIDs under certain circumstances. **All RRID suggestions should be verified;** only the author can know whether the suggestions are correct.

For a full description of scored criteria and tips for improving your score, please see https://www.scicrunch.com/sciscorereport-faq