Document Identifier: **PMC8802728**

SciScore Report

Below you will find your SciScore report containing three tables. Your score is calculated based on adherence to scientific rigor criteria (Table 1) and identification of key biological resources (Table 2). Table 3 contains statistical tests and oligonucleotides but is not scored. If SciScore makes any mistakes, please contact us to help us learn and improve.

Table 1: Rigor Adherence Table

Ethics
IRB: Ethical approval was obtained from the Institutional Review Board vide Letter No. JIP/IEC/ SC/2016/29/925 dated 15/07/2016 and informed written consent was taken from all participants.
Consent: Ethical approval was obtained from the Institutional Review Board vide Letter No. JIP/IEC/ SC/2016/29/925 dated 15/07/2016 and informed written consent was taken from all participants.
Inclusion and Exclusion Criteria
3] Patients aged more than 18 years with CAD who had undergone elective PCI for chronic stable angina or ACS with left ventricular ejection fraction (EF)[3] 50% and above were eligible for enrolment.
Attrition
not detected.
Sex as a biological variable
not detected.
Subject Demographics
Age: 3] Patients aged more than 18 years with CAD who had undergone elective PCI for chronic stable angina or ACS with left ventricular ejection fraction (EF)[3] 50% and above were eligible for enrolment.
Randomization
Materials and Methods A nonblinded randomized control trial was designed to assess the effect of a nurse-led CR program on behavioural, physiological and biochemical parameters control at 12 weeks intervals of the patients with CAD who had undergone elective coronary angioplasty at a tertiary care centre in South India.
Blinding
No blinding was used in this study for outcome assessment.
Power Analysis
The sample size was calculated by comparing the mean difference of total cholesterol (TC) between the Intervention group and control group as –0.50 (0.47) versus –0.17 (0.47) at 3-months' intervals of the group who had nurse-led CR and routine care, with the power of the study as 80% and at 5% level of significance.

Replication
not required.

Table 2: Key Resources Table

Your Sentences	REAGENT or RESOURCE	SOURCE	IDENTIFIER
Software and Algorithms			
The research nurse was trained on CR with American Heart Association (AHA) guidelines by the faculty from medical and nursing discipline.	American Heart Association		Suggestion: American Heart Association (RRID:SCR_007210)(link)
All data were analyzed using IBM SPSS version 22 (IBM SPSS Statistics for Windows, Version 22.0.	SPSS		Suggestion: SPSS, RRID:SCR_002865(link)

Other Entities Detected

Your Sentences	Recognized Entity
Statistical Tests	
Categorical data were summarized using frequency and percentage. Chi-square or Fisher's exact test was applied for the comparison of categorical data.	Chi-square
	Fisher's exact test
A comparison of the impact of counseling between control and intervention groups was done using independent Student's t-test or the alternative nonparametric tests like Mann–Whitney U test.	t-test
	Mann–Whitney U test
Independent student t-test, analysis of variance, and difference in difference (DiD) were used to evaluate the effectiveness of counseling on CR.	analysis of variance

SciScore is an automated tool that is designed to assist expert reviewers by finding and presenting formulaic information scattered throughout a paper in a standard, easy to digest format. ***SciScore is not a substitute for expert review.*** SciScore also checks for the presence and correctness of several unique identifiers, including RRIDs (research resource identifiers) in the manuscript, detects sentences that appear to be missing RRIDs, and can even suggest RRIDs under certain circumstances. **All RRID suggestions should be verified;** only the author can know whether the suggestions are correct.

For a full description of scored criteria and tips for improving your score, please see https://www.scicrunch.com/sciscorereport-faq