Document Identifier: **PMC8814304**

SciScore Report

Below you will find your SciScore report containing three tables. Your score is calculated based on adherence to scientific rigor criteria (Table 1) and identification of key biological resources (Table 2). Table 3 contains statistical tests and oligonucleotides but is not scored. If SciScore makes any mistakes, please contact us to help us learn and improve.

Table 1: Rigor Adherence Table

Ethics
Consent: Patients >=65 years of European descent presenting for elective surgery with an expected duration >60min were included after giving informed consent.
IRB: All procedures were approved by the local medical ethics committees of the study centers in Berlin, Germany (EA209214) and Utrecht, Netherlands (14-469).
Inclusion and Exclusion Criteria
Exclusion criteria were MMSE score <=23 points in the screening, neuropsychiatric morbidity and sensory impairment interfering with neurocognitive testing, centrally acting medication, homelessness or expected unavailability for a follow-up assessment, participation in another clinical intervention study during hospital stay, accommodation in an institution due to official or judicial order (Winterer et al., 2018).
Attrition
not detected.
Sex as a biological variable
(For illustrative purposes, regression coefficients were re-transformed to a natural scale for a female 72-year-old patient in BioCog and a female 59-year-old patient in PHYDELIO.
Subject Demographics
Age: Patients >=65 years of European descent presenting for elective surgery with an expected duration >60min were included after giving informed consent.
Randomization
2.1.2 PHYDELIO PHYDELIO (Perioperative PHYsostigmine prophylaxis for liver resection patients at risk for DELIrium and postOperative cognitive dysfunction) is a single center, double blinded, two-armed, randomized placebo-controlled phase IV trial: Patients were randomized either to receive physostigmine or placebo starting after induction of anesthesia for 24 h.
Blinding
2.1.2 PHYDELIO PHYDELIO (Perioperative PHYsostigmine prophylaxis for liver resection patients at risk for DELIrium and postOperative cognitive dysfunction) is a single center, double blinded, two-armed, randomized placebo-controlled phase IV trial: Patients were randomized either to receive physostigmine or placebo starting after induction of anesthesia for 24 h.

Power Analysis	
not detected.	
Replication	
The screening for delirium was initiated post-surgery and repeated twice per day at 8:00am and 7:00pm (+-1 h) up to seven days after surgery .	
Number: The screening for delirium was initiated post-surgery and repeated twice per day at 8:00am and 7:00pm (+-1 h) up to seven days after surgery .	
Cell Line Authentication	
Authentication: Since plasma IL-8 was not measured in PHYDELIO and thus validation was not possible, the prognostic value for plasma IL-8 has not been analyzed in this instance.	
Protocol Information	
Identifiers: The study was registered at clinicaltrials.gov (NCT02832193).	NCT02832193 Warning: study registration date June 28, 2016 is after study start date March 2009
Identifiers: The study is registered , number ISRCTN18978802 and EudraCT 2008-007237-47 .	2008-007237-47

Table 2: Key Resources Table

Your Sentences	REAGENT or RESOURCE	SOURCE	IDENTIFIER
Software and Algorithms			
Statistical analysis was conducted using the boot, Hmisc, lme4, ROCit and DescTools packages for R 4.0.	DescTools		
Model 5 and 6: Whole blood and plasma IL-8 levels from the first two assessment points of the BioCog study, adjusted for sex and age, surgery type, ASA-PS, anesthesia duration and hemoglobin levels.	BioCog		

Other Entities Detected

Your Sentences	Recognized Entity
Statistical Tests	
In order to compare IL-8 levels at three different sampling time points between patients with postoperative delirium and without delirium, the Wilcoxon rank sum test for independent samples was applied.	Wilcoxon rank sum test
Wilcoxons signed rank test was also used to test for statistically differences in IL-8 levels between patients with different perioperative medication (Supplementary Tables S6 and S7).	Wilcoxons signed rank test

SciScore is an automated tool that is designed to assist expert reviewers by finding and presenting formulaic information scattered throughout a paper in a standard, easy to digest format. ***SciScore is not a substitute for expert review.*** SciScore also checks for the presence and correctness of several unique identifiers, including RRIDs (research resource identifiers) in the manuscript, detects sentences that appear to be missing RRIDs, and can even suggest RRIDs under certain circumstances. **All RRID suggestions should be verified;** only the author can know whether the suggestions are correct.

For a full description of scored criteria and tips for improving your score, please see https://www.scicrunch.com/sciscorereport-faq