Document Identifier: **PMC8819026**

SciScore Report

Below you will find your SciScore report containing three tables. Your score is calculated based on adherence to scientific rigor criteria (Table 1) and identification of key biological resources (Table 2). Table 3 contains statistical tests and oligonucleotides but is not scored. If SciScore makes any mistakes, please contact us to help us learn and improve.

Table 1: Rigor Adherence Table

Ethics
not required.
Inclusion and Exclusion Criteria
not required.
Attrition
not required.
Sex as a biological variable
not detected.
Subject Demographics
Age: not required.
Weight: not required.
Randomization
not detected.
Blinding
not detected.
Power Analysis
not detected.
Replication
All assays were conducted in triplicate, and each experiment was repeated at least thrice.
Number: All assays were conducted in triplicate, and each experiment was repeated at least thrice.
Cell Line Authentication

not detected.

Table 2: Key Resources Table

Your Sentences	REAGENT or RESOURCE	SOURCE	IDENTIFIER
Antibodies			
After blocking with 3 bovine serum albumin, the membranes were incubated with primary antibodies at 4 degC overnight GRP78 (Cat: sc-13539 1:1000), PERK (Cat: sc-377400 1:1000), ATF4 (Cat: sc-390063 1:1000), eIF2a (Cat: sc-133132 1:1000), CHOP (Cat: sc-7351 1:1000), TFAM (Cat: sc-166965 1:1000), NRF1 (Cat: sc-515360 1:1000), b-Actin (Cat: sc-47778 1:1000), (Santa Cruz Biotechnology, Inc, USA), Bax (Cat: 5023 1:1000), Bcl2 (Cat: 3498 1:1000), Caspase 3 (Cat: 9662 1:1000) (CST, USA), Complex I (Cat: (ab110245) 1:1000), Complex II (Cat: (ab110410) 1:1000) and ATP synthase C (Cat: ab181243 1:1000) (Abcam, USA), p-PERK (Cat: ITP07576 1:1000), Mfn-2 (Cat: ITT2740 1:1000), p-eIF2a (Cat: ITP03087 1:1000), (G-biosciences, USA) at 1:1000 dilution in TBST.	PERK		
	ATF4		
	eIF2a		Suggestion: (LSBio (LifeSpan) Cat# LS-C123503-50, RRID:AB_10802395)(link)
	TFAM		
	NRF1		Suggestion: (LSBio (LifeSpan) Cat# LS-C110287-50, RRID:AB_10661254)(link)
	b-Actin		
	Bcl2		
	ab110410		
	p-PERK		
	p-eIF2a		
Then cells were blocked with 3 BSA solution and incubated with PDI (protein disulfide isomerase an ER resident enzyme) and primary antibody GRP78 (Santa Cruz Biotechnology), (1:200) or CHOP (1:50) in 3 BSA (Bovine serum albumin) at 4 degC temperature for 12 h.	CHOP		
Followed by incubation with secondary anti-rat antibody conjugated with rhodamine (Santa Cruz Biotechnology Inc., CA, USA) or FITC (Fluorescein isothiocyanate) Sigma) for 2 h in dark at room temperature.	anti-rat		
Later on cells were incubated with primary antibody solution (1:100) of anti-Mfn2 antibody (rabbit polyclonal, G-biosciences) in PBS.	anti-Mfn2		
After 12 h of incubation cells were washed thrice with PBS, and then subjected to FITC conjugated anti-rabbit secondary antibody solution (1:100) incubation for 2 h in the dark.	anti-rabbit		
Experimental Models: Cell Lines			

2.2 Methods 2.2.1 Experimental design Mouse N2A neuroblastoma cells obtained from NCCS, Pune were cultured in Minimum Essential media (MEM containing 5.5 mM glucose) supplemented with 10 FBS, streptomycinpenicillin (1), glutamine (2 mM) and grown at 37 degC, in a humidified atmosphere of 95 air and 5 CO2.	N2A		
2.2.3 ROS estimation by 2,7-dichlorofluorescin diacetate (DCFDA) staining Cellular Reactive Oxygen Species (ROS) was measured in Neuro2a cells seeded in 6-well plate at a density of 5000 cellswell.	Neuro2a		Suggestion: (RCB Cat# RCB5511, RRID:CVCL_D3XR)(link)
Software and Algorithms			
2 Materials and methods 2.1 Materials GSK2606414 and most of the other chemicals are reagent grade and obtained from Sigma Aldrich, USA.	Sigma Aldrich		Suggestion: Sigma-Aldrich (RRID:SCR_008988)(link)
After blocking with 3 bovine serum albumin, the membranes were incubated with primary antibodies at 4 degC overnight GRP78 (Cat: sc-13539 1:1000), PERK (Cat: sc-377400 1:1000), ATF4 (Cat: sc-390063 1:1000), eIF2a (Cat: sc-133132 1:1000), CHOP (Cat: sc-7351 1:1000), TFAM (Cat: sc-166965 1:1000), NRF1 (Cat: sc-515360 1:1000), b-Actin (Cat: sc-47778 1:1000), (Santa Cruz Biotechnology, Inc, USA), Bax (Cat: 5023 1:1000), Bcl2 (Cat: 3498 1:1000), Caspase 3 (Cat: 9662 1:1000) (CST, USA), Complex I (Cat: (ab110245) 1:1000), Complex II (Cat: (ab110410) 1:1000) and ATP synthase C (Cat: ab181243 1:1000) (Abcam, USA), p-PERK (Cat: ITP07576 1:1000), Mfn-2 (Cat: ITT2740 1:1000), p-eIF2a (Cat: ITP03087 1:1000), (G-biosciences, USA) at 1:1000 dilution in TBST.	Cat		Suggestion: CAT (RRID:SCR_008421)(link)
Chemiluminescence signal was captured using a Fusion-FX imager (VilberLourmat, Germany) and relative band intensities were quantified by densitometry using Image-J software (version 1.48, NIH, USA) (Arruri et al., 2017).	Image-J		Suggestion: ImageJ, RRID:SCR_003070(link)
The intergroup variation was measured by one-way analysis of	Graph Pad Prism		

variance (ANOVA) followed by Bonferronis multiple comparison post-hoc test using the Graph Pad Prism (version-5.0).			

Other Entities Detected

Your Sentences	Recognized Entity
Statistical Tests	
The intergroup variation was measured by one-way analysis of variance (ANOVA) followed by Bonferronis multiple comparison post-hoc test using the Graph Pad Prism (version-5.0).	analysis of variance
	ANOVA
	Bonferronis multiple comparison post-hoc test

SciScore is an automated tool that is designed to assist expert reviewers by finding and presenting formulaic information scattered throughout a paper in a standard, easy to digest format. ***SciScore is not a substitute for expert review.*** SciScore also checks for the presence and correctness of several unique identifiers, including RRIDs (research resource identifiers) in the manuscript, detects sentences that appear to be missing RRIDs, and can even suggest RRIDs under certain circumstances. **All RRID suggestions should be verified;** only the author can know whether the suggestions are correct.

For a full description of scored criteria and tips for improving your score, please see https://www.scicrunch.com/sciscorereport-faq