Document Identifier: **PMC8827068**

SciScore Report

Below you will find your SciScore report containing three tables. Your score is calculated based on adherence to scientific rigor criteria (Table 1) and identification of key biological resources (Table 2). Table 3 contains statistical tests and oligonucleotides but is not scored. If SciScore makes any mistakes, please contact us to help us learn and improve.

Table 1: Rigor Adherence Table

Ethics
IRB: The study was approved by the Ethics Review Board, Uppsala (2018159).
Consent: Exclusion criteria were: celiac disease, functional dyspepsia, Helicobacter pylori infection during the preceding 6 mo, or other functional or inflammatory gastrointestinal disease, previous or ongoing cancer treatment, previous bariatric or abdominal surgery other than appendectomy, treatment of weight reduction, >10 kg body weight change in the preceding year, refusal to give informed consent, unstable medication from 14 d prior to inclusion or during the study, concurrent probiotic or antibiotic medication, reluctance to consume rice porridge daily during 3 separate weeks, pregnancy or lactation, blood donation or participation in other intervention trials within 30 d prior to screening or any time during the study, history of drug or alcohol abuse, smoking, and inability to understand the Swedish language.
Field Sample Permit: Procedures for the collection of blood, urine, and feces samples are described in detail in Supplementary Methods 2 .
Inclusion and Exclusion Criteria
Participants Inclusion criteria of the study were: female and male with moderate to severe IBS (IBS-SSS score >175) (28), BMI 18.5-38 kgm2, age 18-70 y, hemoglobin 120-160 gL, thyroid-stimulating hormone <4 mUL, C-reactive protein <5 mgL, transglutaminase immunoglobulin A <7 UmL, and systolicdiastolic blood pressure <=160<=105 mmHg.
Attrition
Two individuals were mistakenly assigned the sequence ABC .
Sex as a biological variable
Participants Inclusion criteria of the study were: female and male with moderate to severe IBS (IBS-SSS score >175) (28), BMI 18.5-38 kgm2, age 18-70 y, hemoglobin 120-160 gL, thyroid-stimulating hormone <4 mUL, C-reactive protein <5 mgL, transglutaminase immunoglobulin A <7 UmL, and systolicdiastolic blood pressure <=160<=105 mmHg.
Subject Demographics
Weight: Participants Inclusion criteria of the study were: female and male with moderate to severe IBS (IBS-SSS score >175) (28), BMI 18.5-38 kgm2, age 18-70 y, hemoglobin 120-160 gL, thyroid-stimulating hormone <4 mUL, C-reactive protein <5 mgL, transglutaminase immunoglobulin A <7 UmL, and systolicdiastolic blood pressure <=160<=105 mmHg.

Age: Participants Inclusion criteria of the study were: female and male with moderate to severe IBS (IBS-SSS score >175) (28), BMI 18.5-38 kgm2, age 18-70 y, hemoglobin 120-160 gL, thyroid-stimulating hormone <4 mUL, C-reactive protein <5 mgL, transglutaminase immunoglobulin A <7 UmL, and systolicdiastolic blood pressure <=160<=105 mmHg.
Randomization
Methods Study design The double-blind, placebo-controlled, randomized 3-way study with triple crossover design was conducted from September 2018 to June 2019 in Uppsala, Sweden.
Blinding
Participants were then prompted to consume blinded food in the form of rice porridge with added FODMAPs, gluten, or placebo during weeks 3, 5, and 7, but no porridge during weeks 4 and 6 (Figure 1).
Power Analysis
Randomization was restricted to three treatment orders , to ensure sufficient statistical power to allow differentiation of a potential effect of sequencing .
Replication
not required.
Protocol Information

Identifiers: The trial was registered at www.clinicaltrials.gov as NCT03653689.	NCT03653689

Table 2: Key Resources Table

Your Sentences	REAGENT or RESOURCE	SOURCE	IDENTIFIER

Other Entities Detected

Your Sentences	Recognized Entity
Statistical Tests	
Under these conditions, the required number of participants was 64, with the level of significance in a 2-sided test set to 0.053 (Bonferroni correction).	Bonferroni
Comparisons between subtypes at screening were conducted through 1-factor ANOVA.	ANOVA
Proportion tests were performed using McNemars test.	McNemars test

SciScore is an automated tool that is designed to assist expert reviewers by finding and presenting formulaic information scattered throughout a paper in a standard, easy to digest format. ***SciScore is not a substitute for expert review.*** SciScore also checks for the presence and correctness of several unique identifiers, including RRIDs (research resource identifiers) in the manuscript, detects sentences that appear to be missing RRIDs, and can even suggest RRIDs under certain circumstances. **All RRID suggestions should be verified;** only the author can know whether the suggestions are correct.

For a full description of scored criteria and tips for improving your score, please see https://www.scicrunch.com/sciscorereport-faq