Document Identifier: **PMC8850559**

SciScore Report

Below you will find your SciScore report containing three tables. Your score is calculated based on adherence to scientific rigor criteria (Table 1) and identification of key biological resources (Table 2). Table 3 contains statistical tests and oligonucleotides but is not scored. If SciScore makes any mistakes, please contact us to help us learn and improve.

Table 1: Rigor Adherence Table

Ethics
IRB: 2.2 Preclinical study This experiment was based on the ethics requirements of the National Council for Animal Experimentation Control (CONCEA) and was approved by the Ethics Committee on Animal Use (CEUAUNICAMP protocol 4895-12018).
Field Sample Permit: The access to Brazilian genetic heritage was registered at the Ministry of the Environment via SISGEN (protocol A2C81F8).
Euthanasia Agents: Blood was collected by cardiac puncture under anesthesia (ketamine and xylazine) the liver and the sum of adipose tissue (mesenteric , retroperitoneal , and epididymal) were weighted , and liver fragments were immediately frozen in liquid nitrogen .
Inclusion and Exclusion Criteria
not detected.
Attrition
not detected.
Sex as a biological variable
Twenty-four male black mice (C57BL6), 5 weeks old, provided by the Multidisciplinary Center of Biological Investigation (CEMIBUNICAMP), were allocated collectively in polyethylene boxes, under environmental controlled conditions of temperature (22+-2 degC), humidity (60-70), a light-dark cycle of 12 h, with free access to water and food during the whole experimental period.
Subject Demographics
Age: Twenty-four male black mice (C57BL6), 5 weeks old, provided by the Multidisciplinary Center of Biological Investigation (CEMIBUNICAMP), were allocated collectively in polyethylene boxes, under environmental controlled conditions of temperature (22+-2 degC), humidity (60-70), a light-dark cycle of 12 h, with free access to water and food during the whole experimental period.
Randomization
not detected.
Blinding
not detected.

Power Analysis
not detected.
Replication
not required.

Table 2: Key Resources Table

Your Sentences	REAGENT or RESOURCE	SOURCE	IDENTIFIER
Antibodies			
The membranes were blocked within none to 3 bovine serum albumin , incubated with specific antibodies overnight 4 degC mouse anti-TLR4 (sc-53462 , Santa Cruz Biotechnology) rabbit anti-MyD88 (bs-1047R , Bioss)	anti-TLR4		Suggestion: (IMGENEX Cat# IMG-500-TLR4, RRID:AB_10598785)(link)
	sc-53462		
	anti-MyD88		
	bs-1047R		
) mouse anti-nuclear factor-kappa B (NFkB) (ab13594, Abcam) mouse anti-tumor necrosis factor-alpha (TNF-a) (ab48394, Abcam) goat anti-phospho-insulin receptor substrate (IRS) tyrosine (sc-17196, Santa Cruz Biotechnology) and mouse anti-b-actin (ab8227, Abcam), then incubated with specific secondary antibodies for 2 h, at room temperature.	anti-nuclear factor-kappa B (NFkB		
	anti-tumor necrosis factor-alpha (TNF-a		Suggestion: (US Biological Cat# T9160-21P, RRID:AB_2256063)(link)
	anti-phospho-insulin receptor substrate (IRS) tyrosine (sc-17196, Santa Cruz Biotechnology)		
	anti-b-actin		
Experimental Models: Organisms/Strains			
Twenty-four male black mice (C57BL6), 5 weeks old, provided by the Multidisciplinary Center of Biological Investigation (CEMIBUNICAMP), were allocated collectively in polyethylene boxes, under environmental controlled conditions of temperature (22+-2 degC), humidity (60-70), a light-dark cycle of 12 h, with free access to water and food during the whole experimental period.	C57BL6		Suggestion: RRID:IMSR_JAX:000664(link)
Software and Algorithms			
Biochemical analyses (glycemia , insulin , and LPS levels) were measured in the serum using commercial kits according to the manufactures (LaborLab(r) , Millipore(c) , and Thermo scientific(tm)) .	Thermo scientific(tm)		
Alpha (Shannon diversity) and beta (Detrended Correspondence Analysis) diversity analyses were performed on	phyloseq		

the phyloseq package as described (Callahan et al., 2016).			
The taxonomic counts contained in the phyloseq were imported into the edgeR package for normalization of the library sizes of each sample subsequently, the counts were transformed to the base 2 logarithms of the counts per million (logCPM) of each sample (voom transformation).	edgeR		

Other Entities Detected

Your Sentences	Recognized Entity
Statistical Tests	
After adjusting the linear limma model, differential taxonomic abundance was tested for each group with moderate t-tests (Smyth, 2004).	t-tests
The data were submitted to analysis of variance (ANOVA) with post-hoc on Tukeys test with a p-value <0.05 within the experimental groups.	analysis of variance
	ANOVA
	Tukeys test
For metataxonomic analysis, a moderated t-test (p-value <0.05) was used to compare the profile of gut bacteria of every single diet distinctly.	t-test
The Pearson correlation test was carried out comparing the better and the worst phenotypes for glucose metabolism (CJ and HF respectively), considering p-value <0.05 as a significant difference.	Pearson correlation test

SciScore is an automated tool that is designed to assist expert reviewers by finding and presenting formulaic information scattered throughout a paper in a standard, easy to digest format. ***SciScore is not a substitute for expert review.*** SciScore also checks for the presence and correctness of several unique identifiers, including RRIDs (research resource identifiers) in the manuscript, detects sentences that appear to be missing RRIDs, and can even suggest RRIDs under certain circumstances. **All RRID suggestions should be verified;** only the author can know whether the suggestions are correct.

For a full description of scored criteria and tips for improving your score, please see https://www.scicrunch.com/sciscorereport-faq