Document Identifier: **PMC8870008**

SciScore Report

Below you will find your SciScore report containing three tables. Your score is calculated based on adherence to scientific rigor criteria (Table 1) and identification of key biological resources (Table 2). Table 3 contains statistical tests and oligonucleotides but is not scored. If SciScore makes any mistakes, please contact us to help us learn and improve.

Table 1: Rigor Adherence Table

Ethics
not detected.
Inclusion and Exclusion Criteria
not required.
Attrition
not required.
Sex as a biological variable
Materials and Methods Mice Study Twelve-week-old male C57BL6J mice were fed a modified Lieber-DeCarli (LD) liquid diet containing ethanol (ethanol group E1-3, n = 7-10) or the control LD diet in which ethanol was substituted isocalorically by maltose-dextrin (control group C1-3, n = 5-7) for 5 weeks.
Subject Demographics
Age: Materials and Methods Mice Study Twelve-week-old male C57BL6J mice were fed a modified Lieber-DeCarli (LD) liquid diet containing ethanol (ethanol group E1-3, n = 7-10) or the control LD diet in which ethanol was substituted isocalorically by maltose-dextrin (control group C1-3, n = 5-7) for 5 weeks.
Randomization
Male C57BL6J mice (10-12 weeks old) were randomly divided into seven groups.
Blinding
Histopathology A section of the liver from each mouse was processed for hematoxylin and eosin staining using standard procedures and examined by an experienced pathologist blinded to the study.
Power Analysis
not detected.
Replication
not required.

Table 2: Key Resources Table

Your Sentences	REAGENT or RESOURCE	SOURCE	IDENTIFIER
Experimental Models: Organisms/Strains			
Male C57BL6J mice (10-12 weeks old) were randomly divided into seven groups.	C57BL6J		Suggestion: RRID:IMSR_JAX:000664(link)

Other Entities Detected

Your Sentences	Recognized Entity
Statistical Tests	
S9), and after adjusting the P value using the Benjamini-Hochberg correction, no differences between ethanol-fed and control mice were observed (false discovery rate FDR corrected $P < 0.05$) however, calorie intake in weeks 3-5 was slightly higher in control-fed mice and approaching significance.	Benjamini-Hochberg
Score averages, SDs, and P values were determined using the two-sided, unpaired, or unequal variance Student t test.	t test

SciScore is an automated tool that is designed to assist expert reviewers by finding and presenting formulaic information scattered throughout a paper in a standard, easy to digest format. ***SciScore is not a substitute for expert review.*** SciScore also checks for the presence and correctness of several unique identifiers, including RRIDs (research resource identifiers) in the manuscript, detects sentences that appear to be missing RRIDs, and can even suggest RRIDs under certain circumstances. **All RRID suggestions should be verified;** only the author can know whether the suggestions are correct.

For a full description of scored criteria and tips for improving your score, please see https://www.scicrunch.com/sciscorereport-faq