Document Identifier: **PMC8874047**

SciScore Report

Below you will find your SciScore report containing three tables. Your score is calculated based on adherence to scientific rigor criteria (Table 1) and identification of key biological resources (Table 2). Table 3 contains statistical tests and oligonucleotides but is not scored. If SciScore makes any mistakes, please contact us to help us learn and improve.

Table 1: Rigor Adherence Table

Ethics
not detected.
Inclusion and Exclusion Criteria
not required.
Attrition
not required.
Sex as a biological variable
2.5 Wound healing in an animal model 2.5.1 Animals Eight-week-old male Wistar rats (200-230 g) (Liaoning Changsheng Biotechnology , China) were used in a diabetic wound healing animal model and 12-week-old male Wistar rats (400-500 g) for normal wound healing .
Subject Demographics
Age: 2.5 Wound healing in an animal model 2.5.1 Animals Eight-week-old male Wistar rats (200-230 g) (Liaoning Changsheng Biotechnology , China) were used in a diabetic wound healing animal model and 12-week-old male Wistar rats (400-500 g) for normal wound healing .
Weight: 2.5 Wound healing in an animal model 2.5.1 Animals Eight-week-old male Wistar rats (200-230 g) (Liaoning Changsheng Biotechnology , China) were used in a diabetic wound healing animal model and 12-week-old male Wistar rats (400-500 g) for normal wound healing .
Randomization
The normal and diabetic rats were randomly allocated into one of three groups , with 5 normal rats and 10 diabetic rats in each group: (1) the control group , in which wounds healed spontaneously without treatment (2) hot-air treatment group , in which an air heater was used to treat the wounds (3) LTP-treated group , in which a new type of LTP device was used to treat the wounds .
Blinding
not detected.
Power Analysis
not detected.

Replication
not required.

Table 2: Key Resources Table

Your Sentences	REAGENT or RESOURCE	SOURCE	IDENTIFIER
Antibodies			
Slides were stained for antibodies against CD31, CK14, PCNA, and transforming growth factor-beta (TGF-b), the details of which are displayed in Table S1.	CD31		
	CK14		
Experimental Models: Organisms/Strains			
Therefore , a nitritenitrate colorimetric assay kit (Beyotime) was used to measure the total quantity of NO2 - and NO3 - .	NO3 -		
2.5 Wound healing in an animal model 2.5.1 Animals Eight-week-old male Wistar rats (200-230 g) (Liaoning Changsheng Biotechnology , China) were used in a diabetic wound healing animal model and 12-week-old male Wistar rats (400-500 g) for normal wound healing .	Wistar		Suggestion: RRID:RGD_13508588(link)
Software and Algorithms			
Data were obtained and analysed using AvaSoft 7.8 spectroscopy software.	AvaSoft		
ImageJ software was used to count the number of lymphocytes and fibroblasts in the H&E stained images (400 x) .	ImageJ		Suggestion: ImageJ, RRID:SCR_003070(link)

Other Entities Detected

Your Sentences	Recognized Entity
Statistical Tests	
Statistical analyses were performed with one-way analysis of variance (ANOVA) followed by post-hoc t-test.	analysis of variance
	ANOVA
	t-test

SciScore is an automated tool that is designed to assist expert reviewers by finding and presenting formulaic information scattered throughout a paper in a standard, easy to digest format. ***SciScore is not a substitute for expert review.*** SciScore also checks for the presence and correctness of several unique identifiers, including RRIDs (research resource identifiers) in the manuscript, detects sentences that appear to be missing RRIDs, and can even suggest RRIDs under certain circumstances. **All RRID suggestions should be verified;** only the author can know whether the suggestions are correct.

For a full description of scored criteria and tips for improving your score, please see https://www.scicrunch.com/sciscorereport-faq