Document Identifier: **PMC8902179**

SciScore Report

Below you will find your SciScore report containing three tables. Your score is calculated based on adherence to scientific rigor criteria (Table 1) and identification of key biological resources (Table 2). Table 3 contains statistical tests and oligonucleotides but is not scored. If SciScore makes any mistakes, please contact us to help us learn and improve.

Table 1: Rigor Adherence Table

Ethics
IRB: The study was approved by the Research Ethics Boards at all participating hospitals.
Inclusion and Exclusion Criteria
Setting and Participants Participants included adults aged >=18 years who were receiving in-center hemodialysis 3 times per week at 4 tertiary-care hospitals in Montreal, Canada (Jewish General Hospital JGH, Centre Hospitalier de lUniversite de Montreal CHUM, Montreal General Hospital, and Lachine Hospital), and had symptoms of depression andor anxiety as indicated by a score >=6 on the Patient Health Questionnaire-9 (PHQ-9) andor the General Anxiety Disorder-7 (GAD-7).
Attrition
4 Exclusion criteria included significant cognitive impairment suggestive of dementia, suicidal ideation or intent, signs of acute psychosis, difficulty hearing, current psychotherapy, and inability to speak French or English. 4 Of the 400 potential participants at the 4 dialysis units, 112 participants were screened using the PHQ-9 and the GAD-7, and 64 met eligibility criteria.
Sex as a biological variable
not detected.
Subject Demographics
Age: Setting and Participants Participants included adults aged >=18 years who were receiving in-center hemodialysis 3 times per week at 4 tertiary-care hospitals in Montreal, Canada (Jewish General Hospital JGH, Centre Hospitalier de lUniversite de Montreal CHUM, Montreal General Hospital, and Lachine Hospital), and had symptoms of depression andor anxiety as indicated by a score >=6 on the Patient Health Questionnaire-9 (PHQ-9) andor the General Anxiety Disorder-7 (GAD-7).
Randomization
Of the eligible participants, 9 refused and 55 were randomized.
Blinding
The independent statistician transmitted participant group information directly to the interventionists, and the assessors were blinded to the participants group assignment.
Power Analysis

not detected.	
Replication	
not required.	
Data Information	
Identifiers: 4,12,13 The structure of HEP was equivalent to BMI: HEP was comprised of two 20-minute sessions per week for 8 weeks while patients were undergoing dialysis, encouraged the same amount of home practice (implementing health-enhancing habits for 10 minutes per day), allowed 3 to 5 minutes for questions, and was delivered via audio headsets with the interventionist present.	4,12,13
Protocol Information	
Identifiers: 4 Ethical Considerations Prior to recruitment, the trial had been registered (ClinicalTrials.gov Identifier: NCT03406845).	NCT03406845

Table 2: Key Resources Table

Your Sentences	REAGENT or RESOURCE	SOURCE	IDENTIFIER
Software and Algorithms			
Statistical analyses were performed using R (Version 3.5, R Foundation for Statistical Computing, Vienna, Austria) and IBM SPSS Statistics software (Version 27, SPSS, Inc., Chicago, Illinois).	SPSS		Suggestion: SPSS, RRID:SCR_002865(link)

Other Entities Detected

Your Sentences	Recognized Entity
Statistical Tests	
Data Analysis Baseline demographic and clinical characteristics were compared between the treatment and control groups using chi-square tests to assess adequacy of randomization among participants who completed the 6-month follow-up.	chi-square tests
To determine whether BMI is more efficacious than HEP at reducing PHQ-9 depression scores (primary outcome) and GAD-7 anxiety scores (secondary outcome) over the 6-month follow-up period, 2-way mixed analyses of variance 17,18 (ANOVAs) were conducted on 6-month follow-up PHQ-9 and GAD-7 scores separately, with the treatment group (BMI and HEP) as the between-subject factor and time (baseline, 8 weeks, and 6 months) as the within-subject factor.	ANOVAs
Normality of PHQ-9 and GAD-7 scores was assessed using the Kolmogorov-Smirnov test.	Kolmogorov-Smirnov test
19 Post hoc t tests were used to compare treatment groups at each of the 3 time points for any significant interaction effects between time and treatment group, using a Bonferroni correction.	t tests
	Bonferroni
Effect sizes were interpreted as follows: small effect = 0.2, medium effect = 0.5, and large effect = 0.8. 22 To compare the survival function of each group (exploratory outcome ie, to see whether one group was more likely to be hospitalized than the other), Kaplan-Meier (KM) estimates 23 and log-rank tests were performed to examine whether the survival curves for each group were identical or not.	log-rank tests

SciScore is an automated tool that is designed to assist expert reviewers by finding and presenting formulaic information scattered throughout a paper in a standard, easy to digest format. ***SciScore is not a substitute for expert review.*** SciScore also checks for the presence and correctness of several unique identifiers, including RRIDs (research resource identifiers) in the manuscript, detects sentences that appear to be missing RRIDs, and can even suggest RRIDs under certain circumstances. **All RRID suggestions should be verified;** only the author can know whether the suggestions are correct.

For a full description of scored criteria and tips for improving your score, please see https://www.scicrunch.com/sciscorereport-faq