Document Identifier: **PMC8905020**

SciScore Report

Below you will find your SciScore report containing three tables. Your score is calculated based on adherence to scientific rigor criteria (Table 1) and identification of key biological resources (Table 2). Table 3 contains statistical tests and oligonucleotides but is not scored. If SciScore makes any mistakes, please contact us to help us learn and improve.

Table 1: Rigor Adherence Table

Ethics
Field Sample Permit: Animal studies Sixty-six male Sprague-Dawley rats (270-310 g) were maintained in a room at 22 +- 1 degC under a 12-h light:12-h dark cycle in the Institute of Radiation Medicine , Chinese Academy of Medical Sciences (Tianjin , China) .
Euthanasia Agents: All animals were euthanized by barbiturate overdose (intravenous injection , 150 mgkg pentobarbital sodium) for tissue collection .
IACUC: All procedures involving animals were reviewed and approved by the Institutional Animal Care and Use Committee of the Institute of Radiation Medicine , Chinese Academy of Medical Sciences (IACUC Protocol No. IRM-DWLL-2021142) .
Inclusion and Exclusion Criteria
not detected.
Attrition
Four rats were randomly chosen from the PI-IBS and control group to perform the intestinal motility test with carbon solution gavage.
Sex as a biological variable
Animal studies Sixty-six male Sprague-Dawley rats (270-310 g) were maintained in a room at 22 +- 1 degC under a 12-h light:12-h dark cycle in the Institute of Radiation Medicine , Chinese Academy of Medical Sciences (Tianjin , China) .
Subject Demographics
Weight: Animal studies Sixty-six male Sprague-Dawley rats (270-310 g) were maintained in a room at 22 +- 1 degC under a 12-h light:12-h dark cycle in the Institute of Radiation Medicine , Chinese Academy of Medical Sciences (Tianjin , China) .
Randomization
Intervention and evaluation of a rat model of PI-IBS Rats were randomly divided into a PI-IBS group (n = 52) and control group (n = 14) .
Blinding
not detected.

Power Analysis
not detected.
Replication
not required.
Cell Line Authentication
not detected.

Table 2: Key Resources Table

Your Sentences	REAGENT or RESOURCE	SOURCE	IDENTIFIER
Antibodies			
The membranes were blocked in 5 skimmed milk powder at room temperature for 1 h , and incubated with primary rabbit polyclonal antibodies (SERT , Cat. No. ab102048 , Abcam , Cambridge , UK b-actin	SERT		
Experimental Models: Cell Lines			
Cell culture Caco-2 , a human epithelial colorectal adenocarcinoma cell line , was grown in Eagles minimal essential medium (MEM Gibco , New York , NY , United States) supplemented with 20 fetal bovine serum (FBS Gibco) and 1 nonessential amino acids at 37 degC .	Caco-2		Suggestion: RRID:CVCL_0025(link)
The human colonic epithelial carcinoma cell line , HT-29 , was grown in Dulbeccos modified Eagles medium (DMEM Gibco , New York , NY , United States) supplemented with 10 FBS and 1 nonessential amino acids at 37 degC .	HT-29		Suggestion: RRID:CVCL_0320(link)
Experimental Models: Organisms/Strains			
Animal studies Sixty-six male Sprague-Dawley rats (270-310 g) were maintained in a room at 22 +- 1 degC under a 12-h light:12-h dark cycle in the Institute of Radiation Medicine , Chinese Academy of Medical Sciences (Tianjin , China) .	Sprague-Dawley		Suggestion: RRID:RGD_70508(link)
Software and Algorithms			
cDNA was synthesized using the iScript cDNA synthesis kit (Bio-Rad Laboratories , Inc . , Hercules , CA , United States) .	Bio-Rad Laboratories		
The Bio-Rad Image Analysis System (Bio-Rad Laboratories) was utilized to visualize the protein bands.	Bio-Rad Image		
The protein quantitative analysis was undertaken using ImageJ software.	ImageJ		Suggestion: ImageJ, RRID:SCR_003070(link)
All statistical analyses were performed using SPSS 20.0 software (IBM, Armonk, NY, United States).	SPSS		Suggestion: SPSS, RRID:SCR_002865(link)

Other Entities Detected

Your Sentences	Recognized Entity
Statistical Tests	
Statistical analysis The Students t-test was used to assess differences between two groups.	t-test
Oligonucleotides	
Table 1 Primer sequences for RT-PCR Gene Sequence (5-3) rGAPDH Forward: 5-CCATCAACGACCCCTTCATT-3 Reverse: 5-GACCAGCTTCCCATTCTCAG-3 rSERT Forward: 5-ACTGTTACCAAGATGCCCTG-3 Reverse: 5-ATCTTCATTCCTCATCTCCGC- 3 hGAPDH Forward: 5-ACA GCA ACT CCC ATT CTT-3 Reverse: 5-TCC AGG GTT TCT TAC TCC-3 hSERT Reverse: 5-AAT	5-CCATCAACGACCCCTTCATT-3
	5-GACCAGCTTCCCATTCTCAG-3
	5-ACTGTTACCAAGATGCCCTG-3
	5-ATCTTCATTCCTCATCTCCGC- 3
	5-ACA GCA ACT CCC ATT CTT-3
	5-TCC AGG GTT TCT TAC TCC-3
	5-AAT
GGG TAC TCA GCA GTT CC-3 Reverse: 5-CCA CAG CAT AGC CAA TCA C-3 hGAPDH: Human glyceraldehyde-3-phosphate dehydrogenase hSERT: Human serotonin transporter rGAPDH: Rat glyceraldehyde-3-phosphate dehydrogenase rSERT: Rat serotonin transporter .	5-CCA CAG CAT AGC CAA TCA C-3

SciScore is an automated tool that is designed to assist expert reviewers by finding and presenting formulaic information scattered throughout a paper in a standard, easy to digest format. ***SciScore is not a substitute for expert review.*** SciScore also checks for the presence and correctness of several unique identifiers, including RRIDs (research resource identifiers) in the manuscript, detects sentences that appear to be missing RRIDs, and can even suggest RRIDs under certain circumstances. **All RRID suggestions should be verified;** only the author can know whether the suggestions are correct.

For a full description of scored criteria and tips for improving your score, please see https://www.scicrunch.com/sciscorereport-faq