Document Identifier: **PMC8914001**

SciScore Report

Below you will find your SciScore report containing three tables. Your score is calculated based on adherence to scientific rigor criteria (Table 1) and identification of key biological resources (Table 2). Table 3 contains statistical tests and oligonucleotides but is not scored. If SciScore makes any mistakes, please contact us to help us learn and improve.

Table 1: Rigor Adherence Table

Ethics
IRB: MATERIALS AND METHODS All procedures were performed according to the guidelines for the care and use of research animals and were approved by the Third Local Ethics Committee on Animal Experimentation in Warsaw (SGGW Warsaw) (resolution number 29/2010, Warsaw, PL).
Euthanasia Agents: The chickens were slaughtered by decapitation after electrical stunning.
Inclusion and Exclusion Criteria
not detected.
Attrition
not detected.
Sex as a biological variable
Chickens, Breeding, and Bone Sample Collection In this study, fast-growing Cobb 500 (FG) chickens and 2 medium-growing chickens MGH (HubbardJA957) and MGGP (only males) were raised up to 56 D of age.
Subject Demographics
Age: For the Bvol and MvBMD measurements of the bones from 14 to 56 day-old chickens, the VOI was limited by the minimum and maximum densities of the investigated bones at 0 and 3,071 HU, respectively.
Randomization
In each of these growth stages, 10 chickens from each genotype were randomly selected for slaughter.
Blinding
not detected.
Power Analysis
not detected.
Replication
Ten replications consisting of 36 chickens kept in the same pen, with an area of 3 m2, were formed within each genotype.

Number: Ten replications consisting of 36 chickens kept in the same pen, with an area of 3 m2, were formed within each genotype.
Cell Line Authentication
not detected.

Table 2: Key Resources Table

Your Sentences	REAGENT or RESOURCE	SOURCE	IDENTIFIER
Experimental Models: Cell Lines			
Chickens, Breeding, and Bone Sample Collection In this study, fast-growing Cobb 500 (FG) chickens and 2 medium-growing chickens MGH (HubbardJA957) and MGGP (only males) were raised up to 56 D of age.	HubbardJA957		
Software and Algorithms			
All calculations were performed using Statistica 10.0 software (Statistica, 2011) and SPSS 23 (IBM, 2015).	Statistica		Suggestion: STATISTICA (RRID:SCR_014213)(link)
	SPSS		Suggestion: SPSS, RRID:SCR_002865(link)

Other Entities Detected

Your Sentences	Recognized Entity
Statistical Tests	
All examined variables were compared between chickens of different genetic groups within the same slaughter stage and within the genetic group between different weeks of rearing by 1-way analysis of variance (ANOVA) and multiple comparison post-hoc Duncan test.	analysis of variance
	ANOVA
	Duncan test

SciScore is an automated tool that is designed to assist expert reviewers by finding and presenting formulaic information scattered throughout a paper in a standard, easy to digest format. ***SciScore is not a substitute for expert review.*** SciScore also checks for the presence and correctness of several unique identifiers, including RRIDs (research resource identifiers) in the manuscript, detects sentences that appear to be missing RRIDs, and can even suggest RRIDs under certain circumstances. **All RRID suggestions should be verified;** only the author can know whether the suggestions are correct.

For a full description of scored criteria and tips for improving your score, please see https://www.scicrunch.com/sciscorereport-faq