Document Identifier: **PMC8922213**

SciScore Report

Below you will find your SciScore report containing three tables. Your score is calculated based on adherence to scientific rigor criteria (Table 1) and identification of key biological resources (Table 2). Table 3 contains statistical tests and oligonucleotides but is not scored. If SciScore makes any mistakes, please contact us to help us learn and improve.

Table 1: Rigor Adherence Table

Ethics
IRB: 27 The trial was approved by the Ethics Commission of the Canton of Bern, Bern, Switzerland, Nr. 2020-00877 and registered at ClinicalTrials.gov (NCT04510012).
Consent: Patients were included after provision of informed consent.
Inclusion and Exclusion Criteria
Patients with preexisting neuroinflammatory disorders were excluded from the analysis.
Attrition
not detected.
Sex as a biological variable
not detected.
Subject Demographics
Age: not detected.
Weight: not detected.
Randomization
not detected.
Blinding
not detected.
Power Analysis
not detected.
Replication
Samples were separated into triplicates that were automatically and independently processed.
Number: Samples were separated into triplicates that were automatically and independently processed.

Cell Line Authentication	
not detected.	
Protocol Information	
Identifiers: 27 The trial was approved by the Ethics Commission of the Canton of Bern, Bern, Switzerland, Nr. 2020-00877 and registered at ClinicalTrials.gov (NCT04510012).	NCT04510012 Warning: study registration date August 10, 2020 is after study start date March 5, 2020

Table 2: Key Resources Table

Your Sentences	REAGENT or RESOURCE	SOURCE	IDENTIFIER
Antibodies			
Humoral and cell-mediated immune response Anti-spike IgG antibody concentrations in serum were measured using the DiaSorin LIAISON(r) SARS-CoV-2	Anti-spike IgG		
Anti-nucleocapsid antibody concentrations were determined using the Abbott SARS-CoV-2 IgG assay run on the Abbott ARCHITECT i2000 instrument Abbott Diagnostics , Abbott Park , IL , USA positive cutoff of >1.4 SC Index (sample control index ratio) .	Anti-nucleocapsid		
In brief , after washing three times with phosphate-buffered saline (PBS) , 250,000 cryopreserved peripheral blood mononuclear cells (PBMCs) per well were incubated (37degC , humified incubator with 5 CO2) on pre-coated 96-well plates overnight and stimulated with S2N (spike-2 protein , nucleocapsid protein) and SNMO (spike protein , nucleocapsid protein , membrane protein , open reading frame proteins) peptide pools in the presence of costimulatory anti-CD28 antibodies .	SNMO (spike protein , nucleocapsid protein , membrane protein , open reading frame proteins) peptide pools in the presence of costimulatory anti-CD28 antibodies		
	anti-CD28		
Positive (anti-CD3 antibodies) and negative controls (neither anti-CD3 antibodies nor peptide pools) were included.	anti-CD3		
Detection antibodies (anti-IFN-g and anti-IL-2) were added and incubated for 2 h .	anti-IFN-g		
	anti-IL-2		
Experimental Models: Cell Lines			
For the measurement of serum neutralizing SARS-CoV-2 antibody titers , 20,000 Vero E6 cells were seeded in a 96-well plate format .	Vero E6		Suggestion: RRID:CVCL_0574(link)
Software and Algorithms			
Anti-nucleocapsid antibody concentrations were determined using the Abbott SARS-CoV-2 IgG assay run on the Abbott ARCHITECT i2000 instrument Abbott Diagnostics , Abbott Park , IL , USA positive cutoff	Abbott		

of >1.4 SC Index (sample control index ratio) .			
33,34 Statistical analyses were performed using Stata software version 16.0	Stata		
Figures were created using R and GraphPad, version 8.0 (LaJolla, CA, USA).	GraphPad		Suggestion: GraphPad Prism, RRID:SCR_002798(link)

Other Entities Detected

Your Sentences	Recognized Entity
Statistical Tests	
Statistics Demographics were analyzed using descriptive statistics (Fishers exact test) .	Fishers exact test
Mann-Whitney U tests were used to assess the associations between dichotomous COVID-19 severity (mild-moderate and severe) , cytokine levels , antibody concentrations , CMI (SFU250,000 cells) , and sNfL concentrations .	Mann-Whitney U tests
31 Cytokine levels , antibody concentrations , CMI (SFU250,000 cells) , and sNfL concentrations in paired samples over time were compared using Wilcoxon signed-rank test .	Wilcoxon signed-rank test

SciScore is an automated tool that is designed to assist expert reviewers by finding and presenting formulaic information scattered throughout a paper in a standard, easy to digest format. ***SciScore is not a substitute for expert review.*** SciScore also checks for the presence and correctness of several unique identifiers, including RRIDs (research resource identifiers) in the manuscript, detects sentences that appear to be missing RRIDs, and can even suggest RRIDs under certain circumstances. **All RRID suggestions should be verified;** only the author can know whether the suggestions are correct.

For a full description of scored criteria and tips for improving your score, please see https://www.scicrunch.com/sciscorereport-faq