Document Identifier: **PMC8923632**

SciScore Report

Below you will find your SciScore report containing three tables. Your score is calculated based on adherence to scientific rigor criteria (Table 1) and identification of key biological resources (Table 2). Table 3 contains statistical tests and oligonucleotides but is not scored. If SciScore makes any mistakes, please contact us to help us learn and improve.

Table 1: Rigor Adherence Table

Ethics
IRB: Animals The Animal Ethics Committee of the Faculty of Medicine, Chiang Mai University approved the use of mice in our study (protocol No. 47/2020).
Inclusion and Exclusion Criteria
not detected.
Attrition
not detected.
Sex as a biological variable
Male and female mice bred at the Institute of Cancer Research were obtained from the National Animal Institute, Mahidol University, Bangkok, Thailand.
Subject Demographics
Age: Collection of cumulus-oocyte complexes For superovulation, 5- to 9-week-old female mice were injected intraperitoneally (IP) with 10 units of pregnant mare’s serum gonadotropin (Sigma, St. Louis, MO, USA).
Randomization
not detected.
Blinding
not detected.
Power Analysis
not detected.
Replication
not required.

Table 2: Key Resources Table

Your Sentences	REAGENT or RESOURCE	SOURCE	IDENTIFIER
Antibodies			
Washed blastocysts were exposed to rabbit anti-mouse antibody (Sigma M5774; concentration 1:50) for 30 minutes at 37°C , then washed in calcium and magnesium-free PBS , and transferred into a solution containing: (1) guinea pig complement serum (Sigma S1639; concentration 1:4) , (2) 10 µg/mL Hoechst 33342 (Sigma H1399) , and (3) 20 µg/mL propidium iodide (Sigma P4170) for 30 minutes at 37°C .	anti-mouse		
Software and Algorithms			
Washed blastocysts were exposed to rabbit anti-mouse antibody (Sigma M5774; concentration 1:50) for 30 minutes at 37°C , then washed in calcium and magnesium-free PBS , and transferred into a solution containing: (1) guinea pig complement serum (Sigma S1639; concentration 1:4) , (2) 10 µg/mL Hoechst 33342 (Sigma H1399) , and (3) 20 µg/mL propidium iodide (Sigma P4170) for 30 minutes at 37°C .	Sigma		
Statistical analysis The statistical analysis was performed using SPSS ver .	SPSS		Suggestion: SPSS, RRID:SCR_002865(link)

Other Entities Detected

Your Sentences	Recognized Entity
Statistical Tests	
The fertilization rates and blastocyst formation rates among the four experimental groups were compared using the chi-square test.	chi-square test
The mean numbers of ICM, TE, and total cells, as well as the ICM-to-TE ratio, were compared by one-way analysis of variance (ANOVA) when the data distribution was normal, or the Kruskal-Wallis test when normality could not be confirmed.	analysis of variance
	ANOVA
	Kruskal-Wallis test

SciScore is an automated tool that is designed to assist expert reviewers by finding and presenting formulaic information scattered throughout a paper in a standard, easy to digest format. ***SciScore is not a substitute for expert review.*** SciScore also checks for the presence and correctness of several unique identifiers, including RRIDs (research resource identifiers) in the manuscript, detects sentences that appear to be missing RRIDs, and can even suggest RRIDs under certain circumstances. **All RRID suggestions should be verified;** only the author can know whether the suggestions are correct.

For a full description of scored criteria and tips for improving your score, please see https://www.scicrunch.com/sciscorereport-faq