Document Identifier: **PMC8927978**

SciScore Report

Below you will find your SciScore report containing three tables. Your score is calculated based on adherence to scientific rigor criteria (Table 1) and identification of key biological resources (Table 2). Table 3 contains statistical tests and oligonucleotides but is not scored. If SciScore makes any mistakes, please contact us to help us learn and improve.

Table 1: Rigor Adherence Table

Ethics
IRB: MATERIALS AND METHODS The clinical trial study was approved by the ethics committee of Arak University of Medical Sciences (IR.ARAKMU. REC.1397.259) and registered in the Iranian Registry of Clinical Trials (IRCT2019022404282DN1).
Consent: All participants and their parents were briefed about the study and the parents willingly signed informed consent forms.
Inclusion and Exclusion Criteria
The eligibility criteria were as follows: Inclusion criteria: (a) Genetically and systemically healthy children between 4 and 7 years, (b) absence of any dental history or hospitalization, (c) patients with positive or definitely positive behavior according to the Frankl behavior rating scale, (d) children requiring pulpotomy of bilateral primary maxillary first molars.
Attrition
To increase the study power, ten patients were enrolled in each group.
Sex as a biological variable
not detected.
Subject Demographics
Age: Figure 1 CONSORT flow diagram of study procedure Participants, eligibility criteria, and settings The patients were selected among 4-7-year-old children presenting to a pediatric dentistry private office in Arak city, Iran, between July 2019 and December 2019 using convenience sampling.
Randomization
Study design This split-moth crossover randomized clinical trial was conducted on 60 patients.
Blinding
The patients and the operator were not blinded to the group assignments because it was not possible.
Power Analysis

Sample size calculation Sample size was calculated to be 9 in each group according to a study by Ghadimi et al.,9 assuming 0.5-unit level of significance for the FPR scale, 95 confidence interval, 80 study power, and alpha = 0.05.
Replication
not required.

Table 2: Key Resources Table

Your Sentences	REAGENT or RESOURCE	SOURCE	IDENTIFIER
Software and Algorithms			
Statistical analysis The collected data were analyzed using SPSS version 23 through the crossover clinical trial analysis according to the method suggested by Reed (ABBA crossover model).	SPSS		Suggestion: SPSS, RRID:SCR_002865(link)

Other Entities Detected

Your Sentences	Recognized Entity
Statistical Tests	
The means, differences, and sums of PR, SEM, and FPR scales in all groups were calculated in the third and fourth treatment sessions using independent student t-test.	t-test

SciScore is an automated tool that is designed to assist expert reviewers by finding and presenting formulaic information scattered throughout a paper in a standard, easy to digest format. ***SciScore is not a substitute for expert review.*** SciScore also checks for the presence and correctness of several unique identifiers, including RRIDs (research resource identifiers) in the manuscript, detects sentences that appear to be missing RRIDs, and can even suggest RRIDs under certain circumstances. **All RRID suggestions should be verified;** only the author can know whether the suggestions are correct.

For a full description of scored criteria and tips for improving your score, please see https://www.scicrunch.com/sciscorereport-faq