Document Identifier: **PMC8931256**

SciScore Report

Below you will find your SciScore report containing three tables. Your score is calculated based on adherence to scientific rigor criteria (Table 1) and identification of key biological resources (Table 2). Table 3 contains statistical tests and oligonucleotides but is not scored. If SciScore makes any mistakes, please contact us to help us learn and improve.

Table 1: Rigor Adherence Table

Ethics
IRB: Benign human mammary gland tissues were obtained from elective mammoplasty reduction surgeries at the University of California, San Francisco (UCSF cohort A) or Vanderbilt University School of Medicine, which is part of the Cooperative Human Tissue Network (cohort B), in accordance with the institutional review boards approval.
Consent: Tissues were received as deidentified samples, and all subjects provided written informed consent.
Inclusion and Exclusion Criteria
not detected.
Attrition
Seven of 29 samples were subjected to repeat analysis to confirm reproducibility.
Sex as a biological variable
not detected.
Subject Demographics
Age: not detected.
Weight: not detected.
Randomization
not detected.
Blinding
not detected.
Power Analysis
not detected.
Replication
The protein concentration of each replicate was determined using the bicinchoninic acid (BCA) assay.

Table 2: Key Resources Table

Your Sentences	REAGENT or RESOURCE	SOURCE	IDENTIFIER
Antibodies			
Then, the organoids were incubated at 4 degC for 24 h with primary antibodies diluted in IF solution10 FBS (Krt18 Dako M7010 1:100, Krt14 Abcam ab119695 1:200).	Krt18		
	Krt14		
Software and Algorithms			
Briefly, fresh breast tissue was washed with Dulbeccos phosphate-buffered saline (DPBS, Gibco) containing 0.1 gentamycin (Gibco) and 1 penicillinstreptomycin (Gibco).	Gibco		
Organoid size was determined by measuring the maximal cross-sectional organoid area using ImageJFiji.	ImageJFiji		
Proteins were identified using MaxQuant software and the UniProt human database (version 20190501) with isoform sequences.	UniProt		
The SWATH spectral library was built based on MaxQuant search results using Skyline (version 19.1) (81).	MaxQuant		
A total of 3,256 protein groups were quantified based on Skyline searching.	Skyline		

Other Entities Detected

Your Sentences	Recognized Entity
Statistical Tests	
The significance of the branching morphology data was determined by using a two-sample Wilcoxon test.	Wilcoxon test
The average Pearson correlation coefficient of the duplicate protein levels was 0.98.	Pearson correlation
Multiple-sample ANOVA tests were performed with and without patient 16, which results in 1,323 or 657 significant proteins, respectively.	ANOVA
For per patient comparisons among treatments, a two-sample Students t test was performed between each chemical exposure and control culture (DMSO).	t test
Determinations of individual up- or down-regulated proteins were based on the results of two-sample Students t tests.	Students t tests

SciScore is an automated tool that is designed to assist expert reviewers by finding and presenting formulaic information scattered throughout a paper in a standard, easy to digest format. ***SciScore is not a substitute for expert review.*** SciScore also checks for the presence and correctness of several unique identifiers, including RRIDs (research resource identifiers) in the manuscript, detects sentences that appear to be missing RRIDs, and can even suggest RRIDs under certain circumstances. **All RRID suggestions should be verified;** only the author can know whether the suggestions are correct.

For a full description of scored criteria and tips for improving your score, please see https://www.scicrunch.com/sciscorereport-faq