Document Identifier: **PMC8935403**

SciScore Report

Below you will find your SciScore report containing three tables. Your score is calculated based on adherence to scientific rigor criteria (Table 1) and identification of key biological resources (Table 2). Table 3 contains statistical tests and oligonucleotides but is not scored. If SciScore makes any mistakes, please contact us to help us learn and improve.

Table 1: Rigor Adherence Table

Ethics
IRB: Materials and methods Rat model of apical periodontitis in immature permanent teeth The animal experiments were approved by the Ethics Committee of Sun Yat-sen University (Document No. 2021001248).
Euthanasia Agents: The rats were anaesthetized with Thiopental (0.2 mL100 g body weight) and sacrificed by exsanguination (n = 3 each group).
Inclusion and Exclusion Criteria
not detected.
Attrition
not detected.
Sex as a biological variable
not detected.
Subject Demographics
Age: Briefly, 5-week-old Sprague-Dawley rats were anesthetized by intraperitoneally injecting 2 pentobarbital sodium.
Randomization
not detected.
Blinding
not detected.
Power Analysis
not detected.
Replication
Statistical analysis All acquired data are tested by triplicate independent experiments and presented as the mean +- standard deviation.

Number: Statistical analysis All acquired data are tested by triplicate independent experiments and presented as the mean +- standard deviation.
Cell Line Authentication
not detected.

Table 2: Key Resources Table

Your Sentences	REAGENT or RESOURCE	SOURCE	IDENTIFIER
Antibodies			
Cells were then incubated overnight with primary antibody: anti-cytokeratin antibody (1:100 dilution Boster Bio , Hubei , China) and anti-vimentin antibody (1:100 dilution Boster Bio , Hubei , China) and thereafter secondary antibody: DyLight(r) 488 goat anti-mouse IgG (1:200 , EarthOx , CA , USA) .	anti-cytokeratin		
	anti-vimentin		
	anti-mouse IgG		
anti-CD44H-Alexa Fluor 647 , anti-CD45-PE , and anti-CD90-PE antibodies (Biolegend , San Diego , CA , USA) in the dark .	anti-CD45-PE		
	anti-CD90-PE		
After blocking with 5 milk for 1 h at room temperature , PVDF membrane were incubated overnight at 4degC with following primary antibodies: anti-NFIC antibody (1:1000 , Abcam , Burlingame , CA , USA) , anti-DSPP antibody (1:500 , sc-73632 , Santa Cruz , CA , USA) , anti-DMP1 antibody (1:1000 , NBP 1-45525 , Novus Biologicals , Littleton , CO , Massachusetts , USA)	anti-NFIC		
	anti-DSPP		
Sections were deparaffinized , heat retrieved , blocked , and thereafter incubated with the primary antibodies: anti-DMP1 antibody (1:200 , NBP 1-45525 , Novus Biologicals , Littleton , CO , USA)	anti-DMP1		
, anti-TNF-a antibody (1:200 , bs-10802R , Bioss , Beijing , China) , anti-Caspase3 antibody (1:100 , GB11009-1 , Servicebio , Wuhan , China)	anti-TNF-a		
	bs-10802R , Bioss , Beijing , China		
	anti-Caspase3		
anti-CD63 antibody (1:200 , DF2305 , Affinity Biosciences , OH , USA) , overnight at 4degC , followed by washing and incubation with HRP-conjugated secondary antibodies .	anti-CD63		
Experimental Models: Cell Lines			
HEK293FT cells were cultured in high-glucose Dulbeccos modified Eagle medium (DMEM Gibco , Grand Island , NY , USA) containing 10 fetal bovine serum (FBS) and 1 penicillinstreptomycin under 37degC and 5 CO2 .	HEK293FT		Suggestion: RRID:CVCL_6911(link)

Lentivirus was utilized to transfect SCAPs (MOI = 35) and 293T cell line (MOI = 20) at 50-60 confluence with the incubation of 4 mgmL Polybrene (Hanbio , Shanghai , China) .	293T		Suggestion: RRID:CVCL_0063(link)
Experimental Models: Organisms/Strains			
Briefly, 5-week-old Sprague-Dawley rats were anesthetized by intraperitoneally injecting 2 pentobarbital sodium.	Sprague-Dawley		Suggestion: RRID:RGD_70508(link)
Software and Algorithms			
20 Five microliters of LPS from Escherichia coli O111:B4 (10 mgmL, Sigma, USA) diluted in aMEM (Gibco, Grand Island, NY, USA) was injected onto exposed pulp tissue.	Sigma		Suggestion: sigma-tau (RRID:SCR_000488)(link)
Statistics were analyzed with FlowJo software (BD Biosciences , USA)	FlowJo		Suggestion: FlowJo, RRID:SCR_008520(link)
For quantitative analysis of Massons trichrome staining , the blue-stained collagen fiber area was quantified using software ImageJ (NIH , Bethesda , MD , USA) and divided by total tissue area to calculate the collagen volume fraction (n = 3) .	ImageJ		Suggestion: ImageJ, RRID:SCR_003070(link)
Statistical analysis was performed using SPSS 22.0 software (SPSS, Inc., Chicago, IL, USA) and $p < 0.05$ was considered statistically significant.	SPSS		Suggestion: SPSS, RRID:SCR_002865(link)

Other Entities Detected

Your Sentences	Recognized Entity
Statistical Tests	
Differences between two groups were examined by Students t test (normal distribution) or Mann-Whitneys U test (nonnormal distribution).	Students t test
	Mann-Whitneys U test
Multiple group comparisons were conducted by one-way analysis of variance (normal distribution and homogeneity of variance) or Kruskal-Wallis H-test (nonnormal distribution or heterogeneity of variance).	analysis of variance
	Kruskal-Wallis H-test
	heterogeneity of variance
Oligonucleotides	
DMP-1 Rat CTGGTATCAGGTCGGAAGAATC CTCTCATTAGACTCGCTGTCAC	CTGGTATCAGGTCGGAAGAATC
ALP Rat GGAAGGAGGCAGGATTGA TCAGCAGTAACCACAGTCA	GGAAGGAGGCAGGATTGA
NFIC Rat ATGTATTCCTCCCCGCTCTGCC CGGGTTGGACAGCACACAGC b-actin Rat CAGCACTGTGTTGGCATA CGGTCAGGTCATCACTATC NFIC Human GCGGCGATTACTACACTTC CTTGTCCATCTCTGTCTTCTTC	CAGCACTGTGTTGGCATA CGGTCAGGTCATCACTATC NFIC Human GCGGCGATTACTACACTTC CTTGTCCATCTCTGTCTTCTTC
b-actin Human CATTGCTGACAGGATGCAGA CTGCTGGAAGGTGGACAGTGA Histological and immunohistochemistry analysis The mandibles of apical periodontitis rat model and transplanted scaffold tissues were dissected and fixed in 4 buffered paraformaldehyde for 24 h , then decalcified with 10 ethylenediaminetetraacetic acid (EDTA) for 2 months , dehydrated in a gradient alcohol series , and embedded in paraffin .	Human CATTGCTGACAGGATGCAGA CTGCTGGAAGGTGGACAGTGA

SciScore is an automated tool that is designed to assist expert reviewers by finding and presenting formulaic information scattered throughout a paper in a standard, easy to digest format. ***SciScore is not a substitute for expert review.*** SciScore also checks for the presence and correctness of several unique identifiers, including RRIDs (research resource identifiers) in the manuscript, detects sentences that appear to be missing RRIDs, and can even suggest RRIDs under certain circumstances. **All RRID suggestions should be verified;** only the author can know whether the suggestions are correct.

For a full description of scored criteria and tips for improving your score, please see https://www.scicrunch.com/sciscorereport-faq