Document Identifier: **PMC8938331**

SciScore Report

Below you will find your SciScore report containing three tables. Your score is calculated based on adherence to scientific rigor criteria (Table 1) and identification of key biological resources (Table 2). Table 3 contains statistical tests and oligonucleotides but is not scored. If SciScore makes any mistakes, please contact us to help us learn and improve.

Table 1: Rigor Adherence Table

Ethics
Consent: Each patient provided informed consent for the analysis of tissues, which were subjected to paraffin embedding.
IRB: All procedures involved human liver samples in this study were approved by the Ethics Committee of Nanjing Medical University First Affiliated Hospital and adhered to the principles of the Declaration of Helsinki.
IACUC: All animal procedures were performed in accordance with protocols approved by the Institutional Animal Care and Usage Committee of Nanjing Medical university.
Field Sample Permit: HEK293T cells used in this study were purchased from the Type Culture Collection of the Chinese Academy of Sciences (Shanghai, China).
Euthanasia Agents: Mice were euthanized by CO2 inhalation, and the tissues were collected for subsequent experiments.
Inclusion and Exclusion Criteria
Accordingly, liver samples from individuals with excessive alcohol consumption (>140 g for men or >70 g for women, per week), drug abuse or viral infection (e.g., hepatitis B virus and hepatitis C virus) were excluded 21.
Attrition
not detected.
Sex as a biological variable
Accordingly, liver samples from individuals with excessive alcohol consumption (>140 g for men or >70 g for women, per week), drug abuse or viral infection (e.g., hepatitis B virus and hepatitis C virus) were excluded 21.
Subject Demographics
Age: The fatty liver model was established in 7- to 8-week-old male mice by feeding a HFD (D12492, Research Diets) for 16 weeks or a Western diet (WD, D12079B, Research Diets) for 8 weeks.
Randomization
not detected.

Blinding
not detected.
Power Analysis
not detected.
Replication
not required.
Cell Line Authentication
not detected.

Table 2: Key Resources Table

Your Sentences	REAGENT or RESOURCE	SOURCE	IDENTIFIER
Antibodies			
Cells were subsequently stained with the antibody against TLR4 (TLR4-APC, SA15-21, BioLegend) for 30 min on ice.	TLR4		
	SA15-21		
Experimental Models: Cell Lines			
HEK293T cells were cultured in DMEM supplemented with 10 FBS and 1 PS in a 5 CO2 humidified atmosphere.	HEK293T		Suggestion: RRID:CVCL_0063(link)
Experimental Models: Organisms/Strains			
To identify the efficiency of SAA12 knockout, an acute phase response was elicited by an intraperitoneal injection of 1 mg g-1 lipopolysaccharide (LPS, L4516, Sigma-Aldrich) into male SAA12 -- mice and WT mice.	SAA12 --		
Male TLR4 -- (Tlr4lps-del mutmut, C57BL10ScNJNju) mice, and control mice (C57BL10JNju) were purchased from the Model Animal Research Center of Nanjing University (Nanjing, China).	TLR4 --		
	C57BL10ScNJNju		
	C57BL10JNju		
To knock down hepatic SAA12, the AAV8 encoding SAA12-specific short hairpin (sh) RNA (AAV8-shRNA-SAA12: 5-GCGAGCCTACACTGACATGAA-3, Shandong WZ Biosciences Co., Ltd.) was intravenously injected into adult male C57BL6J mice.	C57BL6J		Suggestion: RRID:IMSR_JAX:000664(link)
Recombinant DNA			
2.4 Protein expression Mouse SAA1 (residues 19-122) was cloned into the pET30(a)+ expression vector between NdeI and HindIII restriction endonuclease sites, with a N-terminal hexa-histidine and subsequently transformed into BL21 (DE3) strain of Escherichia coli.	pET30		Suggestion: RRID:Addgene_21267(link)
Software and Algorithms			
The blots were quantified by Image J Software and normalized to levels of GAPDH.	Image J		Suggestion: ImageJ, RRID:SCR_003070(link)
2.14 Statistical analysis Data were analyzed using GraphPad Prism 8.0	GraphPad Prism		

software and presented as mean +- standard error of the mean (SEM).			

Other Entities Detected

Your Sentences	Recognized Entity
Statistical Tests	
Comparisons of two groups were performed by Students t-test, and comparisons among multiple groups were performed using ANOVA followed by Bonferronis post hoc test.	t-test
	ANOVA
	Bonferronis post hoc test
Oligonucleotides	
To knock down hepatic SAA12, the AAV8 encoding SAA12-specific short hairpin (sh) RNA (AAV8-shRNA-SAA12: 5-GCGAGCCTACACTGACATGAA- 3, Shandong WZ Biosciences Co., Ltd.) was intravenously injected into adult male C57BL6J mice.	5-GCGAGCCTACACTGACATGAA- 3

SciScore is an automated tool that is designed to assist expert reviewers by finding and presenting formulaic information scattered throughout a paper in a standard, easy to digest format. ***SciScore is not a substitute for expert review.*** SciScore also checks for the presence and correctness of several unique identifiers, including RRIDs (research resource identifiers) in the manuscript, detects sentences that appear to be missing RRIDs, and can even suggest RRIDs under certain circumstances. **All RRID suggestions should be verified;** only the author can know whether the suggestions are correct.

For a full description of scored criteria and tips for improving your score, please see https://www.scicrunch.com/sciscorereport-faq