Document Identifier: **PMC8965458**

SciScore Report

Below you will find your SciScore report containing three tables. Your score is calculated based on adherence to scientific rigor criteria (Table 1) and identification of key biological resources (Table 2). Table 3 contains statistical tests and oligonucleotides but is not scored. If SciScore makes any mistakes, please contact us to help us learn and improve.

Table 1: Rigor Adherence Table

Ethics
Consent: Ethics Patient clinical and epidemiological data, along with their clinical specimens were collected after informed written consent was obtained under protocols 16-066 and 200829003, which were reviewed and approved by the Scientific Ethics Committee for Health Sciences (CECSaludUC, by its Spanish acronym) at Pontificia Universidad Católica de Chile (PUC).
IRB: Ethics Patient clinical and epidemiological data, along with their clinical specimens were collected after informed written consent was obtained under protocols 16-066 and 200829003, which were reviewed and approved by the Scientific Ethics Committee for Health Sciences (CECSaludUC, by its Spanish acronym) at Pontificia Universidad Católica de Chile (PUC).
Inclusion and Exclusion Criteria
Our study did not have any a priori exclusion criteria and hence all individuals with a laboratory confirmed SARS-CoV-2 infection or that had been vaccinated during the study period were invited to participate in the study.
Attrition
Due to the rapid vaccination campaign in Chile, 36 of these 74 participants were immunized with 1 or 2 doses of either the CoronaVac or BNT162b2 vaccines during the follow up period within 127–398 days (4.2–13.3 months) since onset of symptoms (Supplementary Figure 2 and Supplementary Table 1).
Sex as a biological variable
not detected.
Subject Demographics
Age: not detected.
Weight: not detected.
Randomization
not detected.
Blinding
The samples were assigned an anonymous code and all serological analyses were performed by scientists that were blinded in regards to the subject's clinical condition and time of sample collection.

Power Analysis
not detected.
Replication
Every data point represents results from two technical replicates .
Type: Every data point represents results from two technical replicates .
Number: Every data point represents results from two technical replicates .
Cell Line Authentication
not detected.

Table 2: Key Resources Table

Your Sentences	REAGENT or RESOURCE	SOURCE	IDENTIFIER
Antibodies			
Fig 1 Figure 2 Comparison of neutralizing and anti-N antibody responses after SARS-CoV-2 infection of outpatient and hospitalized individuals over a 12 months period. a . nAb IC50 titres were determined by microneutralization assay of recombinant vesicular stomatitis virus carrying SARS-CoV-2 spike protein (rVSV-SARS2-S) . b .	anti-N		
	SARS-CoV-2 spike protein (rVSV-SARS2-S)		
Fig 2 Figure 3 Longitudinal neutralizing and anti-N antibody titres to SARS-CoV-2 in previously infected before and after CoronaVac or BNT162b2 vaccination . nAb titres (IC50) obtained using a rVSV-SARS2-S microneutralization assay and end-point titres of anti-N IgG were determined by ELISA using a recombinant SARS-CoV-2 nucleocapsid protein for vaccinated previously infected outpatients (a-b; 20 participants) or vaccinated hospitalized patients (c-d; 16 participants) at different time points grouped by weeks (W) or months (M) post-symptom onset (serum samples from: 1W = 1-7 days; 2W = 8-14 days; 3W = 15-21 days; 4W = 22-45 days; 3M = 46-135 days; 6M = 136-225 days; 9M = 226-315 days and 12M = 316-405 days/12-15M = 316-495) .	anti-N IgG		
	days/12-15M = 316-495		
For primary antibody detection a 1:3,000 dilution of goat anti-human IgG–horseradish peroxidase (HRP) conjugated secondary antibody (Thermo Fisher Scientific # SA1-36011 , RRID:AB_1075961) was added to each well for 1 h and SIGMAFAST OPD (o-phenylenediamine dihydrochloride; Sigma–Aldrich #P9187) was used as substrate.	anti-human IgG–horseradish	Thermo Fisher Scientific	(Thermo Fisher Scientific Cat# SA1-36011, RRID:AB_1075961)(link)
	SA1-36011		
CR3022 , a human monoclonal antibody reactive to the RBD of both SARS-CoV-1 and SARS-CoV-2 , 19 , 20 was used as control .	SARS-CoV-2		

For detecting viral infection , a primary mAb 1C7 (anti-SARS nucleoprotein antibody generated in-house) was used at a 1:1,000 dilution and subsequently detected with a 1:3,000 dilution of a goat anti-mouse IgG–HRP (Rockland #KCB002 , RRID:AB_10703407) , and incubation with SIGMAFAST OPD (Sigma-Aldrich) as described above.	anti-SARS nucleoprotein		
	anti-mouse IgG–HRP	Rockland	(Rockland Cat# KCB002, RRID:AB_10703407)(link)
Experimental Models: Cell Lines			
21 Briefly , Vero E6 cells (ATCC #CRL-1586 , RRID:CVCL_0574) were seeded at a density of 20,000 cells per well in a 96-well cell culture plate in complete Dulbecco 's Modified Eagle Medium (cDMEM , Gibco Thermo Fisher Scientific #11995040) .	Vero E6	CLS	(CLS Cat# 305008, RRID:CVCL_0574)(link)
After removing the cell culture media , the Vero E6 cells were incubated with 120 µL of the virus-serum mixture at 37 °C for 1 h .	Vero E6		Suggestion: RRID:CVCL_0574(link)
The serum-virus inoculum was added to Vero E6 hACE2 cells seeded the day before in optical bottom 96-well plates (Thermo Scientific #165305) at 80 % confluence and adsorbed for 2 h at 37 °C .	hACE2		
Recombinant DNA			
Briefly , Vero E6 cells (ATCC # CRL-1586 , RRID:CVCL_0574) grown in 1X MEM (Gibco #11095-080) supplemented with 10 % FBS (Gibco , #16000-044) were transfected with plasmid pCEP4-myc-ACE2 (Addgene catalog # 141185) and stable clones were selected by hygromycin (Invitrogen #10687010) (400 µg/mL) .	pCEP4-myc-ACE2		
Software and Algorithms			
For primary antibody detection a 1:3,000 dilution of goat anti-human IgG–horseradish peroxidase (HRP) conjugated secondary antibody (Thermo Fisher Scientific # SA1-36011 , RRID:AB_1075961) was added to each well for 1 h and SIGMAFAST OPD (o-phenylenediamine dihydrochloride; Sigma–Aldrich #P9187) was used as substrate.	Sigma–Aldrich		

Briefly , Vero E6 cells (ATCC # CRL-1586 , RRID:CVCL_0574) grown in 1X MEM (Gibco #11095-080) supplemented with 10 % FBS (Gibco , #16000-044) were transfected with plasmid pCEP4-myc-ACE2 (Addgene catalog # 141185) and stable clones were selected by hygromycin (Invitrogen #10687010) (400 μg/mL) .	Gibco		
Viral infectivity was quantified by automated enumeration of GFP-positive cells (normalizing against cells stained with DAPI) using a Cytation5 automated fluorescence microscope (BioTek) and segmentation algorithms applied from the ImageJ program.	ImageJ		Suggestion: ImageJ, RRID:SCR_003070(link)
The half-maximum inhibitory concentration (IC50) of the sera, were calculated from data obtained with two technical replicates using non-linear regression analysis and the curve fitting was done using second-order polynomial (quadratic); and linear regression models (using log10 IC50 transformed data) were done with GraphPad Prism 5 software.	GraphPad Prism		
GraphPad Prism 8 was used for statistical analysis: *P < 0·05; **P < 0·01; ***P < 0·001; ****P < 0·0001.	GraphPad		Suggestion: GraphPad Prism, RRID:SCR_002798(link)

Other Entities Detected

Your Sentences	Recognized Entity
Statistical Tests	
Statistical analyses shown at the indicated time points were performed between nAb titres of outpatient and hospitalized using the unpaired two-tailed Mann-Whitney test (*P < 0·05; **P < 0·01; **P < 0·001; ****P < 0·0001; ns , non-significant) .	Mann-Whitney test
[Fisher 's exact test; Mann Whitney test] .	Fisher 's exact test; Mann Whitney test
Association between categorical variables was examined with Chi-squared or Fisher's exact test.	Chi-squared
	Fisher's exact test
Correlation was evaluated calculating the Pearson correlation coefficient.	Pearson correlation
Unresolved RRIDs	
After 48 h at 37 °C , the cells were fixed with 10 % paraformaldehyde (Polysciences # 04018-1) for 24 h at 4 °C , permeabilized with PBS containing 0·1 % Triton X-100 (Sigma-Aldrich # X100) and the plates were and blocked with 3 % milk (American Bio # AB1010901000) in PBST .	AB1010901000

SciScore is an automated tool that is designed to assist expert reviewers by finding and presenting formulaic information scattered throughout a paper in a standard, easy to digest format. ***SciScore is not a substitute for expert review.*** SciScore also checks for the presence and correctness of several unique identifiers, including RRIDs (research resource identifiers) in the manuscript, detects sentences that appear to be missing RRIDs, and can even suggest RRIDs under certain circumstances. **All RRID suggestions should be verified;** only the author can know whether the suggestions are correct.

For a full description of scored criteria and tips for improving your score, please see https://www.scicrunch.com/sciscorereport-faq