Document Identifier: **PMC8968669**

SciScore Report

Below you will find your SciScore report containing three tables. Your score is calculated based on adherence to scientific rigor criteria (Table 1) and identification of key biological resources (Table 2). Table 3 contains statistical tests and oligonucleotides but is not scored. If SciScore makes any mistakes, please contact us to help us learn and improve.

Table 1: Rigor Adherence Table

Ethics
Consent: Informed consent was obtained from all participants, which contained a detailed description of all study procedures, as well as any possible risks andor benefits.
IRB: All data were collected according to ethical guidelines as outlined by the Declaration of Helsinki all procedures were approved by the Northwestern University Feinberg School of Medicine Institutional Review Board.(WAIS-III, Wechsler, 1997).
Inclusion and Exclusion Criteria
not detected.
Attrition
not detected.
Sex as a biological variable
not detected.
Subject Demographics
Age: not detected.
Weight: not detected.
Randomization
not detected.
Blinding
not detected.
Power Analysis
not detected.
Replication
not required.

Table 2: Key Resources Table

Your Sentences	REAGENT or RESOURCE	SOURCE	IDENTIFIER
Software and Algorithms			
Repeated measures multivariate analysis of covariance models (RM-MANCOVA) were used to assess the overall effect of group status on the volume of each brain structure across right and left hemisphere, which included supratentorial volume (derived from FreeSurfer) as a covariate to account for the global effects of whole brain volume.	FreeSurfer		Suggestion: FreeSurfer, RRID:SCR_001847(link)

Other Entities Detected

Your Sentences	Recognized Entity
Statistical Tests	
2.4 Statistical analysis Between-group differences in demographics and cognition were assessed with chi-square and analysis of variance (ANOVA) models while independent samples t-tests were used to evaluate differences in antipsychotic medication between SCZ groups.	chi-square
	analysis of variance
	ANOVA
	t-tests
Repeated measures multivariate analysis of covariance models (RM-MANCOVA) were used to assess the overall effect of group status on the volume of each brain structure across right and left hemisphere, which included supratentorial volume (derived from FreeSurfer) as a covariate to account for the global effects of whole brain volume.	RM-MANCOVA
If a model was significant for the group effect, then follow-up one-way RM-ANCOVA models were conducted as a post hoc analysis to assess each specific group contrast.	RM-ANCOVA

SciScore is an automated tool that is designed to assist expert reviewers by finding and presenting formulaic information scattered throughout a paper in a standard, easy to digest format. ***SciScore is not a substitute for expert review.*** SciScore also checks for the presence and correctness of several unique identifiers, including RRIDs (research resource identifiers) in the manuscript, detects sentences that appear to be missing RRIDs, and can even suggest RRIDs under certain circumstances. **All RRID suggestions should be verified;** only the author can know whether the suggestions are correct.

For a full description of scored criteria and tips for improving your score, please see https://www.scicrunch.com/sciscorereport-faq