Document Identifier: **PMC8968879**

SciScore Report

Below you will find your SciScore report containing three tables. Your score is calculated based on adherence to scientific rigor criteria (Table 1) and identification of key biological resources (Table 2). Table 3 contains statistical tests and oligonucleotides but is not scored. If SciScore makes any mistakes, please contact us to help us learn and improve.

Table 1: Rigor Adherence Table

Ethics
IACUC: Animal experiments All animal experiments were carried out in accordance with the guidelines of the Institutional Animal Care and Use Committee (IACUC).
Inclusion and Exclusion Criteria
not detected.
Attrition
not detected.
Sex as a biological variable
not detected.
Subject Demographics
Age: not detected.
Weight: not detected.
Randomization
not detected.
Blinding
not detected.
Power Analysis
not detected.
Replication
not required.
Cell Line Authentication
not detected.

Table 2: Key Resources Table

Your Sentences	REAGENT or RESOURCE	SOURCE	IDENTIFIER
Antibodies			
For surface LAMP-5 detection , cells were incubated with 3 mg anti-human LAMP-5 therapeutic antibody overnight and then stained with anti-mouse IgG1-PE .	anti-human LAMP-5		
	anti-mouse IgG1-PE		
Cells were incubated with LAMP-5 therapeutic antibody (Creative Biolabs) and anti-mouse immunoglobulin G (IgG) Fc-DM1 antibody with non-cleavable linker (Moradec) at 5 nguL and 1 nguL final concentrations , respectively .	anti-mouse immunoglobulin G (IgG		
	Fc-DM1		
Cells were stained overnight with primary antibodies against LAMP-5, MYD88, and LAMP-1.	LAMP-5, MYD88		
	LAMP-1		
Experimental Models: Cell Lines			
In xenograft experiments bone marrow samples were collected 4 weeks after transplantation as well as when signs of leukemia were present bone marrow aspirates were analyzed via flow cytometry for the presence of human CD45+ cells and the presence of short hairpin RNA (shRNA)-transduced Venus+ cells Retroviral and lentiviral transductions Retroviral and lentiviral supernatants were generated by transfection of HEK293T cells using the FuGENE 6 reagent (Promega) or Lipofectamine 3000 (Thermo Fisher Scientific) according to the manufacturers recommendations.	HEK293T		Suggestion: RRID:CVCL_0063(link)
, THP-1 , and Kasumi-1 cells were plated at 10,000 cells per well in a 96-well plate.	Kasumi-1		Suggestion: RRID:CVCL_0589(link)
Experimental Models: Organisms/Strains			
For xenograft experiments with MV411 and MLL-AF10 primary patient cells, immunocompromised NOD-Rag1null IL2rg null (NRG) (Jackson Laboratories, stock no. 007799) recipient mice were conditioned with busulfan and transplanted 24 hours later.	NOD-Rag1null IL2rg		
Software and Algorithms			

Cells transduced with constructs containing a fluorescent marker (Venus) were sorted 4-5 days after transduction by using MoFlo XDP (Beckman Coulter), FACSAria (BD Biosciences), or a SONY SH800S (Sony	BD Biosciences		
Data were acquired on a BD FACSCanto analyzer and results were analyzed using FlowJo Version 10 (BD Biosciences)	FlowJo		Suggestion: FlowJo, RRID:SCR_008520(link)
Statistical analysis was performed using GraphPad Prism.	GraphPad Prism		

Other Entities Detected

Your Sentences	Recognized Entity
Statistical Tests	
t-tests were two-tailed unless otherwise stated.	t-tests

SciScore is an automated tool that is designed to assist expert reviewers by finding and presenting formulaic information scattered throughout a paper in a standard, easy to digest format. ***SciScore is not a substitute for expert review.*** SciScore also checks for the presence and correctness of several unique identifiers, including RRIDs (research resource identifiers) in the manuscript, detects sentences that appear to be missing RRIDs, and can even suggest RRIDs under certain circumstances. **All RRID suggestions should be verified;** only the author can know whether the suggestions are correct.

For a full description of scored criteria and tips for improving your score, please see https://www.scicrunch.com/sciscorereport-faq