Document Identifier: **PMC8969300**

SciScore Report

Below you will find your SciScore report containing three tables. Your score is calculated based on adherence to scientific rigor criteria (Table 1) and identification of key biological resources (Table 2). Table 3 contains statistical tests and oligonucleotides but is not scored. If SciScore makes any mistakes, please contact us to help us learn and improve.

Table 1: Rigor Adherence Table

Ethics
IACUC: 21 DNA Collection and Mutation Analysis Blood samples were collected at each referral center for both diagnostic and research purposes, according to the protocols of the respective Institutional Ethical Committees, and sent for genomic DNA extraction from peripheral leukocytes and molecular analyses andor confirmation to 1 of the 2 major diagnostic participating centers
Consent: Standard Protocol Approvals and Patient Consents The study was conducted according to the criteria set by the Declaration of Helsinki, and each patient or family member, in case of minors, signed an informed consent.
IRB: The study protocol was approved by the Ethical Committee of the Fondazione Policlinico Universitario A. Gemelli IRCCS (Ethical Board approving number: ID 3998).
Inclusion and Exclusion Criteria
not detected.
Attrition
Eight pedigrees included in this study had been already described elsewhere. 13 - 20 Strengthening the Reporting of Observational Studies in Epidemiology cross-sectional reporting guidelines were used.
Sex as a biological variable
The Mann-Whitney U test was used to evaluate the presence of any associations between specific features and various SPAST-HSP groups of patients (malesfemales, probandsfamily members, early-onsetlate-onset cases, short-long-DD cases, and missense variant carrierstruncating-variant carriers).
Subject Demographics
Age: not detected.
Weight: not detected.
Randomization
not detected.
Blinding
not detected.

Power Analysis
not detected.
Replication
not required.

Table 2: Key Resources Table

Your Sentences	REAGENT or RESOURCE	SOURCE	IDENTIFIER
Software and Algorithms			
(IRCCS Fondazione Stella Maris , Pisa , and Ospedale Pediatrico Bambino Gesu , Rome)	Pisa		Suggestion: PISA (RRID:SCR_015749)(link)
23 To define the impact of missense variants on protein function , we used an in silico pipeline encompassing 19 prediction tools including Mutation Assessor , 24 FATHMM , 25 LRT , 26 Deo-gen2 , 26 Eigen , 27 Eigen PC , 27 Sorting Intolerant From Tolerant (SIFT) , 28 SIFT4G , 29 Provean , 30 Minimum Viable Product , 26 Revel , 26 Primate AI , 26 MetaSVM , 26 MetalR , 26 Genomic Evolutionary Rate Profiling , 26 PolyPhen-2 HumDir , 31 PolyPhen-2 HumVar , 31 UMD Predictor , 32 and CADD .	Mutation Assessor		Suggestion: Mutation Assessor (RRID:SCR_024502)(link)
	SIFT		Suggestion: SIFT (RRID:SCR_012813)(link)
	PolyPhen-2		
33 Splicing variants and synonymous variants close to splicing sites were also tested using Human Splicing Finder 3.1 34 and NNSPLICE 0.9 .	Human Splicing Finder		Suggestion: Human Splicing Finder (RRID:SCR_005181)(link)
35 We included variants with minor allele frequency <0.1 in the ExAC , 36 gnomAD , 37 1000 Genomes Project , 38 and dbSNP 39 databases .	ExAC		
	1000 Genomes Project		
Statistical analysis was performed by SPSS (Statistical Package for Social Science, IBM SPSS Statistics, Version 24.0.	SPSS		Suggestion: SPSS, RRID:SCR_002865(link)

Other Entities Detected

Your Sentences	Recognized Entity
Statistical Tests	
Normality of continuous variables was checked using the Kolmogorov-Smirnov test.	Kolmogorov-Smirnov test
One-way analysis of variance (ANOVA) was performed to compare AAO among the generations.	analysis of variance
	ANOVA
The Mann-Whitney U test was used to evaluate the presence of any associations between specific features and various SPAST-HSP groups of patients (malesfemales, probandsfamily members, early-onsetlate-onset cases, short-long-DD cases, and missense variant carrierstruncating-variant carriers).	Mann-Whitney U test
The Kruskal-Wallis test with pairwise comparisons was used to compare qualitative variables when >2 subgroups of patients were analyzed (slowly vs typically vs rapidly progressing cases).	Kruskal-Wallis test
The kh2 test (or Fisher 2-tailed exact test when required) was used to compare categorical variables between the groups.	Fisher 2-tailed exact test

SciScore is an automated tool that is designed to assist expert reviewers by finding and presenting formulaic information scattered throughout a paper in a standard, easy to digest format. ***SciScore is not a substitute for expert review.*** SciScore also checks for the presence and correctness of several unique identifiers, including RRIDs (research resource identifiers) in the manuscript, detects sentences that appear to be missing RRIDs, and can even suggest RRIDs under certain circumstances. **All RRID suggestions should be verified;** only the author can know whether the suggestions are correct.

For a full description of scored criteria and tips for improving your score, please see https://www.scicrunch.com/sciscorereport-faq