Document Identifier: **PMC8973833**

SciScore Report

Below you will find your SciScore report containing three tables. Your score is calculated based on adherence to scientific rigor criteria (Table 1) and identification of key biological resources (Table 2). Table 3 contains statistical tests and oligonucleotides but is not scored. If SciScore makes any mistakes, please contact us to help us learn and improve.

Table 1: Rigor Adherence Table

Ethics
IACUC: Xenograft tumor models All mouse experiments were carried out by Beijing Viewsolid Biotech Co., Ltd., and the animal ethics approval form was 201901003.
Inclusion and Exclusion Criteria
not detected.
Attrition
not detected.
Sex as a biological variable
Male BALBc mice (6 weeks old, 18 g) were purchased from Sibf Biotechnology Co., Ltd. (Beijing, P.R. China).
Subject Demographics
Age: Male BALBc mice (6 weeks old, 18 g) were purchased from Sibf Biotechnology Co., Ltd. (Beijing, P.R. China).
Weight: Male BALBc mice (6 weeks old, 18 g) were purchased from Sibf Biotechnology Co., Ltd. (Beijing, P.R. China).
Randomization
not detected.
Blinding
not detected.
Power Analysis
not detected.
Replication
not required.

Cell Line Authentication	
not detected.	
Data Information	
Identifiers: Expression validation of SVIL-AS1 in GEO datasets The datasets GSE146461 and GSE85716 downloaded from the GEO database of NCBI (https:www.ncbi.nlm.nih.govgeo) , which were based on GPL20115 (Agilent-067406	GSE146461
	GSE85716

Table 2: Key Resources Table

Your Sentences	REAGENT or RESOURCE	SOURCE	IDENTIFIER
Antibodies			
Two micrograms of anti-m6A antibody (ab151230 , Abcam , Burlingame , CA , USA) or anti-IgG (Cell Signaling Technology) were conjugated to the protein AG magnetic beads at 4degC for 4 h in IP buffer (20 mM Tris pH 7.5 , 140 mM NaCl , 1 NP-40 , 2 mM EDTA) .	anti-m6A		
	anti-IgG (Cell Signaling Technology)		
Then , the cells were incubated with BrdU antibody (1:1000) and incubated at 4degC overnight .	BrdU		
The membranes were blocked with 5 nonfat milk for 1 h at room temperature, and then incubated with specific primary antibodies against E2F1 (1:500, Bioss, Beijing, China), METTL3 (1:1000, Abcam), and GAPDH (1:1000, Bioss) overnight at 4degC.	E2F1		
	METTL3		
	GAPDH		
Experimental Models: Cell Lines			
A549 cells were cultured in Hams F-12K medium supplemented with 10 fetal bovine serum (FBS) and 100 UmL penicillin-streptomycin .	A549		Suggestion: RRID:CVCL_0023(link)
H1650 cells were maintained in RPMI-1640 medium containing 10 FBS and 1 PS.	H1650		Suggestion: RRID:CVCL_1483(link)
Experimental Models: Organisms/Strains			
Male BALBc mice (6 weeks old, 18 g) were purchased from Sibf Biotechnology Co., Ltd. (Beijing, P.R. China).	BALBc		
Software and Algorithms			
Overall survival analysis of lncRNAs from AL731577.2 and NORAD was performed using R software (3.6.3) 31 survminer package (0.4.9).	NORAD		
The expression of SVIL-AS1 in LUAD tissues was analyzed using GEPIA2 (http:gepia2.cancer-pku.cngeneral) 32 and R software (3.6.3) 31 ggplot2 package (	ggplot2		Suggestion: ggplot2 (RRID:SCR_014601)(link)
Cell cycle analysis was performed using a FACScan Flow Cytometer (Becton-Dickinson) and analyzed using FlowJo LLC software (Ashland , OR) .	FlowJo		Suggestion: FlowJo, RRID:SCR_008520(link)

Statistical analysis Data were analyzed using the GraphPad Prism version 8.	GraphPad Prism		

Other Entities Detected

Your Sentences	Recognized Entity
Statistical Tests	
Materials and methods Data analysis Pearson correlation analysis was performed to identify co-expressed lncRNAs related to METTL3 with a cutoff of Pearson R > 0.3 and P < 0.05, in The Cancer Genome Atlas (TCGA)-LUAD dataset.	Pearson correlation
The expression of SVIL-AS1 was analyzed in LUAD and normal tissues using the unpaired Students t-test .	t-test
The two-tailed Students t-test was used to compare gene expression between the two groups.	Students t-test
One-way analysis of variance (ANOVA) was used to analyze the differences among multiple groups.	analysis of variance
	ANOVA
Pearsons correlation analysis was used to analyze the correlation between METTL3 and lncRNAs in LUAD samples.	Pearsons correlation
Oligonucleotides	
Primers for qRT-PCR Genes Sequences METTL3 Forward 5-TTGTCTCCAACCTTCCGTAGT-3 Reverse 5-CCAGATCAGAGAGGTGGTGTAG-3 E2F1 Forward 5-ACGCTATGAGACCTCACTGAA-3 Reverse 5-TCCTGGGTCAACCCCTCAAG-3 CCNE1 Forward 5-GCCAGCCTTGGGACAATAATG-3 Reverse 5-CTTGCACGTTGAGTTTGGGT-3 MYBL2 Forward 5-CTTGAGCGAGTCCAAAGACTG-3 Reverse 5-AGTTGGTCAGAAGACTTCCCT-3 CDC25A Forward 5-CTCCTCCGAGTCAACAGATTCA-3 Reverse 5-CAACAGCTTCTGAGGTAGGGA-3 p27 Forward 5-ATCACAAACCCCTAGAGGGCA-3 Reverse 5-GGGTCTGTAGTAGAACTCGGG-3 p15 Forward 5-GGGACTAGTGGAGAAGGTGC-3 Reverse 5-CATCATCATGACCTGGATCGC-3 SVIL-AS1 Forward 5-ACAGAAATCCTAACGGGGCG-3 Reverse 5-CCATGATCTGATCCCTGCCC-3 GAPDH Forward 5-	5-CCATGATCTGATCCCTGCCC-3

CTGGGCTACACTGAGCACC-3 Reverse 5-AAGTGGTCGTTGAGGGCAATG-3 E2F1 promoter (F-5TTCGCGGCAAAAAGGATTT R- 5GCCGCTGCCTGCAAAGT) , GGGACTAGTGGAGAAGGTGC	

SciScore is an automated tool that is designed to assist expert reviewers by finding and presenting formulaic information scattered throughout a paper in a standard, easy to digest format. ***SciScore is not a substitute for expert review.*** SciScore also checks for the presence and correctness of several unique identifiers, including RRIDs (research resource identifiers) in the manuscript, detects sentences that appear to be missing RRIDs, and can even suggest RRIDs under certain circumstances. **All RRID suggestions should be verified;** only the author can know whether the suggestions are correct.

For a full description of scored criteria and tips for improving your score, please see https://www.scicrunch.com/sciscorereport-faq