Document Identifier: **PMC8983315**

SciScore Report

Below you will find your SciScore report containing three tables. Your score is calculated based on adherence to scientific rigor criteria (Table 1) and identification of key biological resources (Table 2). Table 3 contains statistical tests and oligonucleotides but is not scored. If SciScore makes any mistakes, please contact us to help us learn and improve.

Table 1: Rigor Adherence Table

Ethics
Field Sample Permit: Study approval All experiments and procedures described in this study were conducted under an approved protocol of the Institutional Committee for the Good Animal Experimentation of the San Raffaele Scientific Institute and were reported to the Ministry of Health as required by Italian law (Institutional Animal Care and Use Committee - IACUC 791 and 1145).
IRB: Study approval All experiments and procedures described in this study were conducted under an approved protocol of the Institutional Committee for the Good Animal Experimentation of the San Raffaele Scientific Institute and were reported to the Ministry of Health as required by Italian law (Institutional Animal Care and Use Committee - IACUC 791 and 1145).
IACUC: Study approval All experiments and procedures described in this study were conducted under an approved protocol of the Institutional Committee for the Good Animal Experimentation of the San Raffaele Scientific Institute and were reported to the Ministry of Health as required by Italian law (Institutional Animal Care and Use Committee - IACUC 791 and 1145).
Inclusion and Exclusion Criteria
Mice dying within 2-3 weeks after busulfan conditioning were excluded from subsequent analyses.
Attrition
not detected.
Sex as a biological variable
not detected.
Subject Demographics
Age: BMT Total BM was flushed from the tibiae, femora, and humeri of 4- to 8-week-old tgGFP mice as described previously.
Randomization
not detected.
Blinding
not detected.

Power Analysis
not detected.
Replication
not required.
Cell Line Authentication
Authentication: The remaining PB sample aliquots were analyzed using an automated hemocytometer (ProCyte Dx Hematology Analyzer, IDEXX Laboratories, Westbrook, ME, USA) to determine the complete blood count variables.

Table 2: Key Resources Table

Your Sentences	REAGENT or RESOURCE	SOURCE	IDENTIFIER
Antibodies			
The following primary antibodies were used: chicken anti-GFP (1:500, ab-13970, Abcam, Cambridge, UK), glial fibrillary acidic protein (polyclonal, 1:1,000, ZO334, Dako-Agilent, Santa Clara, CA, USA monoclonal, 1:1,000, MAB 3402, Millipore, Burlington, MA, USA), rabbit anti Iba1 (1:100, 019-19741, Wako Pure Chemicals Industries, Neuss, Germany), rat anti CD68 (1:200, clone FA-11MCA1957, Bio-Rad, Hercules, CA, USA), rabbit anti LAMP1 (1:200, AB24170 Abcam), rat anti-LAMP1 hybridoma (1:300, 1D4B, Developmental Studies Hybridoma Bank at The University of Iowa), mouse anti-GM2 (1:500, A2576, TCI Europe), and rabbit anti-Calbindin (1:1,000, CB-38A, Swant, Burgdorf, Switzerland).	anti-GFP		
	glial fibrillary acidic protein		
	anti Iba1		Suggestion: (Synaptic Systems Cat# 234 308, RRID:AB_2924932)(link)
	anti CD68		
	anti LAMP1		
	anti-LAMP1		
	anti-GM2		
	anti-Calbindin		
The secondary antibodies were from Thermo Fisher Scientific : Alexa 488 (1:1,000, anti-rabbit A11008, anti-mouse A11001, and anti-chicken A11039), Alexa 546 (1:2,000, anti-mouse A11039 and anti-rabbit A11010), and Alexa 633 (1:500, anti-rat A21094 and anti-mouse A21050).	anti-rabbit		
	anti-chicken		
	anti-mouse		
	anti-rat		
Primary antibodies were as follows: rabbit anti-LAMP1 (1:100, AB24170, Abcam), rabbit anti-GFAP (1:100,000, ZO334, Dako), mouse anti-MBP (1:4,000, clone SMI 94SMI 99, BioLegend), rabbit anti-MAG (1:1,000, 34-6200, Thermo Fisher Scientific), rabbit anti-calnexin (1:5,000, C4731, Sigma-Aldrich, St. Louis, MO, USA), and mouse anti-actin (1:10,000 sc-1616r, Santa Cruz Biotechnology, Dallas, TX, USA).	anti-GFAP		
	anti-MBP		
	anti-MAG		
	anti-calnexin		
	anti-actin		
Secondary antibodies (Chemicon, Temecula, CA, USA) were as follows: goat anti-rabbit horseradish peroxidase (HRP) (1:10,000, AP132P) and goat anti-mouse HRP (1:10,000, AP124P).	anti-mouse HRP		
Experimental Models: Organisms/Strains			

CAG GFP transgenic mice (C57BL6-Tg (CAG-EGFP)1OsbJ) were purchased from The Jackson Laboratory (Bar Harbor, ME USA).	C57BL6-Tg (CAG-EGFP)1OsbJ		
BMT Total BM was flushed from the tibiae, femora, and humeri of 4- to 8-week-old tgGFP mice as described previously.	tgGFP		
Software and Algorithms			
Data were analyzed with Fiji software (ImageJ, National Institutes of Health, Bethesda, MD, USA), 94 LasX software (Leica Application Suite X, RRID: SCR_013673), and Imaris viewer software (Oxford Instruments, Abingdon-on-Thames, UK).	Fiji		
	ImageJ		Leica Application Suite X (RRID:SCR_013673)(link)
	Imaris		
Images were imported into Adobe Photoshop 2021 for brightness and contrast adjustments and to merge the channels.	Adobe Photoshop		Suggestion: Adobe Photoshop, RRID:SCR_014199(link)
All data were analyzed using the FlowJo software.phycoerytrin Morphological analysis of the SC and optic nerve Semithin section analysis of SCs was performed as reported previously.	FlowJo		Suggestion: FlowJo, RRID:SCR_008520(link)
Lipid bands were detected by digital acquisition and quantified using ImageJ software.	ImageJ		Suggestion: ImageJ, RRID:SCR_003070(link)
Statistics Data were analyzed with GraphPad Prism version 8.0 for Macintosh (GraphPad, San Diego, CA, USA) and expressed as the mean +- standard error of the mean (SEM).	GraphPad Prism		Suggestion: GraphPad Prism (RRID:SCR_002798)(link)
	GraphPad		Suggestion: GraphPad Prism, RRID:SCR_002798(link)

Other Entities Detected

Your Sentences	Recognized Entity
Statistical Tests	
Unpaired Students t test and one-way or two-way ANOVA followed by post-tests were used when appropriate (statistical significance: $p < 0.05$).	t test
	ANOVA
Survival curves were analyzed with log rank (Mantel-Cox) test.	Mantel-Cox) test
Unresolved RRIDs	
The following primary antibodies were used: chicken anti-GFP (1:500, ab-13970, Abcam, Cambridge, UK), glial fibrillary acidic protein (polyclonal, 1:1,000, ZO334, Dako-Agilent, Santa Clara, CA, USA monoclonal, 1:1,000, MAB 3402, Millipore, Burlington, MA, USA), rabbit anti Iba1 (1:100, 019-19741, Wako Pure Chemicals Industries, Neuss, Germany), rat anti CD68 (1:200, clone FA-11MCA1957, Bio-Rad, Hercules, CA, USA), rabbit anti LAMP1 (1:200, AB24170 Abcam), rat anti-LAMP1 hybridoma (1:300, 1D4B, Developmental Studies Hybridoma Bank at The University of Iowa), mouse anti-GM2 (1:500, A2576, TCI Europe), and rabbit anti-Calbindin (1:1,000, CB-38A, Swant, Burgdorf, Switzerland).	AB24170
Primary antibodies were as follows: rabbit anti-LAMP1 (1:100, AB24170, Abcam), rabbit anti-GFAP (1:100,000, ZO334, Dako), mouse anti-MBP (1:4,000, clone SMI 94SMI 99, BioLegend), rabbit anti-MAG (1:1,000, 34-6200, Thermo Fisher Scientific), rabbit anti-calnexin (1:5,000, C4731, Sigma-Aldrich, St. Louis, MO, USA), and mouse anti-actin (1:10,000 sc-1616r, Santa Cruz Biotechnology, Dallas, TX, USA).	AB24170,

SciScore is an automated tool that is designed to assist expert reviewers by finding and presenting formulaic information scattered throughout a paper in a standard, easy to digest format. ***SciScore is not a substitute for expert review.*** SciScore also checks for the presence and correctness of several unique identifiers, including RRIDs (research resource identifiers) in the manuscript, detects sentences that appear to be missing RRIDs, and can even suggest RRIDs under certain circumstances. **All RRID suggestions should be verified;** only the author can know whether the suggestions are correct.

For a full description of scored criteria and tips for improving your score, please see https://www.scicrunch.com/sciscorereport-faq