Document Identifier: **PMC8989626**

SciScore Report

Below you will find your SciScore report containing three tables. Your score is calculated based on adherence to scientific rigor criteria (Table 1) and identification of key biological resources (Table 2). Table 3 contains statistical tests and oligonucleotides but is not scored. If SciScore makes any mistakes, please contact us to help us learn and improve.

Table 1: Rigor Adherence Table

Ethics
IACUC: EXPERIMENTAL MODELS AND SUBJECT DETAILS Animal experiments All animal work and procedures were approved by the Northwestern University Institutional Animal Care and Use Committee.
Inclusion and Exclusion Criteria
Mice that developed extra-pancreatic diseases such as thymoma , prolapse , or inoperable papillomas were excluded from the study .
Attrition
not detected.
Sex as a biological variable
All mice were bred on a C57/BL6 background , with ages ranging from 3–20 months and from both genders .
Subject Demographics
Age: Primary cell lines from KCG+/+ (2005 , 7-month-old female; 2685 , 6-month-old male) , KCGfl/fl (1930 , 9-month-old male; 1999 , 10-month-old male) , KPCG+/+ (2138 , 3-month-old male; 3213 , 3-month-old female) , and KPCGfl/fl (3458 , 5-month-old female; 3566 , 4-month-old female) C57/BL6 mice were generated as described below .
Randomization
not detected.
Blinding
not detected.
Power Analysis
not detected.
Replication
not required.
Cell Line Authentication

Contamination: All cells in culture were tested and confirmed to be free of mycoplasma using the Myco Alert Plus Kit (Lonza LT07–701) .
Data Information
Availability: Data and code availability This paper analyzes publicly available data from the TCGA and cBioportal databases.

Table 2: Key Resources Table

Your Sentences	REAGENT or RESOURCE	SOURCE	IDENTIFIER
Antibodies			
Tissue sections were incubated with antibodies for: E-cadherin (Cell Signaling #3195 , RRID: AB_2291471 , 1:500) , Cytokeratin 19 (Abcam ab52625 , RRID: AB_2281020 , 1:500) , Ki67 (Cell Signaling #12202 , RRID: AB_2620142 , 1:1000)	E-cadherin	Cell Signaling Technology	(Cell Signaling Technology Cat# 3195, RRID:AB_2291471)(link)
	Cytokeratin 19		
	Cytokeratin	Abcam	(Abcam Cat# ab52625, RRID:AB_2281020)(link)
	Ki67		
	Ki67 (Cell	Cell Signaling Technology	(Cell Signaling Technology Cat# 12202, RRID:AB_2620142)(link)
, cleaved caspase-3 (Cell Signaling #9664 , RRID: AB_2070042 , 1:800)	cleaved caspase-3	Cell Signaling Technology	(Cell Signaling Technology Cat# 9664, RRID:AB_2070042)(link)
, mTOR (Cell Signaling #2983 , RRID: AB_2105622 , 1:1000) ,	mTOR	Cell Signaling Technology	(Cell Signaling Technology Cat# 2983, RRID:AB_2105622)(link)
p-mTOR (Abcam ab109268 , RRID: AB_10888105 , 1:2000) , RPS6 (Cell Signaling #2217 , RRID: AB_331355 , 1:1000) , and p-RPS6 (Cell Signaling #4858 , RRID: AB_916156 , 1:1000) overnight at 4°C .	p-mTOR	Abcam	(Abcam Cat# ab109268, RRID:AB_10888105)(link)
			(Cell Signaling Technology Cat# 2217, RRID:AB_331355)(link)
	p-RPS6	Cell Signaling Technology	(Cell Signaling Technology Cat# 4858, RRID:AB_916156)(link)
Antibody binding was detected using HRP-conjugated anti-rabbit secondary antibody and visualized using ImmPACT DAB Peroxidase Substrate kit (VectorLabs	anti-rabbit		
Tissue sections were incubated with a mixture of antibodies for CK19 (DSHB TROMA-III , RRID: AB_2133570,1:100) and Ki67 (Cell Signaling #12202 , RRID: AB_2620142 , 1:500) for 2 h at room temperature.	CK19		
			(DSHB Cat# TROMA-III, RRID:AB_2133570)(link)
After washing , tissue sections were incubated with a mixture of goat anti-rat Alexa Fluor 594 (ThermoFisher , RRID: AB_10561522 #A-11007 , 1:200) and goat anti-rabbit AlexaFluor 488Plus (ThermoFisher #A-32731 , RRID: AB_2633280 1:200) secondary antibodies and protected from light .	anti-rat	Thermo Fisher Scientific	(Thermo Fisher Scientific Cat# A-11007, RRID:AB_10561522)(link)
Primary antibodies used include Gα13 (Santa Cruz #sc-293424 , 1:1,000) , E-cadherin (Cell Signaling #3195 , RRID:AB_2291471 , 1: 5,000) , PDK1 (Cell Signaling #5662 , AB_10839264 , 1:1,000) ,	PDK1		
	PDK1 (Cell	Cell Signaling Technology	(Cell Signaling Technology Cat# 5662, RRID:AB_10839264)(link)
	p-PDK1		

p-PDK1 (Cell Signaling #3438 , RRID:AB_2161134 1:1,000			
HRP-conjugated rabbit (#A6667 , RRID: AB_258307 , 1:10,000) or mouse (#A4416 , RRID: AB_258167 , 1:10,000) secondary antibody (Millipore-Sigma St . Louis , MO) was used with SuperSignal West Pico PLUS (Thermo Fisher Scientific) for protein detection .	rabbit	Sigma-Aldrich	(Sigma-Aldrich Cat# A6667, RRID:AB_258307)(link)
	mouse	Sigma-Aldrich	(Sigma-Aldrich Cat# A4416, RRID:AB_258167)(link)
Experimental Models: Organisms/Strains			
The following mice were used in the study: Pdx1-Cre mice (Jackson Laboratory #014647), Gna13 (kindly provided by Stefan Offermans, Max Planck Institute) (Moers et al., 2003), LSL-KRasG12D/+ (Jackson Laboratory #019104	Pdx1-Cre		
	Gna13		
	LSL-KRasG12D/+		
Primary cell lines from KCG+/+ (2005 , 7-month-old female; 2685 , 6-month-old male) , KCGfl/fl (1930 , 9-month-old male; 1999 , 10-month-old male) , KPCG+/+ (2138 , 3-month-old male; 3213 , 3-month-old female) , and KPCGfl/fl (3458 , 5-month-old female; 3566 , 4-month-old female) C57/BL6 mice were generated as described below .	C57/BL6		Suggestion: RRID:IMSR_JAX:000664(link)
METHOD DETAILS Conditional knockout Mice with loss of Gα13 in the pancreas were generated by crossing Pdx1-Cre mice (Jackson Laboratory #014647) to mice expressing the floxed allele of Gna13 (kindly provided by Stefan Offermans , Max Planck Institute) (Moers et al. , 2003) , to generate CGα13fl/+ and CGα13fl/fl mice.	CGα13fl/fl		
) mutant allele to generate KCGα13fl/+ (KCGfl/+) or KCGα13fl/fl (KCGfl/fl) mice.	KCGα13fl/fl		
These mice were further crossed with mice expressing LSL-Trp53R172H/+ (Jackson Laboratory 008652) to generate KPCGfl/+ and KPCGfl/fl mice.	KPCGfl/fl		
Cell implantation For subcutaneous implantation , cells (2.5 × 105) were suspended in 100 µL of 1:1 mixture of PBS: Matrigel and then injected under the skin in the flank of C57BL/6 mice.	C57BL/6		Suggestion: RRID:IMSR_JAX:000664(link)

Generation of cell lines from genetically engineered mouse models KCG+/+ , KCGfl/+ , or KCGfl/fl mice more than 6 months of age were euthanized , and the harvested pancreatic tissue was placed in 5 mL of cold DMEM using aseptic technique.	KCGfl/fl		
Software and Algorithms			
Data and code availability This paper analyzes publicly available data from the TCGA and cBioportal databases.	cBioportal		Suggestion: cBioPortal (RRID:SCR_014555)(link)
At least 10 sections were acquired per tissue section , and ImageJ was used to analyze the double-positive CK19 and Ki67 areas .	ImageJ		Suggestion: ImageJ, RRID:SCR_003070(link)
All statistical analyses were done using GraphPad Prism.	GraphPad Prism		

Other Entities Detected

Your Sentences	Recognized Entity
Statistical Tests	
QUANTIFICATION AND STATISTICAL ANALYSIS The in vivo and in vitro results were compared using one-way ANOVA with Bonferroni correction and 2-tailed t test analysis.	ANOVA
	Bonferroni
	t test
Unresolved RRIDs	
After washing , tissue sections were incubated with a mixture of goat anti-rat Alexa Fluor 594 (ThermoFisher , RRID: AB_10561522 #A-11007 , 1:200) and goat anti-rabbit AlexaFluor 488Plus (ThermoFisher #A-32731 , RRID: AB_2633280 1:200) secondary antibodies and protected from light .	RRID: AB_2633280 1
Primary antibodies used include Gα13 (Santa Cruz #sc-293424 , 1:1,000) , E-cadherin (Cell Signaling #3195 , RRID:AB_2291471 , 1: 5,000) , PDK1 (Cell Signaling #5662 , AB_10839264 , 1:1,000) , p-PDK1 (Cell Signaling #3438 , RRID:AB_2161134 1:1,000	RRID:AB_2161134 1
Gapdh (Millipore Sigma #MAB374 , RRID: AB_2107445 1:5,000) , and Hsp90 (Santa Cruz Biotechnology sc-7947 , RRID: AB_2121235 1:5,000)	RRID: AB_2107445 1
	RRID: AB_2121235 1

SciScore is an automated tool that is designed to assist expert reviewers by finding and presenting formulaic information scattered throughout a paper in a standard, easy to digest format. ***SciScore is not a substitute for expert review.*** SciScore also checks for the presence and correctness of several unique identifiers, including RRIDs (research resource identifiers) in the manuscript, detects sentences that appear to be missing RRIDs, and can even suggest RRIDs under certain circumstances. **All RRID suggestions should be verified;** only the author can know whether the suggestions are correct.

For a full description of scored criteria and tips for improving your score, please see https://www.scicrunch.com/sciscorereport-faq