Document Identifier: **PMC8993640**

SciScore Report

Below you will find your SciScore report containing three tables. Your score is calculated based on adherence to scientific rigor criteria (Table 1) and identification of key biological resources (Table 2). Table 3 contains statistical tests and oligonucleotides but is not scored. If SciScore makes any mistakes, please contact us to help us learn and improve.

Table 1: Rigor Adherence Table

Ethics
IRB: 2 MATERIALS AND METHODS 2.1 Participants and plasma samples This study was approved by the Ethics Committee of The First Affiliated Hospital of Guangxi Medical University (No. 2021; KY-E-156).
Consent: Parental informed consent was obtained for this study for each subject.
Inclusion and Exclusion Criteria
The inclusion criteria were as follows: (1) age <14 years and (2) patient underwent cardiac catheterization before intervention.
Attrition
not detected.
Sex as a biological variable
not detected.
Subject Demographics
Age: The inclusion criteria were as follows: (1) age <14 years and (2) patient underwent cardiac catheterization before intervention.
Randomization
not detected.
Blinding
not detected.
Power Analysis
not detected.
Replication
not required.

Table 2: Key Resources Table

Your Sentences	REAGENT or RESOURCE	SOURCE	IDENTIFIER
Software and Algorithms			
The circMIR software was used to predict circRNA-miRNA interactions, based on the RNAhybrid (https://bibiserv.cebitec.uni-bielefeld.de/rnahybrid/) and miRanda (http://www.microrna.org/) databases.	circMIR		
	http://www.microrna.org/		
The miRWalk database (http://zmf.umm.uniheidelberg.de/apps/zmf/mirwalk/) was used to predict the target genes of miRNAs targeting hsa_circ_0003416, and the top 30 genes of each miRNA were selected.	miRWalk		Suggestion: miRWalk (RRID:SCR_016509)(link)
Gene Ontology (GO) and Kyoto Encyclopedia of Genes and Genomes (KEGG) analyses were conducted using the DAVID database (http://david.abcc.ncifcrf.gov), based on the selected target genes.	DAVID		Suggestion: DAVID (RRID:SCR_001881)(link)
The GO terms and KEGG pathways with p-value <0.05 were considered significantly enriched.	KEGG		Suggestion: KEGG (RRID:SCR_012773)(link)
Receiver operating characteristic (ROC) curves were established using SPSS 25.0 to assess the diagnostic values.	SPSS		Suggestion: SPSS, RRID:SCR_002865(link)

Other Entities Detected

Your Sentences	Recognized Entity
Statistical Tests	
Fisher's exact or chi-square test was used to test the categorical variables.	Fisher's exact
	chi-square test
Data normality was tested using the Shapiro–Wilk normality test.	Shapiro–Wilk normality test
If a quantitative variable conformed to a normal distribution, a Student's t-test or analysis of variance was applied; otherwise, the Mann–Whitney U-test or the Kruskal–Wallis H-test was used.	t-test
	analysis of variance
	Mann–Whitney U-test
	Kruskal–Wallis H-test
Correlation was calculated using Spearman correlation analysis.	Spearman correlation

SciScore is an automated tool that is designed to assist expert reviewers by finding and presenting formulaic information scattered throughout a paper in a standard, easy to digest format. ***SciScore is not a substitute for expert review.*** SciScore also checks for the presence and correctness of several unique identifiers, including RRIDs (research resource identifiers) in the manuscript, detects sentences that appear to be missing RRIDs, and can even suggest RRIDs under certain circumstances. **All RRID suggestions should be verified;** only the author can know whether the suggestions are correct.

For a full description of scored criteria and tips for improving your score, please see https://www.scicrunch.com/sciscorereport-faq