Document Identifier: **PMC8995061**

SciScore Report

Below you will find your SciScore report containing three tables. Your score is calculated based on adherence to scientific rigor criteria (Table 1) and identification of key biological resources (Table 2). Table 3 contains statistical tests and oligonucleotides but is not scored. If SciScore makes any mistakes, please contact us to help us learn and improve.

Table 1: Rigor Adherence Table

Ethics
not required.
Inclusion and Exclusion Criteria
not required.
Attrition
not required.
Sex as a biological variable
not detected.
Subject Demographics
Age: 4.3 Expression analysis of X . oryzae pv . oryzae effectors in planta TN1 rice plants (60 days old) were clip-inoculated with X . oryzae pv . oryzae BXO43 at OD600 = 1 .
Randomization
not detected.
Blinding
not detected.
Power Analysis
not detected.
Replication
Each set was repeated three times.
Number: Each set was repeated three times.
Data Information

Identifiers: solanacearum MolK2 is CAQ37632.1 .	CAQ37632.1

Table 2: Key Resources Table

Your Sentences	REAGENT or RESOURCE	SOURCE	IDENTIFIER
Antibodies			
Western blot analysis was performed using alkaline phosphatase-conjugated anti-His antibody (Sigma Aldrich) and anti-GFP antibody (Abcam).	anti-His		
	anti-GFP		
For immunoblotting, alkaline phosphatase conjugated to anti-rabbit immunoglobulin G secondary antibody (Sigma Aldrich), 4-nitroblue tetrazolium (NBT Roche), and 5-bromo-4-chloro-3-indolyl-phosphate 4-toluidine salt (BCIP Roche) were used.	anti-rabbit immunoglobulin G		
Recombinant DNA			
The pENTR::xopG plasmid was used as template.	pENTR::xopG		
The xopG gene of X. oryzae pv. oryzae was disrupted by homologous plasmid integration using the vector pK18mob (Schafer et al., 1994).	pK18mob		
To complement xopG -, the wild-type xopG gene, xopG E76A, or xopG E85A were cloned into the HindIII andKpnI sites of the pHM1 vector (Innes et al., 1988) using gene-specific primers (	pHM1		
4.6 Yeast two-hybrid assays Wild-type xopQ , xopQ T222A , or xopX cloned in the yeast two-hybrid vector pDEST32 (Invitrogen) was used from a previous study (Deb et al. , 2019 , 2020) . xopU , xopV , xopP , xopG , xopG E76A , xopG E85A , and avrBs2 were cloned in the yeast two-hybrid vector pDEST22 (Invitrogen) by Gateway cloning (Invitrogen) .	pDEST32		
	pDEST22		
4.7 BiFC The wild-type copy of xopQ, xopQ T222A, or xopX cloned in the BiFC vector pDEST-VYNE(R)GW carrying nVFP (Gehl et al., 2009) was used from a previous study (Deb et al., 2019, 2020). xopU, xopV, xopP, xopG, xopG E76A, xopG E85A, and avrBs2 were cloned in the BiFC vector pDEST-VYCE(R)GW (Gehl et al., 2009) carrying the C-terminal region of VFP (cVFP) by Gateway cloning	pDEST-VYNE(R)GW		
	pDEST-VYCE(R)GW		

(Invitrogen) to yield the constructs as listed in Table S2.			
Briefly, xopU, xopV, and xopG were cloned with the EGFP tag in the pH7WGF2 vector.	pH7WGF2		
EGFP::gus was used from a previous study as a control (Deb et al., 2020). xopX was cloned with a 6xHis tag in the pMDC7 vector (Table S2). pMDC7::xopQ-6xHis was used from a previous study (Deb et al., 2020).	pMDC7		
	pMDC7::xopQ-6xHis		
Software and Algorithms			
The obtained sequences were subjected to homology searches using the BLAST algorithm in the National Centre for Biotechnology Information database (Altschul et al., 1990).	BLAST		Suggestion: WU-BLAST (RRID:SCR_011824)(link)
Quantitative PCR was performed using Power SYBR Green Master Mix (Thermo Fisher Scientific) in a CFX384 RT-PCR instrument (Bio-Rad Laboratories) .	Bio-Rad Laboratories		Suggestion: Bio-Rad Laboratories (RRID:SCR_008426)(link)
4.4 Bioinformatic analysis of XopG Multiple sequence alignment of XopG from various bacterial strains was carried out using Clustal Omega (Madeira et al. , 2019) and visualized using ESPript software (Robert & Gouet , 2014) .	Clustal Omega		Suggestion: Clustal Omega (RRID:SCR_001591)(link)
Images were analysed using ZEN software.	ZEN		
Quantification of PI internalization was carried out using ImageJ software (Schneider et al., 2012).	ImageJ		Suggestion: ImageJ, RRID:SCR_003070(link)

Other Entities Detected

Your Sentences	Recognized Entity
Statistical Tests	
Statistical analysis for significance was conducted using the unpaired two-sided Students t test.	Students t test

SciScore is an automated tool that is designed to assist expert reviewers by finding and presenting formulaic information scattered throughout a paper in a standard, easy to digest format. ***SciScore is not a substitute for expert review.*** SciScore also checks for the presence and correctness of several unique identifiers, including RRIDs (research resource identifiers) in the manuscript, detects sentences that appear to be missing RRIDs, and can even suggest RRIDs under certain circumstances. **All RRID suggestions should be verified;** only the author can know whether the suggestions are correct.

For a full description of scored criteria and tips for improving your score, please see https://www.scicrunch.com/sciscorereport-faq