Document Identifier: **PMC8996088**

SciScore Report

Below you will find your SciScore report containing three tables. Your score is calculated based on adherence to scientific rigor criteria (Table 1) and identification of key biological resources (Table 2). Table 3 contains statistical tests and oligonucleotides but is not scored. If SciScore makes any mistakes, please contact us to help us learn and improve.

Table 1: Rigor Adherence Table

Ethics
IRB: Ethics statement This study was approved by the Scientific Research Ethics Committee of Near East University (Research Project Evaluation Report No: 2020/85-1188).
Consent: Written informed consent was obtained from the participants.
Inclusion and Exclusion Criteria
While the study was in progress, 7 patients were excluded because 5 of these patients had previously undetected type 2 diabetes mellitus and 2 had hypertension.
Attrition
not detected.
Sex as a biological variable
not detected.
Subject Demographics
Age: A total of 112 men between the ages of 22 and 45 years, who had been hunting regularly (20 times a year, with an average of 10 shots per day) since the age of 18, were included in the study.
Randomization
not detected.
Blinding
not detected.
Power Analysis
not detected.
Replication
not required.

Table 2: Key Resources Table

Your Sentences	REAGENT or RESOURCE	SOURCE	IDENTIFIER
Software and Algorithms			
Demographic characteristics of individuals such as age, gender, occupation, and audiometry results were recorded in an Excel format, and data were analyzed using SPSS V23 (IBM Corp., Armonk, NY, USA).	SPSS		Suggestion: SPSS, RRID:SCR_002865(link)

Other Entities Detected

Your Sentences	Recognized Entity
Statistical Tests	
Conformity to a normal distribution was evaluated using the Kolmogorov-Smirnov and Shapiro-Wilk tests.	Kolmogorov-Smirnov
	Shapiro-Wilk tests
The chi-square test was used to compare categorical variables across groups.	chi-square test
Independent two-sample t-tests were used to compare normally distributed data according to paired groups, and the Mann-Whitney U test was used to compare non-normally distributed data.	t-tests
	Mann-Whitney U test
One-way analysis of variance was used to compare normally distributed quantitative data according to groups of three or more, and the Kruskal-Wallis test was used to compare non-normally distributed data.	analysis of variance
	Kruskal-Wallis test
Spearman's rho correlation coefficient was used to examine the relationship between the non-normally distributed data.	Spearman's rho correlation

SciScore is an automated tool that is designed to assist expert reviewers by finding and presenting formulaic information scattered throughout a paper in a standard, easy to digest format. ***SciScore is not a substitute for expert review.*** SciScore also checks for the presence and correctness of several unique identifiers, including RRIDs (research resource identifiers) in the manuscript, detects sentences that appear to be missing RRIDs, and can even suggest RRIDs under certain circumstances. **All RRID suggestions should be verified;** only the author can know whether the suggestions are correct.

For a full description of scored criteria and tips for improving your score, please see https://www.scicrunch.com/sciscorereport-faq