

CURRENT STATUS OF FAT

P. KAMPHUIS

1. INTRODUCTION

This document provides an overview of the current version of **FAT** related to the tests provided in Kamphuis et al. (2015). It merely consist of a set of figures which are described in said paper. Typically the current version works much better then the version from 2015. However, as we have also reduced the applied errors these plot show that performance is comparable, i.e. the fits are equally accurate with smaller errors. As **FAT** now employs a Monte Carlo method in the regularisation of the parameters the final model can sometimes differ slightly. However this should always be within the calculated errors. If not please open an issue on the GitHub. Also, if you are looking for specific functionality or find that **FAT** is not performing well despite the galaxy having regular rotation please do not hesitate to contact me.

It is important to remember that **FAT** is truly meant for batch fitting. Hence the aim of the code is to provide tilted ring models that are accurate in a large fraction of galaxies. Ideally, **FAT** should identify galaxies that are not fitted well however this feature is not optimal yet. When fitting individual galaxies it is recommended to run **FAT** and then fine tune the model by hand. In most cases such finetuning will be limited to a few outer rings but in the case of complex galaxies with significant non-cylindrically symmetric motions the models can fail.

FAT is not a replacement of the extended Functionality of **TiRiFiC**. **FAT** fits simple rotationally symmetric discs with assymetric warps and surface brightness distributions. However, **TiRiFiC** has a much more extended functionality and should be used for identifying significant non-cylindrically symmetric motions, thick discs, bars and other such Hifeatures. When modelling such galaxies ideally **FAT** provides a base model with merely a thin disc and subsequently other features are added with **TiRiFiC**. For examples of such modelling please see Kamphuis et al. (2011), Zschaechner et al. (2011), Kamphuis et al. (2013), Gentile et al. (2013).

2. LVHIS GALAXIES

This section shows Figures 7, 11 and 12 from Kamphuis et al. (2015).




FIGURE 1. Differences between the final fits with **FAT**, ROTCUR and DISKFIT. Left hand side of each panel: the difference between ROTCUR and DISKFIT, i.e. ROTCUR - DISKFIT, Right hand side of each panel: blue: the difference between ROTCUR and **FAT**, red: the difference between DISKFIT and **FAT**. The solid grey lines show the weighted average and the dashed lines show the 1σ deviation from this average in each plot. Stars correspond to final fits outside the previously determined reliable range (see Section ??) and circles to ones inside these limits. The errors on the points are calculated as the average of the error for all rings divided by \sqrt{N} with N the number of rings in the model. This is Figure 7 from Kamphuis et al. (2015).




FIGURE 2. Comparison between the PA and inclination of the optical values and the different fits, black - FAT, blue - ROTCUR, red - DISKFIT. Stars are galaxies where the final FAT model is in a part of parameter space that is considered unreliable. The grey dash line denote the scatter on the deviations between literature and the ROTCUR interactive fit. This is Figure 11 from Kamphuis et al. (2015).

3. ARTIFICIAL GALAXIES

This shows the results on the set of Artificial Database as described in Kamphuis et al. (2015).




FIGURE 3. Rotation curves for all the galaxies in the LVHIS Sample. The plots are ordered according to their **FAT** inclination, with the lowest inclination in the bottom left. Black lines: **FAT**, Red Lines: **DISKFIT**, Blue lines: **ROTCUR**. The error bars are the formal errors derived by **DISKFIT** and **ROTCUR** for the red and blue lines respectively. For **FAT** the error is an empirical estimate derived from the regularisation process (see § ??). Stars indicate curves outside the reliably fitted range. This is Figure 12 from Kamphuis et al. (2015).




FIGURE 4. Rotation curves used and fitted in the artificial galaxy database. Left panel: dwarf galaxy and intermediate galaxy. Right panel: massive galaxy. The black circles and lines indicate the input rotation curves, grey solid lines are the reliable output curves, grey dashed lines are from artificial galaxies outside the reliable fitting range. Figure 4 in Kamphuis et al. (2015)

4. EXTENDED ARTIFICIAL GALAXIES

This shows the results on an extended set of Artificial Database as described in Kamphuis et al. (2015). Typically these galaxies are warped, as opposed to flat like the artificial database of the paper, and they explore a range of different velocity resolutions. This addition of the database was made to show that the accuracy of the code should really be expressed in beams and channels and not in arcsec and km s^{-1} . Table 1 is a copy of Table one in Kamphuis et al. (2015) for this extended set of galaxies.




FIGURE 5. Difference between the artificial galaxy inputs and **FAT** output. On the y-axis the difference between the model input and best fitting **FAT** values is shown for various parameters. The solid line shows the mean deviation over the whole sample and the dashed lines show the 1σ offsets from this average. The errors on the points are calculated as the average of the **FAT** error for all rings divided by \sqrt{N} , where N is the number of rings in the model. The stars show galaxies outside the reliable range for **FAT** whereas circles are inside this range. The colour coding indicates: blue - dwarf galaxy, black - intermediate galaxy, red - massive galaxy. Figure 6 in Kamphuis et al. (2015)

Parameter	Database Galaxies		Variations
	Base 1	Base 2	Sample Values
Beams across the major axis	16	16	32
Inclination ($^{\circ}$)	60	50	-
PA ($^{\circ}$)	45	55	-
Inner-Outer Dispersion (km s^{-1})	8.0-8.0	8.0-8.0	-
Inner-Outer $\Delta \theta$ (rad)	0-0.05	0-0.05	0-0.15
Inner-Outer $\Delta \phi$ (rad)	0-0.2	0-0.2	0-0.05
Inner-Outer scale height (kpc)	0.3-0.3	0.2-0.2	-
Channel width (km s^{-1})	2	2	4, 8
Signal-to-Noise	8	8	-
Rotation Curve	intermediate	massive	-

TABLE 1. Values for the two base galaxies (columns 2 & 3) and the values by which they are varied (column 4). We change only one parameter at a time while all other parameters for the base galaxies are unchanged.




FIGURE 6. Rotation curves used and fitted in the artificial galaxy database. Left panel: dwarf galaxy and intermediate galaxy. Right panel: massive galaxy. The black circles and lines indicate the input rotation curves, grey solid lines are the reliable output curves, grey dashed lines are from artificial galaxies outside the reliable fitting range. Figure 4 in Kamphuis et al. (2015)




FIGURE 7. Difference between the artificial galaxy inputs and [FAT](#) output. On the y-axis the difference between the model input and best fitting [FAT](#) values is shown for various parameters. The solid line shows the mean deviation over the whole sample and the dashed lines show the 1σ offsets from this average. The errors on the points are calculated as the average of the [FAT](#) error for all rings divided by \sqrt{N} , where N is the number of rings in the model. The stars show galaxies outside the reliable range for [FAT](#) whereas circles are inside this range. The colour coding indicates: blue - dwarf galaxy, black - intermediate galaxy, red - massive galaxy. Figure 6 in Kamphuis et al. (2015)