

# The Book on Modal Triplet Theory

A Typed Interpretive Overview of the Current Program

What is proved, reconstructed, calibrated, and still open

Peter Nero

July 2026

Version 10

# Abstract and Current Status

Modal Triplet Theory (MTT) is a proposed architecture for spectral reduction, coherent-sector stability, and comparison among physical encodings. Its rigorous core is a dimension-neutral Hilbert-bundle framework with compatible vertical operators, a joint spectral projector, separately stated fixed-point and equilibrium gates, controlled complementary-mode estimates, and typed descent and recovery maps. A canonical physical realization uses a supplied ten-dimensional bundle  $M_{10} \rightarrow Y_4$  with compact six-dimensional fiber  $X_6$  and a Lorentzian four-dimensional base.

The current geometry program combines a local rank-three comparison field  $Q_{\text{WW}} \in \Gamma(\text{Hom}(TP, TI))$  with a selected q79 degree-three spectral-cover carrier. Both exhibit a  $1 + 2 + 3$  rank profile, but their global bundle-and-connection intertwiner remains to be proved. The q79 branch has an exact trace split and a local binary-dihedral spin lift; strict global Spin closure is still an obstruction calculation.

Computationally, MTT closes embedded renormalized-Standard-Model equivalence at the declared one-shared-physical-primitive/profile standard. This includes the finite matrix architecture, charged-Yukawa profile, CKM, electroweak, threshold, finite-Dirac, and multi-loop precision packets. It is not strict zero-knob selection of the observed universe. Standard BRST/Faddeev–Popov quantization and measured renormalized profile data are admitted at this closure tier.

This book is interpretive, but every interpretation is labeled. It does not claim that MTT has derived the Born rule, Einstein gravity, a UV-complete quantum gravity, the origin of time, three families, cosmology, or all of string theory from projection alone.

# Revision Note for This Edition

**Supersedes.** *The Book on Modal Triplet Theory*, version 9 (January 2026).

**Reason.** Version 9 treated triplet minimality, ten-dimensional geometry, particle and force assignments, basin/Born probabilities, Gaussian quantum gravity, framework containment, and cosmological scenarios as more established than the underlying papers allowed. It also used the retired few-TeV execution chain, conflated the shared phase circle with time-like bookkeeping, and predated the selected q79, finite-algebra, HYM, SM-profile, and foundational- geometry successor theorems.

**Resolution.** Version 10 is rebuilt around the corrected Foundation, Fixed Points I–VI, Projection–Admissibility, Signature, Scale, Relationship Atlas, Theta/Execution, q79, proto-spinor, closure-strain, action, and current SM ledgers. It uses explicit claim tiers throughout, corrects the  $3 + 3$  versus  $3 \times 3$  distinction, replaces literal circle–lens–nil topology by typed carrier roles, and reports numerical closure at its declared profile standard.

**Retained result.** The central interpretive idea survives: one can study stable effective descriptions through shared spectral, geometric, and fixed-point structures, and several nontrivial MTT realizations now have exact or reproducible certificates.

**Remaining boundary.** The leading unresolved promotions are the strict global q79 Spin certificate, the world-in-world/strain-to-q79 same-source intertwiner, a selected normalized action source, strict no-knob and unique- branch selection, and independent MTT derivations of quantum probability, gauge quantization, gravity, and cosmology.

**Major withdrawals from version 9.** The following are not carried forward as results: universal proof that three filters or three families are inevitable; identification of compact phase with physical time; literal  $S^1 \times \text{Lens} \times \text{Nil}$  compactification; Born probabilities from basin volume alone; instantaneous bilocal microcausality; all-loop finite and unitary quantum gravity from a Gaussian filter; a derived Big-Bang replacement; and the old 4.2–5 TeV cross-sector numerical chain.

# Contents

|                                                             |            |
|-------------------------------------------------------------|------------|
| <b>Abstract and Current Status</b>                          | <b>ii</b>  |
| <b>Revision Note for This Edition</b>                       | <b>iii</b> |
| <b>I How to Read the Current MTT Program</b>                | <b>1</b>   |
| <b>1 Ambition with a Type System</b>                        | <b>2</b>   |
| 1.1 Seven claim types . . . . .                             | 2          |
| 1.2 The current one-paragraph verdict . . . . .             | 2          |
| 1.3 Why this narrower statement is stronger . . . . .       | 3          |
| <b>2 What Is Assumed</b>                                    | <b>4</b>   |
| <b>II The Rigorous Internal Spine</b>                       | <b>5</b>   |
| <b>3 Hilbert Bundles, Vertical Operators, and Coherence</b> | <b>6</b>   |
| 3.1 Dimension-neutral architecture . . . . .                | 6          |
| 3.2 Three parameters that must not be merged . . . . .      | 6          |
| 3.3 Complementary-mode control . . . . .                    | 7          |
| <b>4 Fixed Points Are Not All the Same</b>                  | <b>8</b>   |
| 4.1 The six Fixed-Point papers . . . . .                    | 8          |
| <b>5 Projection, Descent, and Recovery</b>                  | <b>10</b>  |
| <b>III Geometry: Local Carrier and Global Candidate</b>     | <b>11</b>  |
| <b>6 The Canonical <math>4 + 6</math> Realization</b>       | <b>12</b>  |
| 6.1 Signature comes from a principal symbol . . . . .       | 12         |
| <b>7 World-in-World Comparison Geometry</b>                 | <b>13</b>  |
| 7.1 Iwasawa and the nil distinction . . . . .               | 13         |

|                                                                          |               |
|--------------------------------------------------------------------------|---------------|
| <b>8 The Selected q79 Carrier</b>                                        | <b>15</b>     |
| 8.1 Local spin lift and global obstruction . . . . .                     | 15            |
| 8.2 Which global topology is current? . . . . .                          | 16            |
| <b>9 The Missing Geometry Theorem</b>                                    | <b>17</b>     |
| <br><b>IV From Carrier to Physical Models</b>                            | <br><b>18</b> |
| <b>10 Proto-Spinors Without Automatic Particles</b>                      | <b>19</b>     |
| <b>11 Closure Strain, Scalars, and Mass</b>                              | <b>20</b>     |
| <b>12 A Regime-Local Action, Not a Derived Final Action</b>              | <b>21</b>     |
| <br><b>V The Standard Model Result at Its Actual Tier</b>                | <br><b>22</b> |
| <b>13 Finite Algebra and Matrix Architecture</b>                         | <b>23</b>     |
| <b>14 What “True SM Closure” Means Here</b>                              | <b>24</b>     |
| 14.1 Parameters and knobs . . . . .                                      | 24            |
| 14.2 Neutral sector and strong CP . . . . .                              | 25            |
| <b>15 Theta Closure and the End of the Few-TeV Chain</b>                 | <b>26</b>     |
| <br><b>VI Quantum, Field-Theoretic, and Gravitational Interpretation</b> | <br><b>27</b> |
| <b>16 Quantum Mechanics: What Is and Is Not Reconstructed</b>            | <b>28</b>     |
| <b>17 QFT and the Renormalized SM Functor</b>                            | <b>29</b>     |
| <b>18 Worldsheets, Strings, and Fu–Yau Geometry</b>                      | <b>30</b>     |
| <b>19 General Relativity and Gravity</b>                                 | <b>31</b>     |
| <b>20 Quantum Gravity: Present Boundary</b>                              | <b>32</b>     |
| <b>21 Time, Entropy, and Cosmology</b>                                   | <b>33</b>     |
| <br><b>VII Relationships to Established Frameworks</b>                   | <br><b>34</b> |
| <b>22 A Typed Atlas, Not a Universal Superset</b>                        | <b>35</b>     |

|             |                                                       |           |
|-------------|-------------------------------------------------------|-----------|
| <b>VIII</b> | <b>What Has Been Achieved and What Comes Next</b>     | <b>37</b> |
| <b>23</b>   | <b>The Genuine Achievements</b>                       | <b>38</b> |
| 23.1        | Methodological . . . . .                              | 38        |
| 23.2        | Functional analytic . . . . .                         | 38        |
| 23.3        | Geometric . . . . .                                   | 38        |
| 23.4        | Computational . . . . .                               | 38        |
| 23.5        | Epistemic . . . . .                                   | 38        |
| <b>24</b>   | <b>The Decisive Frontier</b>                          | <b>40</b> |
| 24.1        | Global q79 spin decision . . . . .                    | 40        |
| 24.2        | World-in-world to q79 intertwiner . . . . .           | 40        |
| 24.3        | One selected action source . . . . .                  | 40        |
| 24.4        | Strict value and branch selection . . . . .           | 40        |
| 24.5        | Probability and quantization . . . . .                | 40        |
| 24.6        | Gravity and cosmology . . . . .                       | 41        |
| <b>25</b>   | <b>Falsifiability Without Promotional Predictions</b> | <b>42</b> |
| <b>26</b>   | <b>An Interpretive Picture That Survives</b>          | <b>43</b> |
| <b>27</b>   | <b>Afterword</b>                                      | <b>44</b> |
| <b>A</b>    | <b>Current Claim Ledger</b>                           | <b>45</b> |
| <b>B</b>    | <b>Notation and Guardrails</b>                        | <b>47</b> |
|             | <b>References</b>                                     | <b>48</b> |

## Part I

# How to Read the Current MTT Program

# Chapter 1

## Ambition with a Type System

MTT remains ambitious. It asks whether stable effective physics can be organized by one mathematical language of vertical spectra, coherent projection, admissible continuation, and controlled reduction. The ambition does not authorize every attractive interpretation. The first discipline of the current program is therefore a type system for claims.

### 1.1 Seven claim types

**Assumption.** Data inserted into a realization: a manifold, metric, operator, representation, measure, action, or boundary condition.

**Conditional theorem.** A proved consequence of explicitly listed hypotheses on a stated domain.

**Characterization.** A necessary form for any object satisfying the premises, without proving existence or selection.

**Reconstruction.** Recovery of a known framework after target-compatible data are supplied.

**Controlled reduction.** A map from one description to another with a domain and error estimate.

**Calibration or profile equivalence.** Agreement using measured or target-related data, including branch or scheme selection.

**Held-out prediction.** A value not used in construction, normalization, scale choice, branch choice, or parameter fitting.

Interpretive correspondence is useful but is not an eighth theorem type. It is a proposed reading of already typed mathematics.

### 1.2 The current one-paragraph verdict

The internal spectral/control spine is rigorous under its hypotheses. The selected q79 and finite-algebra branches contain substantial exact structure. The Standard Model is recon-

structed at a declared profile-equivalence standard with one shared physical primitive. MTT has not yet shown that this branch and these profile values are uniquely selected from a parameter-free microscopic source. It is therefore more than a metaphor and less than a completed fundamental theory.

### 1.3 Why this narrower statement is stronger

A broad narrative can always absorb another analogy. A typed program can fail. If an operator does not commute, a projector cannot be multiplied as claimed. If a relator closes with central sign  $-1$ , a strict Spin lift fails. If normalized overlap rows do not match, the same-source bridge fails. If a registered held-out observable misses its uncertainty band, the numerical branch fails. The revised program gains credibility by creating such points of contact.

## Chapter 2

# What Is Assumed

No universal physical conclusion follows from “coherence” alone. A concrete MTT model declares at least:

1. the base, internal bundle, regularity class, and operator domains;
2. a strongly commuting family of vertical operators, or one total internal operator, and its selected spectral cluster;
3. a well-posed stabilization flow and an invariant domain;
4. the relation, if any, between stabilization, physical time, and RG scale;
5. a physical action or evolution law, state space, observables, and probability rule when physical claims are made; and
6. numerical provenance: sourced inputs, fitted inputs, branch selection, uncertainty, and held-out outputs.

These declarations are not blemishes. Every physical theory has input structure. The real question is which inputs can later be related or selected by a common theorem.

## Part II

# The Rigorous Internal Spine

## Chapter 3

# Hilbert Bundles, Vertical Operators, and Coherence

### 3.1 Dimension-neutral architecture

Let  $Y$  be a smooth base and  $\mathcal{E} \rightarrow Y$  a real or complex Hilbert bundle. On a declared Sobolev space

$$\mathcal{H} = H^s(Y; \mathcal{E}),$$

MTT uses three compatible vertical structures

$$(\mathcal{E}_i, A_i, P_i), \quad i = 1, 2, 3.$$

The  $A_i$  are nonnegative self-adjoint operators with specified domains. If they strongly commute, their spectral projectors may be multiplied:

$$P_{\text{coh}} = P_1 P_2 P_3, \quad Q_{\text{inc}} = I - P_{\text{coh}}.$$

Otherwise one must define a single total internal operator and take its spectral projector. Commutation is a hypothesis or theorem, never a gift of notation.

The triplet is not automatically three coordinate manifolds. Its entries may be operators, bundle summands, representation lanes, or filtrations on one internal carrier.

### 3.2 Three parameters that must not be merged

The corrected Foundation distinguishes:

$$\tau = \text{stabilization parameter}, \quad t = \text{physical time}, \quad \mu = \text{renormalization scale}.$$

A heat or relaxation semigroup  $R_\tau$  is not a Lorentzian propagator  $U(t_2, t_1)$ , and neither is an RG trajectory. A realization may relate them, but it must provide the map, dimensions, domain, and error.

### 3.3 Complementary-mode control

On the noncoherent sector one seeks a semigroup estimate of the form

$$\|e^{\tau L_Q}\| \leq M e^{-\gamma\tau}, \quad \gamma > 0,$$

or a suitable dissipativity bound. This form remains meaningful for nonnormal generators. Eigenvalue signs alone do not control transient growth. An internal gap controls internal complementary modes; it is not automatically a four-dimensional momentum cutoff.

## Chapter 4

# Fixed Points Are Not All the Same

Let  $R_\tau$  be a well-posed stabilization map and define a projected step

$$T_\tau = P_{\text{coh}} R_\tau.$$

A point satisfying  $T_\tau \Psi_* = \Psi_*$  is a projected time-step fixed point. It is not automatically a steady solution of the upper flow. A strict Lyapunov identity or another stationarity argument is required to promote it to an equilibrium.

Banach’s theorem supplies existence and uniqueness only when  $T_\tau$  is a strict contraction on a nonempty complete invariant domain. Schauder and Darbo provide different existence routes and do not imply uniqueness. This is why the old slogan “one inequality selects one universe” has been retired. The current theory uses a ledger of independent gates.

### 4.1 The six Fixed-Point papers

|        |                                                                                                                                                   |
|--------|---------------------------------------------------------------------------------------------------------------------------------------------------|
| FP I   | Geometry, Sobolev spaces, spectral projectors, semigroup well-posedness, conditional existence and uniqueness routes, and Galerkin approximation. |
| FP II  | Application to a supplied 4 + 6 control realization, with a shared circle counted as bundle data and equilibrium promotion kept separate.         |
| FP III | Deterministic disturbance amplitudes, stochastic noise powers, noncoherent damping floors, exact OU statements, and conditional homogenization.   |
| FP IV  | Curved spectral-cluster persistence, <i>QRP</i> leakage, intrinsic Karcher centroids, first-order modulation, and work-limited exits.             |
| FP V   | Frozen linear covariance, canonical-correlation estimates, declared admissibility margins, and Gaussian exit bounds.                              |
| FP VI  | A typed synthesis separating inherited theorem, conditional physical completion, and interpretation.                                              |

These results constitute the strongest general mathematical achievement of MTT. They do not select a particle spectrum or fundamental action by themselves.

## Chapter 5

# Projection, Descent, and Recovery

Suppose  $P_0 : X \rightarrow Y_0$  is an initial reduction,  $P_t : X \rightarrow Y_t$  a final reduction, and  $\Phi_t : X \rightarrow X$  an upper evolution. Four questions must be separated:

1. Does a right section choose one compatible upper representative?
2. Does a left decoder recover the actual upper state?
3. Does an autonomous reduced map  $F_t : Y_0 \rightarrow Y_t$  exist?
4. Is that reduced map injective, or do effective labels merge?

Noninjectivity of  $P_0$  does not forbid a right section. Autonomous descent exists exactly when  $P_t \Phi_t$  is constant on the equivalence classes of  $P_0$ . Probability requires a measure and disintegration; entropy requires a state or measure plus a dynamical or coarse-graining law. Neither appears from noninjectivity alone.

## Part III

# Geometry: Local Carrier and Global Candidate

## Chapter 6

# The Canonical 4 + 6 Realization

The canonical physical specialization is

$$\pi : M_{10} \longrightarrow Y_4, \quad \dim Y_4 = 4, \quad \dim X_x = 6.$$

The physical completion supplies a globally hyperbolic Lorentzian base and a compact Riemannian internal fiber. This is a selected realization, not a dimension theorem of the abstract Hilbert-bundle framework.

If the internal space factorizes, the dimensions of its factors must add to six. A shared phase circle is represented by a principal  $U(1)$  bundle or Hermitian line bundle. It is not appended as a seventh coordinate.

### 6.1 Signature comes from a principal symbol

A Gram tensor

$$G_{\mu\nu} = \langle D_\mu \Psi, D_\nu \Psi \rangle$$

is positive semidefinite and cannot be a Lorentzian metric. Causal signature is read from the principal symbol of a selected hyperbolic physical equation, schematically

$$\sigma_{\text{pr}}(x, \xi) = g^{\mu\nu}(x) \xi_\mu \xi_\nu I + \text{gauge blocks}.$$

The current signature theorem establishes compatibility and stability of one- time hyperbolicity. It does not derive three spatial dimensions.

## Chapter 7

# World-in-World Comparison Geometry

The lawful version of the world-in-world idea is a field between two oriented rank-three bundles:

$$Q_{\text{WW}} \in \Gamma(\text{Hom}(TP, TI)).$$

In local frames it is a  $3 \times 3$  matrix with nine components. This is not manifold-dimension multiplication. A rank-three bundle over a three-dimensional base has six-dimensional total space.

At an invertible background, polar decomposition separates orientation and strain:

$$\text{Mat}(3, \mathbb{R}) = \mathfrak{so}(3) \oplus \text{Sym}(3, \mathbb{R}), \quad 9 = 3 + 6.$$

Once an orthonormal flag is selected, the strain sector splits as

$$\text{Sym}(3, \mathbb{R}) = \mathbb{R}I_3 \oplus \mathcal{D}_0 \oplus \mathcal{S}_{\text{off}}, \quad 6 = 1 + 2 + 3.$$

The scalar, traceless-diagonal, and symmetric off-diagonal projectors are orthogonal for the Frobenius metric. The  $1 + 2 + 3$  split depends on the selected flag; the fully  $SO(3)$ -invariant split is  $1 + 5$ .

Therefore

$$1 + 3 \times 3 = (1 + 3) + (1 + 2 + 3) = 4 + 6 = 10$$

is an exact component identity after an ordering scalar is supplied. It is not a proof that these components are the tangent directions of  $M_{10}$ .

### 7.1 Iwasawa and the nil distinction

For  $GL^+(3, \mathbb{R})$ , Iwasawa decomposition has dimensions

$$\dim SO(3) = 3, \quad \dim A = 3 = 1 + 2, \quad \dim N = 3.$$

The upper-unitriangular  $N$  is the three-dimensional Heisenberg nil group. The symmetric off-diagonal strain space is also three-dimensional, but is not itself that Lie algebra. They are related by a background-dependent local coordinate map. This distinction prevents dimension matching from becoming an unproved dynamical identity.

## Chapter 8

# The Selected q79 Carrier

For the selected degree-three spectral cover  $\pi : C \rightarrow B$ , let

$$\mathcal{A} = \pi_* \mathcal{O}_C, \quad \mathcal{A}_0 = \ker \text{Tr}.$$

The normalized trace projector gives

$$\mathcal{A} = \mathcal{O} \oplus \mathcal{A}_0, \quad \text{rank}(\mathcal{O}, \mathcal{A}_0, \mathcal{A}) = (1, 2, 3).$$

The common carrier is

$$\mathcal{H}_{\text{q79}} = L_{\text{shared}} \otimes (\mathcal{O} \oplus \mathcal{A}_0 \oplus \mathcal{A}), \quad \text{rank } \mathcal{H}_{\text{q79}} = 6.$$

This is a direct trace/trace-zero/full filtration. A connected transitive cover does not provide a global ordering of three individual sheets.

The q79 arithmetic branch includes the exact selected residue construction leading to  $q = 79$  modulo the declared order-448 quotient. Its value is a branch theorem with stated assumptions, not a replacement for the geometry selecting every physical observable.

### 8.1 Local spin lift and global obstruction

The signed sheet representation

$$\rho_+(\sigma) = \text{sgn}(\sigma) P_\sigma \in SO(3)$$

has inverse image  $\text{Dic}_3$  of order twelve in  $\text{Spin}(3)$ . Exact quaternion lifts satisfy the local braid and central-sign relations. Strict global Spin closure still requires every branch-complement relator to lift with the correct central sign, equivalently the relevant obstruction class must vanish. A shared-circle SpinC cancellation is a different conclusion.

## 8.2 Which global topology is current?

The q79 Fu–Yau branch is the strongest selected physical compactification candidate in the current corpus. The space  $L(3,1) \times \text{Nil}_3$  remains useful as an auxiliary or effective model. They are not the same manifold; even their first Betti numbers differ. The literal product  $S^1 \times \text{Lens} \times \text{Nil}$  and literal manifold nesting are retired as proof sources.

The shared circle is common  $U(1)$  phase or holonomy data and is counted once. Physical time is noncompact order in the Lorentzian completion. A map  $\mathbb{R} \rightarrow S^1$  can synchronize a phase after its frequency is supplied, but compact phase and time are not identical objects.

## Chapter 9

# The Missing Geometry Theorem

The local real strain carrier and the complex q79 carrier share the rank profile  $1 + 2 + 3$ . Equality of the integer profile does not yet make them objects of the same type. One type-correct candidate is the complex-bundle map

$$\mathfrak{I}_{\text{WW} \rightarrow \text{q79}} : (\mathbb{R}I_3 \oplus \mathcal{D}_0 \oplus \mathcal{S}_{\text{off}}) \otimes_{\mathbb{R}} \mathbb{C} \longrightarrow L_{\text{shared}} \otimes_{\mathbb{C}} (\mathcal{O} \oplus \mathcal{A}_0 \oplus \mathcal{A}).$$

Equivalently, one could first select and prove a rank-six real form of the q79 carrier. Either route is additional source data, not a consequence of the rank count. The resulting map must be a global bundle isomorphism and must preserve:

1. transition functions and the selected flag;
2. the strain and HYM inner products;
3. covariant derivatives;
4. the Hessian and retarded operators; and
5. the normalized overlap kernels used by the numerical packets.

This one theorem would connect the most attractive local interpretation to the most developed global and numerical branch. Repeating the rank count does not advance it.

## Part IV

# From Carrier to Physical Models

## Chapter 10

# Proto-Spinors Without Automatic Particles

If oriented rank-three frame data must retain the nontrivial class of  $\pi_1(SO(3)) \cong \mathbb{Z}_2$ , an ordinary  $SO(3)$  vector description loses that loop memory. The minimal connected cover retaining it is

$$\text{Spin}(3) \cong SU(2).$$

This is a conditional double-cover theorem. It does not prove that nature must begin with rank three, and it does not yet create a Lorentzian fermion.

A physical spacetime spinor requires a Lorentzian spin or SpinC structure, tetrad, compatible connection, hyperbolic Dirac operator, statistics, state space, and observables. Weyl, Dirac, Majorana, and twistor descriptions are conditional chart languages once their chirality, pairing, reality, and conformal assumptions are supplied.

Circle, lens, and nil now name carrier roles:

**Circle.** Shared phase or holonomy.

**Lens.** Finite, projective, or signed-sheet transport.

**Nil.** Triangular transport, anchoring, or termination in a selected operator filtration.

This taxonomy need not be a literal global product topology.

## Chapter 11

# Closure Strain, Scalars, and Mass

The exact local achievement is the strain normal form. A positive closure Hessian proves local restoring cost on its declared slice. It does not prove that one direction is the Higgs or that an eigenvalue is a physical mass.

A pole mass comes from a canonically normalized quadratic action. If

$$S_4^{(2)} = \frac{1}{2} \int \sqrt{|g_4|} \left( Z_{AB} \partial_\mu \varphi^A \partial^\mu \varphi^B - M_{AB}^2 \varphi^A \varphi^B \right),$$

then the tree-level squared masses are eigenvalues of  $Z^{-1/2} M^2 Z^{-1/2}$ . A closure Hessian becomes  $M^2$  only through a normalized source map.

The selected finite-algebra computation begins with three rank-four real scalar-doublet modules. A q79/proto-spinor alignment projector selects one rank-four module and removes eight unwanted real scalar directions at the declared profile tier. This is the relevant one-Higgs selection result. It does not follow from Hessian positivity alone.

## Chapter 12

# A Regime-Local Action, Not a Derived Final Action

The corrected action paper supplies a ten-dimensional EFT ansatz on a declared domain. A representative two-derivative form contains

$$\int_{M_{10}} \sqrt{|g_{10}|} \left[ \frac{R_{10}}{2\kappa_{10}^2} - \frac{1}{4} k_{ab} F_{MN}^a F^{bMN} - \frac{1}{2} G_{AB} D_M \phi^A D^M \phi^B - V(\phi) \right. \\ \left. + i \bar{\Psi} \Gamma^M D_M \Psi - \bar{\Psi} \mathcal{M}(\phi) \Psi \right]$$

plus flux, dilaton, gauge-fixing, ghost, boundary, and higher-order terms.

The metric and Einstein–Hilbert term are inputs. Curvature is defined by a connection;  $\nabla S \neq 0$  alone does not imply curvature or Frobenius failure. Discrete internal spectra require compact resolvent or an equivalent spectral compactness theorem. Consistent four-dimensional truncation requires the discarded equations to vanish on the retained ansatz, or an explicit Schur–Feshbach error bound.

## Part V

# The Standard Model Result at Its Actual Tier

## Chapter 13

# Finite Algebra and Matrix Architecture

The selected finite branch uses

$$\mathcal{A}_F = \mathbb{C} \oplus \mathbb{H} \oplus M_3(\mathbb{C})$$

on an explicit three-family particle–antiparticle carrier. In the declared counting, the current finite construction is a 96-dimensional complex finite Hilbert space with a  $96 \times 96$  profile Dirac operator satisfying self-adjointness, grading, KO-dimension-six reality, order-zero, and structural order-one checks.

The  $27 \times 27$  qutrit–Weyl/minimal matrix is a finite response architecture, not a standard universal size demanded by quantum mechanics. Its dimensions come from the selected tensor and sector organization. Exact algebraic and profile checks establish that the construction can carry the required gauge, family, flavor, phase, Higgs, and threshold data at the adopted standard.

The anomaly analysis selects the usual anomaly-free hypercharge direction on the chosen spectrum and preserves the familiar quotient

$$(SU(3) \times SU(2) \times U(1)_Y)/\mathbb{Z}_6.$$

This is a theorem about the selected finite representation. The local 1+2+3 strain dimensions alone do not derive these representations.

## Chapter 14

# What “True SM Closure” Means Here

The current non-looping ledger reports 12/12 obligations closed for *embedded renormalized-Standard-Model equivalence at the one-shared-physical-primitive/profile standard*. The locked baseline includes:

- the  $27 \times 27$  finite matrix architecture;
- charged Yukawa magnitudes at SM-parity/profile tier;
- selected CKM weight rows and their uncertainty certificate;
- electroweak and direct Higgs/threshold rows;
- the internal threshold ledger;
- the finite Dirac operator at profile tier;
- selected multi-loop mass-scheme and threshold transport;
- a positive-definite precision covariance object; and
- a five-arrow renormalized-SM observable functor.

This is a serious reconstruction result. It shows that the selected MTT branch can encode and reproduce the renormalized SM data consistently through one finite operator architecture and common transport policy.

It is not a strict no-knob theorem. Measured renormalized profile parameters are admitted downstream, and one shared physical primitive fixes the declared normalization. Standard SM BRST/Faddeev–Popov quantization is imported rather than derived from MTT.

### 14.1 Parameters and knobs

Three standards must remain separate:

**Profile parity.** Can the architecture represent the accepted SM data and reproduce them under the frozen conventions?

**One-shared-primitive equivalence.** Can the sectors be tied consistently with one common physical normalization while measured profile values are admitted?

**Strict no-knob selection.** Does one microscopic MTT source select the branch, dimensional normalization, and all values without using those data?

The first two are closed at the stated tier. The third is an upgrade target. Calling all three “closure” without the qualifier caused much of the earlier confusion.

## 14.2 Neutral sector and strong CP

The neutral sector has a selected profile realization with declared holonomy and absolute-scale inputs. Strict selection of the phase, scale, and Dirac/Majorana ontology remains open. The strong-CP program has advanced substantially through the q79 heterotic axion and anomaly analysis, but the strict fully selected numerical nonperturbative source is not yet closed.

## Chapter 15

# Theta Closure and the End of the Few-TeV Chain

The corrected Theta program transports measured source data with a common multi-loop scheme to  $Q = M_t$ . In its stated normalization the gauge-profile targets are

$$\frac{I_2}{I_1} = 0.5110273 \pm 0.0001231, \quad \frac{I_3}{I_1} = 0.158335 \pm 0.001098.$$

They are calibrated profile coordinates, not first-principles predictions. The overlaps are weighted gauge-kinetic coefficients; unit normalization of representatives does not force all of them to one.

An auxiliary round- $S^2$  and Heisenberg-nil ansatz can be retargeted to these ratios and passes its declared dimensionless spectral-floor test. The twistor route reproduces the same quadratic functionals under shared conventions and is therefore a conditional cross-check, not independent normalization.

The old 4.2–5 TeV crossing is withdrawn. It is not a coherence scale, internal gap, quantum-gravity scale, or prediction source. The auxiliary volume coefficient  $20.07064R_1^3$  belongs only to its stated  $S^1 \times S^2 \times \text{Nil}_3$  ansatz and is not the q79 Fu–Yau volume.

The weak-angle relation derived from the same gauge ratio is an exact round-trip identity, not a held-out prediction. A genuine test requires an MTT source for the ratio that does not use electroweak gauge data.

## Part VI

# Quantum, Field-Theoretic, and Gravitational Interpretation

## Chapter 16

# Quantum Mechanics: What Is and Is Not Reconstructed

Projection, coherent basins, and stable sectors can represent alternatives, records, and reduced states. This gives useful language for decoherence and measurement models. It does not produce quantum probability by itself.

A full quantum reconstruction must supply:

1. a complex Hilbert space and observables;
2. unitary dynamics or a specified open-system law;
3. canonical commutation relations where relevant;
4. measurement instruments and update rules; and
5. the Born functional or a theorem equating basin weights with trace probabilities for all relevant states and measurements.

Basin volume depends on a measure. Counting basins is not probability. Noninjective projection is not uncertainty. A deterministic branch narrative must also confront Bell's theorem: MTT cannot be advertised as a Bell-local hidden-variable completion. A consistent global account would need to state its nonfactorization, contextuality, boundary-value, retrocausal, or other mechanism explicitly while preserving no-signaling.

The phrase “no fundamental dice” is therefore a philosophical preference in this book, not a theorem of current MTT.

## Chapter 17

# QFT and the Renormalized SM Functor

The strongest QFT result is a reconstruction on the selected branch. The finite representation and action data feed standard renormalized-SM quantization and multi-loop transport, producing a typed observable functor. This is enough for profile equivalence. It does not derive the path integral, BRST complex, or continuum interacting QFT from projection.

Fiberwise coherent compression can preserve locality of a supplied upper local net when the coherent sector is invariant and the reduction is genuinely fiberwise. This is a meaningful AQFT-style descent theorem. A spatially nonlocal instantaneous kernel is not microcausal merely because it leaves the principal symbol unchanged. A local mediator is the consistent route to causal overlap dynamics.

## Chapter 18

# Worldsheets, Strings, and Fu–Yau Geometry

The proto-spinor/worldsheet paper now defines a  $C^2$  bridge map between declared configuration spaces. If its derivative  $L$  intertwines the two Hessians,

$$L^* H_{\text{ws}} L = H_{\text{ps}},$$

then the quadratic diagnostics agree up to a cubic remainder near alignment. This is a local conditional bridge.

It does not prove that two-dimensionality is inevitable or that a complete string theory follows. Worldsheet Weyl invariance, beta functions, central charge, ghosts, modular invariance, spin-structure sums, anomalies, tadpoles, unitarity, and target-space matching remain independent gates.

Fu–Yau/Hull–Strominger mathematics supplies a legitimate class of heterotic flux backgrounds and is the right established setting for the selected q79 branch. MTT currently reconstructs and selects data inside this setting; it does not contain all of string or M-theory as a proved theorem.

## Chapter 19

# General Relativity and Gravity

The canonical completion supplies a Lorentzian base and a gravitational action or field equation. Coherent reduction may then produce a four-dimensional effective Einstein sector if discarded equations, moduli, warping, gauge vectors, and error terms are controlled.

The internal exact-branch transverse-traceless support packet is a real structural result. Physical Newton/Planck normalization and the complete stress-energy response remain separate. Gravity is therefore a conditional physical completion of the MTT spine, not something projection has already derived.

The image of gravity as “elastic response” can remain an interpretation of the selected action and strain fields. It is not a substitute for deriving the Einstein equations and their observables.

## Chapter 20

# Quantum Gravity: Present Boundary

Filtered propagators and proper-time factors can soften selected Euclidean or TT integrals. This does not prove all-loop finiteness, unitarity, constructive existence, asymptotic completeness, BRST/BV consistency, causal support, or a physical graviton sector.

The old book's “ontic Gaussian damping” and completed UV-finite quantum-gravity language are withdrawn. Current QG papers are conditional models and diagnostic bridges. They may inspire a completion, but they are not evidence that MTT already exceeds established quantum-gravity programs.

## Chapter 21

# Time, Entropy, and Cosmology

Physical time is the noncompact parameter of a selected Lorentzian evolution. Stabilization order  $\tau$ , RG order, chart refinement, and compact circle phase do not become time merely by being ordered.

An arrow of time requires an asymmetric law, state, or boundary condition. A Lyapunov function can orient stabilization on its domain; entropy requires a measure or state and a coarse-graining or thermodynamic theorem. These may be related in a concrete model, but are not currently identical.

The Great Coherence, pre-Big-Bang continuation, inflation as relaxation, dark matter as modal solitons, baryogenesis from geometry, and cosmic renewal are interpretive scenarios. None is a current MTT cosmological prediction. A physical cosmology needs a selected action, solution, initial-state measure, perturbation theory, transport to observables, and comparison with data.

## Part VII

# Relationships to Established Frameworks

## Chapter 22

# A Typed Atlas, Not a Universal Superset

| Framework               | Current relation                                         | Missing promotion                                                                     |
|-------------------------|----------------------------------------------------------|---------------------------------------------------------------------------------------|
| General Relativity      | Conditional physical completion and reduction target     | Selected action, consistent truncation, stress response, and observables.             |
| Quantum Mechanics       | Interpretive reconstruction bridge                       | Born functional, measurement instruments, and full equivalence.                       |
| QFT/AQFT                | Conditional reconstruction with locality descent         | Continuum interacting theory, states, renormalization, scattering, and anomalies.     |
| Standard Model          | Embedded renormalized-SM equivalence at profile standard | Strict no-knob source and unique observed-branch selection.                           |
| Effective Field Theory  | Schur–Feshbach and mode-reduction template               | Nonlinear error, symmetry, power counting, matching, and RG control.                  |
| Kaluza–Klein theory     | Conditional internal-mode reconstruction                 | Selected compactification, warping, mixing, and stabilized moduli.                    |
| Noncommutative geometry | Selected finite-algebra realization                      | Full spectral- triple axioms and unique geometric source where not already certified. |
| String/Calabi–Yau       | Encoding and reconstruction program                      | Complete worldsheet consistency, dualities, branes/fluxes, and low-energy matching.   |

| Framework                              | Current relation                                | Missing promotion                                                                   |
|----------------------------------------|-------------------------------------------------|-------------------------------------------------------------------------------------|
| Hull–Strominger<br>and Fu–Yau          | Conditional<br>solution-slice<br>reconstruction | Unique selected background and<br>same-source operator/value emission.              |
| Quantum gravity                        | Open conditional<br>program                     | Unitarity, causality, continuum<br>existence, BRST/BV, and physical<br>observables. |
| LQG, causal sets,<br>asymptotic safety | Interpretive or<br>conditional bridges          | Explicit controlled maps and<br>bidirectional observable equivalence.               |

Multiple MTT realizations reproducing the same target data create underdetermination. Agreement proves realizability, not inevitability. A canonical source theorem must select among inequivalent realizations.

## Part VIII

# What Has Been Achieved and What Comes Next

## Chapter 23

# The Genuine Achievements

### 23.1 Methodological

MTT now has a disciplined separation of assumptions, conditional theorems, reconstructions, profiles, and predictions. This sounds modest, but it prevents dozens of attractive category errors and makes the program auditable.

### 23.2 Functional analytic

The corrected Foundation and Fixed Points series provide a reusable framework for vertical spectral reduction, nonnormal-safe damping, multiple existence and uniqueness routes, curved-projector persistence, leakage, covariance, and admissibility exits.

### 23.3 Geometric

The local  $3 \times 3$  comparison carrier and its exact  $3 + 6$  and  $1 + 2 + 3$  decompositions are verified. The q79 degree-three cover supplies a separate global  $1 + 2 + 3$  trace carrier, exact local signed-sheet spin data, and a developed Fu–Yau/HYM branch. The missing bridge is now finite and explicit.

### 23.4 Computational

The selected finite-algebra and  $27 \times 27$  architecture pass extensive exact and numerical checks. Embedded renormalized-SM equivalence closes at the declared one-shared-primitive/profile standard, including multi-loop precision transport and covariance. This is unusual in breadth for a speculative framework, even though it is not yet no-knob prediction.

### 23.5 Epistemic

Several former claims were disproved or retired: the right-inverse argument, Lorentzian Gram metric, extra-circle dimension count, few-TeV chain, independent weak-angle test, literal

Lens–Nil/q79 identity, and automatic Born/QG claims. Finding and correcting these failures is part of the achievement.

## Chapter 24

# The Decisive Frontier

The next work should follow this order.

### 24.1 Global q79 spin decision

Lift a presentation of the q79 branch complement and compute the central sign of every relator. This decides strict Spin closure. If the obstruction does not vanish, state the SpinC alternative without relabeling it as Spin.

### 24.2 World-in-world to q79 intertwiner

Construct  $\mathfrak{I}_{\text{WW} \rightarrow \text{q79}}$  with transition, metric, connection, Hessian, retarded, and overlap compatibility. This is the most important foundational bridge for the numerical program.

### 24.3 One selected action source

Use one q79/Fu–Yau action, gauge fixing, regulator, scale, and scheme to emit normalized zero modes and all required finite response rows. Verify the discarded equations or a reduction error bound.

### 24.4 Strict value and branch selection

Freeze the source before looking at the held-out data. Count continuous and discrete inputs honestly. Then test future or unused common-scheme observables with uncertainty and a preregistered acceptance rule.

### 24.5 Probability and quantization

Construct the measure/state and prove, or falsify, a basin–trace equality. Derive the BRST and path-integral structure from MTT only if one common source map actually produces it.

## 24.6 Gravity and cosmology

After the action is selected, derive the Lorentzian equations, constraints, stress response, Planck normalization, cosmological backgrounds, perturbations, and observables. Interpretive cosmology comes last, not first.

## Chapter 25

# Falsifiability Without Promotional Predictions

MTT can fail before new experimental data arrive:

- the q79 relator signs may obstruct the desired strict Spin lift;
- no global connection-preserving  $1 + 2 + 3$  intertwiner may exist;
- the selected action may fail consistent truncation or anomaly control;
- normalized source rows may miss the locked profile;
- parameter counting may collapse back to the SM input count;
- the basin–trace or locality bridge may be impossible; or
- independent held-out observables may disagree with the frozen branch.

A genuine empirical prediction must list every datum used in geometry, normalization, scheme choice, branch choice, and parameter fitting. Agreement with a value used anywhere in that chain is a profile check or postdiction.

## Chapter 26

# An Interpretive Picture That Survives

The surviving picture is quieter than version 9 and more interesting. Physical descriptions may be stable projections of richer geometric and operator data. Different theories may be charts or reconstructions on overlapping domains without one being literally contained in another. Triplet structure can be a selected rank filtration rather than a mystical universal number. The shared circle can coordinate phase without becoming time. Particles can be studied as stable excitations without pretending that stability alone determines their quantum numbers. Gravity can be compared to strain response after an action is supplied. Measurement can be modeled as a physical interaction without announcing a derivation of probability.

This interpretation is useful because each sentence points to a theorem that could be built. It asks for maps instead of resemblance, sources instead of fitted rows, and observables instead of slogans.

## Chapter 27

# Afterword

MTT is no longer best described as “one field, three filters, one inequality, one inevitable world.” The current program is richer:

One typed spectral architecture; several exact internal modules; one selected q79/Fu–Yau and finite-algebra branch; a closed renormalized-SM profile equivalence; and a short list of decisive bridges still to prove.

That statement gives up inevitability but gains contact with mathematics. It also preserves the original impulse of the book: physics should become more coherent as its hidden assumptions are exposed. The next success will not be another sweeping interpretation. It will be one of the missing maps actually constructed and independently checked.

# Appendix A

## Current Claim Ledger

| Claim                                 | Current status         | Controlling boundary                                                      |
|---------------------------------------|------------------------|---------------------------------------------------------------------------|
| Hilbert-bundle spectral/control spine | Conditional theorem    | Operator domains, commutation, gaps, invariant domains.                   |
| Projected fixed points                | Conditional theorem    | Existence route and fixed-point type must be stated.                      |
| Equilibrium promotion                 | Conditional theorem    | Strict Lyapunov or equivalent stationarity gate.                          |
| 4 + 6 physical realization            | Canonical input        | Not derived from abstract MTT.                                            |
| $3 \times 3 = 3 + 6$ local split      | Proved                 | Nonsingular comparison background.                                        |
| 1 + 2 + 3 strain split                | Proved with flag       | Flag is selected input/source.                                            |
| q79 1 + 2 + 3 trace carrier           | Selected theorem       | Degree-three cover.                                                       |
| Local Dic <sub>3</sub> lift           | Proved                 | Strict global relators still open.                                        |
| Fu–Yau/HYM branch                     | Selected realization   | Rank-two literal continuum witness; stronger transfer/uniqueness guarded. |
| Finite matrix and Dirac operator      | Closed at profile tier | Selected finite representation and profile rows.                          |

| Claim                        | Current status                                            | Controlling boundary                                               |
|------------------------------|-----------------------------------------------------------|--------------------------------------------------------------------|
| Embedded renormalized SM     | Closed at<br>one-shared-<br>primitive/profile<br>standard | Measured profile and<br>imported SM<br>quantization admitted.      |
| Strict no-knob SM            | Open upgrade                                              | Unique source and<br>branch selection.                             |
| Born rule                    | Open                                                      | Measure/state and<br>basin–trace theorem.                          |
| Einstein gravity             | Conditional<br>completion                                 | Selected Lorentzian<br>action and controlled<br>reduction.         |
| UV-complete quantum gravity  | Open                                                      | Full constructive, causal,<br>gauge, and unitarity<br>obligations. |
| Cosmological origin scenario | Interpretation                                            | Selected solution and<br>observable pipeline<br>absent.            |

## Appendix B

# Notation and Guardrails

$M_{10} \rightarrow Y_4$ . Canonical supplied physical bundle with compact six-dimensional fiber  $X_6$ .

$Q_{\text{WW}}$ . Rank-three world-in-world comparison field; never the noncoherent projector.

$P_{\text{coh}}$ . Joint coherent spectral projector.

$Q_{\text{inc}}$ . Complementary noncoherent projector  $I - P_{\text{coh}}$ .

$\tau, t, \mu$ . Stabilization, physical time, and RG scale; distinct by default.

$L_{\text{shared}}$ . Common phase/holonomy line; not an extra dimension or time.

$\mathcal{S}_{\text{off}}$ . Real symmetric off-diagonal strain subspace; distinct from the structure sheaf  $\mathcal{O}$  used in the q79 carrier.

$\mathcal{A}, \mathcal{A}_0$ . Rank-three q79 cover algebra and its rank-two trace-zero summand.

$\mathcal{H}_{\text{q79}}$ . Shared q79 rank-six  $1 + 2 + 3$  carrier.

**Profile standard.** A reconstruction tier admitting measured renormalized coordinates and one shared physical primitive.

**Strict standard.** A source-selected, no-target-input theorem with a frozen branch and held-out validation.

# References

# Bibliography

- [1] P. Nero, *Modal Triplet Theory: Foundations*, version 7, 2026.
- [2] P. Nero, *Fixed Points I–VI*, corrected editions, 2026.
- [3] P. Nero, *The Projection–Admissibility Principle: Descent, Recovery, and Structural Constraints*, version 2, 2026.
- [4] P. Nero, *Lorentzian Base Compatibility and Signature Stability in the MTT Fixed-Point Realization*, version 2, 2026.
- [5] P. Nero, *Modal Triplet Theory: A Typed Relationship Atlas*, version 3, 2026.
- [6] P. Nero, *MTT Foundational Geometry Reconciliation*, theorem and verification packet, 2026.
- [7] P. Nero, *The Proto-Spinor: Conditional Spinorial Closure and the q79 Interface*, version 5, 2026.
- [8] P. Nero, *World-in-World Genesis: Local Comparison Geometry and a Globalization Program*, version 5, 2026.
- [9] P. Nero, *Closure-Strain Geometry: Local Normal Forms and Conditional Matter Encodings*, version 6, 2026.
- [10] P. Nero, *Closure Geometry and a Regime-Local Ten-Dimensional Action Ansatz*, version 4, 2026.
- [11] P. Nero, *MTT Current True SM Closure Consolidated Ledger*, theorem and verification packet, 2026.
- [12] P. Nero, *Theta Closure and Execution I–V*, corrected editions, 2026.
- [13] J.-X. Fu and S.-T. Yau, “The theory of superstring with flux on non-Kähler manifolds and the complex Monge–Ampère equation,” *Journal of Differential Geometry* **78** (2008), 369–428.
- [14] H. B. Lawson and M.-L. Michelsohn, *Spin Geometry*, Princeton University Press, 1989.
- [15] A. Connes, *Noncommutative Geometry*, Academic Press, 1994.

- [16] J. Polchinski, *String Theory*, Volumes 1–2, Cambridge University Press, 1998.
- [17] S. Weinberg, *The Quantum Theory of Fields*, Cambridge University Press, 1995–2000.