

From Modal Triplet Theory to Noncommutative Geometry: A Typed Almost-Commutative Encoding and Its Current Finite-Triple Frontier

Peter Nero

July 2026, Version 5

Abstract

We analyze the relation between Modal Triplet Theory (MTT) and noncommutative geometry (NCG). A coherent fixed-point projector does not by itself derive a spectral triple, the Standard Model finite algebra, a Wick rotation, or the spectral action. We prove the precise compression theorem that is available when a projector reduces an already specified real even spectral triple. We then record the stronger finite results now available in the selected MTT branch. The family factor is separated from the rank $1 < 2 < 3$ gauge carrier; an explicit three-family chiral representation and the faithful group

$$(SU(3) \times SU(2) \times U(1)_Y)/\mathbb{Z}_6$$

are obtained; and the finite real-even triple is completed at profile tier by the additional neutral complex summand $\mathbb{C}_N$. Direct enumeration of all 676 finite one-form basis pairs gives a raw rank-twelve scalar space, consisting of three Higgs-doublet modules. The selected q79/proto-spinor alignment projector has rank four and leaves one complex Higgs doublet. These are finite, reproducible results, but the entries of D_F remain profile data. The selected representation also derives the one-loop coefficient vector $(41/10, -19/6, -7)$, and all ten matter, scalar, and gauge/ghost internal spectrum rows are now exact finite calculations with no added continuous parameter. Their normalized nonzero spectral shape is universal, however, so they do not generate the missing nonuniversal gauge thresholds. The four-dimensional compact Riemannian spin triple, Wick dictionary, cutoff function, and spectral moments are not derived by the fixed-point construction. Moreover, the selected pure-Standard-Model running branch rules out one universal gauge normalization: the best common-scale coupling ratio remains 1.046656046. A positive overlap kinetic metric reconstructs the profile couplings exactly but uses two measured relative coordinates. Thus MTT currently supplies a typed almost-commutative profile encoding with an executed finite factor, not a first-principles derivation of NCG or a no-knob derivation of the Standard Model.

Version 5 Revision Note

Supersedes *From Modal Triplet Theory to Noncommutative Geometry: Spectral Triples, Spectral Action, and the Standard Model Embedding*, corrected version 4.

Reason Version 4 correctly replaced the original first-principles claims by a typed finite profile encoding, but it predated the completed gauge-fixed fluctuation complex and the exact ten-row internal spectrum calculation.

Resolution Version 5 retains the A44–A52 correction and adds the verified A57–A62 finite gauge/ghost chain. It records the exact representation-theoretic one-loop coefficients and all ten exact internal spectrum rows, while also recording the resulting no-go for obtaining nonuniversal gauge thresholds from their common normalized shape.

Retained result Given the declared Euclidean spin triple and the selected finite profile triple, standard almost-commutative NCG reconstructs the corresponding bosonic operator content.

Remaining boundary A physical Lorentzian-to-Euclidean dictionary, source-derived D_F values, a selected proper-time measure and overlap metric, a physical continuum HYM threshold source, nonperturbative quantum transport, and a same-source physical q79 continuum realization remain open.

Contents

Version 5 Revision Note	1
1 Status and logical scope	3
2 The spectral-triple contract	4
2.1 What a fixed-point projection can preserve	4
3 The finite MTT carrier and the algebra no-go	5
3.1 The qutrit algebra is not the Standard Model algebra	5
3.2 Native automorphisms and the global gauge group	5
4 The completed finite real-even triple	6
4.1 Why the neutral summand is required	6
4.2 One physical abelian circle	6
5 Finite one-forms and the Higgs sector	7
6 Euclidean product triple and the Wick contract	7
7 Spectral action: conditional output and no-go results	8
7.1 What the finite execution closes	8
7.2 Gauge-fixed heat data and the ten finite spectra	8
7.3 Universal gauge normalization is excluded	9
7.4 Moment identifiability	9
8 Relation to q79, fixed points, and quantum field theory	10
8.1 The shared circle	10
8.2 Renormalization	10
9 Tiered synthesis theorem	10
10 Reproducibility and remaining frontier	11
11 Conclusion	11

1 Status and logical scope

This paper concerns a relation among three different constructions:

1. a coherent or fixed-point projection in MTT;
2. a finite real-even spectral triple encoding the selected particle representation; and
3. a four-dimensional Euclidean almost-commutative product triple and its spectral action.

They are not interchangeable. In particular, a projector is not an algebra, a spectral gap is not a Dirac operator, and a finite representation is not a Wick rotation.

Definition 1.1 (Status vocabulary). In this paper:

- *exact* means an algebraic identity or finite calculation verified from the cited packet without measured continuous values;
- *profile* means a reconstruction using an accepted Standard Model parameter profile;
- *imported* means a standard NCG or QFT theorem used after its hypotheses are supplied independently; and
- *open* means that the current MTT source does not emit the required object.

Exact carrier. The rank- $1 < 2 < 3$ carrier and weak real structure are selected up to unitary or phase equivalence; the family factor remains separate.

Exact representation. The 48-state family-diagonal representation, anomaly table, gauge group, and global $\mathbb{Z}_6$ quotient are executed.

Mixed finite triple. The representation and finite axioms are exact finite calculations; physical entries of D_F use the accepted profile.

Mixed scalar sector. The one-form enumeration and rank statements are finite calculations; the Yukawa traces use profile D_F .

Exact finite fluctuation spectra. The representation-theoretic one-loop coefficient vector and all ten matter, scalar, and gauge/ghost internal spectrum rows are exact finite calculations. Their universal normalized shape is also an exact obstruction to using these rows alone as nonuniversal gauge thresholds.

Imported/open spacetime. The Euclidean product triple is standard NCG input; MTT has not derived its physical Wick dictionary.

Open/profile normalization. Only certain spectral products are identifiable; the current overlap metric uses measured relative gauge coordinates.

A concrete picture of the finite construction

A finite spectral triple may be read as a small, structured quantum-mechanical system. The algebra $\mathcal{A}_F$ lists the internal observables, the Hilbert space $\mathcal{H}_F$ lists the fermionic states on which they act, and the matrix D_F connects states of opposite chirality. The commutator $[D_F, a]$ measures how an internal observable a changes across those connections. Linear combinations of terms $a[D_F, b]$ are the finite one-forms; after the standard reality completion they are the possible scalar and gauge fluctuations of the supplied finite geometry.

In the present construction one family has sixteen chiral particle states. Three families give forty-eight, and adjoining the charge-conjugate states gives the 96-dimensional space on which the completed real-even triple acts. For example, the weak doublet $Q = (u_L, d_L)$ carries the quaternionic $SU(2)$ action, its three color copies carry the $M_3(\mathbb{C})$ action, and its abelian charge comes from the selected anomaly-free complex phase. Entries of D_F connect these left-handed states to the corresponding right-handed singlets. Algebraically those entries have the role of Yukawa and neutral mass matrices; numerically they remain profile data in this paper.

This picture also explains the Higgs calculation. Enumerating all $a[D_F, b]$ asks which scalar bridges the finite algebra permits. The unrestricted completed triple permits three real rank-four modules, or three complex weak doublets. The selected q79/proto-spinor alignment projector retains one rank-four module and removes the other eight real scalar directions. The later ten spectrum rows ask a different question: what are the eigenvalues of the selected finite heat operators in the seven matter/scalar and three gauge/ghost sectors? They certify the internal finite shape; they are not yet physical spacetime masses or continuum thresholds.

2 The spectral-triple contract

Definition 2.1 (Real even spectral triple). A real even spectral triple is data

$$(\mathcal{A}, \mathcal{H}, D, J, \Gamma, \pi)$$

where $\mathcal{A}$ is a unital involutive algebra represented faithfully by bounded operators $\pi(a)$ on the complex Hilbert space $\mathcal{H}$, D is self-adjoint with compact resolvent, and

$$[D, \pi(a)] \in \mathcal{B}(\mathcal{H}) \quad (a \in \mathcal{A}).$$

The grading and real structure obey the declared KO-sign table, and the order-zero and order-one conditions are

$$[\pi(a), J\pi(b)J^{-1}] = 0, \quad [[D, \pi(a)], J\pi(b)J^{-1}] = 0.$$

Orientability and Poincare duality are additional requirements; they do not follow merely from compact resolvent and bounded commutators.

This definition makes the missing data in a projection-only argument visible. A coherent subspace can be useful without carrying a faithful algebra action, a self-adjoint Dirac operator, a real structure, or a nondegenerate intersection form.

2.1 What a fixed-point projection can preserve

Theorem 2.2 (Reducing-projector spectral-triple theorem). *Let $(\mathcal{A}, \mathcal{H}, D, J, \Gamma, \pi)$ be a real even spectral triple and let P be an orthogonal projection such that:*

1. *P commutes with $\pi(a)$ for every $a \in \mathcal{A}$;*
2. *P reduces D , equivalently it commutes with the spectral projections of D ;*
3. *P commutes with J and Γ ; and*
4. *the restricted representation $\pi_P(a) = P\pi(a)|_{P\mathcal{H}}$ is faithful.*

Then

$$(\mathcal{A}, P\mathcal{H}, D_P, J_P, \Gamma_P, \pi_P), \quad D_P = D|_{P\mathcal{H}},$$

is a real even spectral triple with the inherited KO signs and order-zero and order-one relations. Orientability and Poincare duality are inherited only when the compressed Hochschild cycle and intersection pairing satisfy the corresponding axioms.

Proof. Because P reduces D , the restriction D_P is self-adjoint. On $P\mathcal{H}$,

$$(D_P - i)^{-1} = P(D - i)^{-1}P,$$

which is compact. For $a \in \mathcal{A}$,

$$[D_P, \pi_P(a)] = P[D, \pi(a)]P,$$

so the commutator is bounded. Commutation of P with J and Γ restricts the real and graded relations, while the order-zero and order-one identities follow by compression. Faithfulness is assumed explicitly. Neither a Hochschild orientation cycle nor nondegeneracy of the finite intersection form is implied by these operator restrictions. $\square$

Corollary 2.3 (Limit of fixed-point reasoning). *A fixed-point or harmonic projector alone does not construct a spectral triple. It must either act on prior spectral-triple data as in Theorem 2.2, or an independent theorem must construct the algebra, representation, Dirac domain, real structure, grading, and finite axioms on its image.*

This is the precise separation between the MTT fixed-point architecture and the NCG encoding. The corrected MTT Foundation supplies typed Hilbert-bundle, projector, gap, and stability hypotheses; it does not automatically supply the inputs of Theorem 2.2.

3 The finite MTT carrier and the algebra no-go

3.1 The qutrit algebra is not the Standard Model algebra

The selected finite qutrit package is

$$\mathcal{A}_Q = \mathbb{C}_{\text{family}}^3 \otimes M_3(\mathbb{C}) \cong M_3(\mathbb{C}) \oplus M_3(\mathbb{C}) \oplus M_3(\mathbb{C}).$$

The original three-summand Standard Model candidate is

$$\mathcal{A}_F^{(0)} = \mathbb{C} \oplus \mathbb{H} \oplus M_3(\mathbb{C}).$$

Proposition 3.1 (No direct algebra identification). *There is no real involutive-algebra isomorphism $\mathcal{A}_Q \cong \mathcal{A}_F^{(0)}$.*

Proof. The real dimensions are

$$\dim_{\mathbb{R}} \mathcal{A}_Q = 54, \quad \dim_{\mathbb{R}} \mathcal{A}_F^{(0)} = 24.$$

Their centers also have different real dimensions:

$$\dim_{\mathbb{R}} Z(\mathcal{A}_Q) = 6, \quad \dim_{\mathbb{R}} Z(\mathcal{A}_F^{(0)}) = 5.$$

Either invariant rules out an isomorphism. $\square$

The lane-wise corner construction of A44 is an exact abstract reduction, but A45 corrects its physical interpretation: the $\mathbb{C}^3$ label is the family/character factor and cannot simultaneously be used as three family-dependent gauge-rank lanes. The physical construction therefore keeps

$$\mathcal{H}_{\text{chiral}} = \mathbb{C}_{\text{family}}^3 \otimes \mathcal{H}_{16}, \quad \rho_{\text{gauge}} = \mathbf{1}_3 \otimes \rho_{16},$$

and builds the gauge factor separately from the selected rank flag.

3.2 Native automorphisms and the global gauge group

On the selected rank-one, rank-two, and determinant-trivial rank-three carriers, the automorphism groups are

$$U(1), \quad USp(2) \cong SU(2), \quad SU(3).$$

The rank-two result uses the selected symplectic antiunitary $J = \epsilon K$, and the rank-three result uses the determinant-trivial carrier. On the explicit chiral representation the diagonal central kernel is $\mathbb{Z}_6$. Thus the faithful low-energy group is

$$G_{\text{SM}} = \frac{SU(3) \times SU(2) \times U(1)_Y}{\mathbb{Z}_6}.$$

The associated 48-state three-family chiral representation contains the rows Q, u^c, d^c, L, e^c, N^c and passes the local and global anomaly tests in the A46–A47 packets [5].

4 The completed finite real-even triple

4.1 Why the neutral summand is required

The explicit particle-antiparticle construction has complex dimension 96. For the accepted charged and neutral profile matrices, the finite Dirac operator satisfies self-adjointness, oddness, KO-dimension-six reality, order zero, and order one. The three-summand candidate $\mathcal{A}_F^{(0)}$, however, fails two remaining finite axioms:

- the $N_R : \mathbb{C} - \mathbb{C}$ self-edge obstructs the required orientation; and
- the antisymmetric 3×3 intersection form has determinant zero.

The minimal selected completion is

$$\mathcal{A}_F = \mathbb{C} \oplus \mathbb{H} \oplus M_3(\mathbb{C}) \oplus \mathbb{C}_N,$$

where

$$\mathbb{C}_N = \text{End}_{\mathbb{C}}(1_M), \quad 1_M = N^c.$$

Only the neutral edge is changed, from $\mathbb{C} - \mathbb{C}$ to $\mathbb{C}_N - \mathbb{C}$. The completion has an explicit 17-term Hochschild zero-cycle and intersection determinant 4 per family, hence 324 for the three-family product. No continuous parameter or new state is added.

4.2 One physical abelian circle

Let (α, μ, ν) denote the phases of $\mathbb{C}, M_3(\mathbb{C}), \mathbb{C}_N$. The independent anomaly equations are

$$\alpha + 3\mu = 0, \quad \alpha - \nu = 0.$$

Their primitive integer null vector is

$$(\alpha, \mu, \nu) = (3, -1, 3),$$

which gives

$$6Y(Q, u^c, d^c, L, e^c, N^c) = (1, -4, 2, -3, 6, 0).$$

The cubic anomaly vanishes on the same line, whereas an independent $\mathbb{C}_N$ phase is anomalous. Distinct algebra sheets therefore share one anomaly-free physical hypercharge circle.

Theorem 4.1 (Current finite-triple result). *The A45–A50 packet chain defines an explicit three-family real-even finite triple on the completed algebra $\mathcal{A}_F$ at the declared profile tier. The representation, KO signs, order-zero and order-one identities, orientability, Poincare pairing, anomaly-free hypercharge line, and global $\mathbb{Z}_6$ quotient are finite executable statements. The numerical entries of D_F , including Yukawa and Majorana data, remain accepted profile coordinates rather than source-derived MTT predictions.*

Proof. The representation and anomaly table are A46; the native automorphism group and central quotient are A47; the explicit bimodule is A48; the finite Dirac and three-summand obstruction are A49; and the neutral completion and unique abelian anomaly-free line are A50. Their hash-addressed artifacts and independent verifiers are recorded in the curated result repository [5]. The profile status of the entries of D_F is part of the A49 contract and is not changed by the algebraic checks. $\square$

5 Finite one-forms and the Higgs sector

For the completed finite triple, define

$$\Omega_{D_F}^1(\mathcal{A}_F) = \text{span}\{\rho(a)[D_F, \rho(b)] : a, b \in \mathcal{A}_F\}.$$

The A51 calculation evaluates all $26 \times 26 = 676$ real-algebra basis pairs and then imposes self-adjoint real fluctuations

$$A_{\text{real}} = A + J_F A J_F^{-1}.$$

Proposition 5.1 (Raw scalar multiplicity and selected projection). *The unrestricted real fluctuation space has real rank 12 and decomposes into three rank-four scalar-doublet modules,*

$$H_{\text{up}}, \quad H_{\text{down/charged lepton}}, \quad H_{\text{neutrino}}.$$

Consequently the raw completed finite triple is not automatically the one-Higgs Standard Model. The selected q79/proto-spinor alignment rule

$$H_{\text{up}} = H_{\text{neutrino}} = H, \quad H_{\text{down}} = H_{\text{charged lepton}} = -\epsilon \bar{H}$$

defines a self-adjoint idempotent rank-four submodule and removes the eight unwanted real scalar directions.

The finite execution reports inclusion residual 6.152×10^{-15} in its floating-point realization. The surviving module is one complex $SU(2)$ doublet of hypercharge $+1/2$, with pseudoreality supplying its conjugate channels. This is an executed finite selection. It is not yet a derivation of the same projector from the physical nonzero-Chern q79 HYM connection.

For three families, the finite gauge traces are

$$k_Y : k_2 : k_3 = 10 : 6 : 6.$$

After the conventional $5/3$ hypercharge normalization they become $6 : 6 : 6$. The accepted profile D_F gives

$$a = \text{Tr}(Y^\dagger Y) = 3.15667873398489, \quad b = \text{Tr}((Y^\dagger Y)^2) = 3.31696406124945,$$

$$\frac{b}{a^2} = 0.332874093710992.$$

These are profile traces at M_Z , not high-scale predictions.

6 Euclidean product triple and the Wick contract

Let (Y_E, g_E) be a compact oriented four-dimensional Riemannian spin manifold with spin triple

$$(C^\infty(Y_E), L^2(Y_E, S_E), D_E^{(4)}, J_E, \gamma_5).$$

Given the finite factor of Theorem 4.1, the standard product data are

$$\begin{aligned} \mathcal{A} &= C^\infty(Y_E) \otimes \mathcal{A}_F, \\ \mathcal{H} &= L^2(Y_E, S_E) \otimes \mathcal{H}_F, \\ D_{\text{prod}} &= D_E^{(4)} \otimes \mathbf{1} + \gamma_5 \otimes D_F. \end{aligned}$$

Theorem 6.1 (Conditional product-triple theorem). *If the Euclidean spin triple and the finite real-even triple satisfy the standard sign, order, orientation, and nondegeneracy hypotheses, then their graded product is an almost-commutative real even spectral triple. Inner fluctuations generate the associated gauge fields and finite scalar modules.*

This is a standard NCG theorem [2, 3, 6]; it is not a theorem that MTT has selected (Y_E, g_E) or its Dirac operator.

Definition 6.2 (Physical Wick contract). A physical use of Theorem 6.1 requires, at minimum:

1. a Lorentzian spin background and causal field domain;
2. a declared analytic continuation to the Euclidean background;
3. maps for spinors, gauge fields, scalar fields, actions, and boundary conditions;
4. a contour and inverse domain controlling singularities and zero modes; and
5. a theorem relating Euclidean correlation functions or determinants to Lorentzian observables.

The spectral-action formulation is intrinsically Euclidean at this stage. Wick rotation and the removal of unphysical fermion doubling require a separate prescription [4]. A compact phase circle or the q79 shared differential line is not physical time and does not supply this contract automatically.

7 Spectral action: conditional output and no-go results

For a positive cutoff profile f and scale Λ , write the bosonic spectral action in the convention

$$S_B(f, \Lambda) = \text{Tr } f(D_{\text{prod}}^2/\Lambda^2).$$

Under the usual heat-kernel hypotheses,

$$S_B(f, \Lambda) \sim \Lambda^4 f_4 a_0(D_{\text{prod}}^2) + \Lambda^2 f_2 a_2(D_{\text{prod}}^2) + f_0 a_4(D_{\text{prod}}^2) + \cdots,$$

where, in this convention,

$$f_4 = \int_0^\infty f(u) u \, du, \quad f_2 = \int_0^\infty f(u) \, du, \quad f_0 = f(0).$$

The heat-kernel theorem produces cosmological, Einstein–Hilbert, Yang–Mills, scalar kinetic and potential, and nonminimal curvature terms from the supplied product triple [1, 7].

7.1 What the finite execution closes

Combining Theorem 6.1 with Theorem 5.1 closes the bosonic *operator content* at profile tier: three gauge terms, one Higgs covariant kinetic term and potential, and the standard gravitational terms. It does not fix absolute coefficients until the base geometry, cutoff data, field normalization, and RG transport are supplied.

7.2 Gauge-fixed heat data and the ten finite spectra

The later finite execution sharpens this statement without changing its tier. From the selected Q, u, d, L, e, N chiral representations, the one-Higgs scalar module, the adjoint gauge blocks, and Faddeev–Popov ghosts, A57 derives the standard one-loop coefficient vector

$$b = \left(\frac{41}{10}, -\frac{19}{6}, -7 \right)$$

by representation indices. The accepted QFT vector is not used to select this result. Its interpretation as a physical running law still imports the usual gauge-fixed one-loop QFT theorem.

A58–A62 then execute the seven Q, u, d, L, e, N, H rows and the three $U(1)$, $SU(2)$, and $SU(3)$ gauge/ghost rows. The final two nonabelian rows are

	0	$4\pi^2/9$	$8\pi^2/9$
$SU(2)$	3	12	12
$SU(3)$	8	32	32

with the table entries giving multiplicities. The $U(1)$ adjoint self-interaction row vanishes, as required by $C_2(U(1)) = 0$. These ten internal spectra are exact and add no continuous scale.

This completion has a useful negative consequence. After normalization per representation or adjoint rank, every nontrivially weighted row has the same finite spectral shape. It can supply a common finite heat factor, but not the different threshold corrections needed to repair the universal-boundary failure below. A nonuniversal physical threshold still requires a same-source continuum HYM or mass-dependent operator, not a relabeling of the finite spectra.

7.3 Universal gauge normalization is excluded

Equal normalized finite traces with one universal f_0 and identity kinetic metric would require

$$g_1^{\text{GUT}}(Q) = g_2(Q) = g_3(Q)$$

at one scale. On the selected pure-Standard-Model running branch, A52 finds the best one-loop point

$$Q_{\text{best}} = 1.71709699661 \times 10^{14} \text{ GeV},$$

$$(g_1, g_2, g_3) = (0.556290097, 0.535101507, 0.560067228), \quad \frac{\max g_i}{\min g_i} = 1.046656046.$$

The archived multi-loop trajectory has the same failure.

Proposition 7.1 (Universal-normalization no-go). *On the selected pure-Standard-Model profile branch, no single universal spectral moment with identity gauge kinetic metric reproduces all three running gauge couplings.*

The exact profile reconstruction instead uses

$$K_{\text{gauge}} = \text{diag}(1.9568425763574, 1, 0.309837025907476), \quad f_0 = 0.397408225924864,$$

in $g_i^{-2} = 6f_0K_i$, with residual 4.578×10^{-16} . Its two relative entries are the two measured relative coupling coordinates. They add no parameters beyond the Standard Model profile but are not MTT predictions.

7.4 Moment identifiability

The displayed action determines only

$$f_0K_i, \quad f_2\Lambda^2, \quad f_4\Lambda^4.$$

Proposition 7.2 (Spectral-moment identifiability). *The spectral action above cannot separately determine Λ, f_2, f_4 .*

Proof. For every $c > 0$, the transformation

$$\Lambda \mapsto c\Lambda, \quad f_2 \mapsto f_2/c^2, \quad f_4 \mapsto f_4/c^4$$

leaves $f_2\Lambda^2$ and $f_4\Lambda^4$ unchanged. □

The former claim that a fixed-point gap alone fixes the cutoff function, spectral moments, Newton normalization, and vacuum term is therefore withdrawn. A selected proper-time measure and source-derived overlap metric would be needed for that stronger result.

8 Relation to q79, fixed points, and quantum field theory

8.1 The shared circle

The finite triple has distinct complex algebra sheets but one anomaly-free hypercharge circle. The corrected Foundation and Proto-Spinor papers also identify a universal flat q79 differential line and finite carrier actions. The equality of these roles is not automatic. A physical identification must preserve the selected connections, covariant derivatives, Hessians, and holonomies. The nonzero-Chern HYM realization and its local strain-to-q79 continuum intertwiner remain open.

8.2 Renormalization

The A03 observable functor recovers perturbative Standard Model observables after importing the standard gauge-fixed renormalized action and BRST/Faddeev–Popov quantization. That result is compatible with the finite NCG encoding but does not derive quantization from the spectral triple. Conversely, the Euclidean spectral action does not establish a nonperturbative Lorentzian QFT, a quantum BV pushforward, or ultraviolet completeness. The v3 claim that FRG, pAQFT, and the MTT quantum-gravity program “match” from the common vector Θ is therefore replaced by this typed compatibility statement.

9 Tiered synthesis theorem

Theorem 9.1 (Selected MTT–NCG encoding theorem). *Assume:*

1. *the exact selected finite carrier, chiral representation, native gauge automorphisms, neutral completion, and anomaly-free circle of A45–A50;*
2. *the accepted profile D_F and the selected rank-four scalar projector executed in A51;*
3. *an independently supplied compact four-dimensional Euclidean spin triple; and*
4. *a cutoff profile, scale, field normalization, and RG convention.*

Then the standard almost-commutative product construction gives a real even spectral triple and reproduces the bosonic Standard Model operator content with one Higgs doublet at the declared profile tier.

The conclusion does not derive:

1. *the physical Lorentzian spacetime or its Wick rotation;*
2. *the accepted entries of D_F from source geometry;*
3. *a universal spectral normalization on the pure-SM branch;*
4. *a unique cutoff function, scale, or spectral moments;*
5. *a no-knob Standard Model parameter point; or*
6. *the renormalized quantum theory or ultraviolet completion.*

Proof. The finite premises give Theorem 4.1 and Theorem 5.1. The imported Euclidean premise gives Theorem 6.1; the standard heat-kernel expansion then gives the bosonic operator content. The six exclusions follow respectively from the missing Wick contract, the profile status of D_F , the no-go in Theorem 7.1, the degeneracy in Theorem 7.2, the open source-value program, and the absence of a quantum transport theorem. $\square$

10 Reproducibility and remaining frontier

The finite results used here are archived in the curated calculation repository [5]. The principal result identifiers are:

- **A46:** `typed_family_representation`; the 48-state representation and anomaly table.
- **A47:** `native_gauge_group`; native $U(1)$, $SU(2)$, $SU(3)$ automorphisms and the $\mathbb{Z}_6$ quotient.
- **A49:** `physical_df_96`; the explicit profile 96×96 finite Dirac operator and finite-axiom tests.
- **A50:** `neutral_summand_hypercharge`; the neutral completion and unique anomaly-free hypercharge line.
- **A51:** `finite_inner_fluctuation`; the 676-pair one-form execution, selected one-Higgs projector, and finite traces.
- **A57:** `gauge_fixed_fluctuation_complex`; the selected gauge-fixed complex and exact one-loop coefficient vector.
- **A62:** `ten_spectrum_closure`; completion of all ten exact finite internal spectrum rows and the universal-shape limitation.

The next strict mathematical exits are:

1. construct the physical nonpullback q79 HYM endpoints and the connection-preserving continuum intertwiner;
2. derive the entries of D_F and the overlap kinetic metric without measured profile coordinates;
3. derive a sector-dependent physical threshold operator; the completed finite spectra alone cannot provide it;
4. select a proper-time measure and determine the identifiable spectral products from source geometry;
5. supply the Lorentzian/Wick and physical-state contracts; and
6. complete renormalized quantum transport with an uncertainty and observable-comparison theorem.

11 Conclusion

The corrected result is narrower than v3 and materially stronger than a disclaimer. MTT now has an executed finite almost-commutative profile encoding: the family and gauge carriers are correctly typed, the finite real-even triple is completed, the anomaly-free circle and global gauge group are explicit, and the finite one-form calculation explains why a selected projector is needed to obtain one Higgs doublet. The gauge-fixed complex now also derives the one-loop representation coefficients and closes all ten finite internal spectrum rows. The same analysis identifies exact obstructions: the qutrit algebra cannot simply be renamed as the Standard Model algebra, the uncompleted three-summand finite triple fails orientation and duality, the raw fluctuation space has three doublets, the normalized finite spectra cannot generate nonuniversal thresholds, and a single universal spectral normalization fails on the selected pure-SM running branch. What remains is not “derive NCG again.” It is to source the Euclidean/Lorentzian bridge, finite values, physical threshold operator, overlap metric, and spectral moments from one physical MTT geometry.

References

- [1] Ali H. Chamseddine and Alain Connes. The spectral action principle. *Communications in Mathematical Physics*, 186:731–750, 1997.
- [2] Alain Connes. *Noncommutative Geometry*. Academic Press, 1994.
- [3] Alain Connes and Matilde Marcolli. *Noncommutative Geometry, Quantum Fields and Motives*. American Mathematical Society, 2007.
- [4] Francesco D’Andrea, Maxim A. Kurkov, and Fedele Lizzi. Wick rotation and fermion doubling in noncommutative geometry. *Physical Review D*, 94:025030, 2016.
- [5] Peter Nero. MTT Results Repro: Curated theorem and calculation packets. <https://github.com/PeterNero/mtt-results-repro>, 2026. Authorities A44–A52 and their hash-addressed result packets, accessed 2026-07-28.
- [6] Walter D. van Suijlekom. *Noncommutative Geometry and Particle Physics*. Springer, 2015.
- [7] Dmitri V. Vassilevich. Heat kernel expansion: User’s manual. *Physics Reports*, 388:279–360, 2003.