

# Theta Closure in Modal Triplet Theory II: Direct Geometric Realization of Selected Gauge-Profile Overlaps

Peter Nero

July 2026

## Abstract

We re-execute the Route A geometric overlap ansatz against the selected multi-loop gauge-profile targets of Paper I. In its stated dimensionless normalization, the round- $S^2$  effective lens base and isotropic compact Heisenberg nilmanifold can be retargeted analytically to

$$(f_2 R_{\text{lens}})^2 = 0.2555137 R_1, \quad c = 0.9948493 R_1.$$

The associated spectral lower bounds remain above the assumed dimensionless admissibility floor. This proves existence of a calibrated representative in the declared auxiliary ansatz. It does not prove that the full lens space reduces to this  $S^2$  model, identify the Lens–Nil representative with the selected q79/Fu–Yau compactification, select a geometry uniquely by MTT, or make the gauge couplings held-out predictions. The shared circle is common  $U(1)$  phase/holonomy data, not an additional factor in a  $S^1 \times L(3, 1) \times \text{Nil}_3$  manifold.

## Revision note for this edition

**Supersedes.** *Theta Closure in Modal Triplet Theory II: Direct Geometric Realization of Non-abelian Overlaps*, first edition.

**Reason.** The previous numerical target came from the retired few-TeV crossing and the auxiliary Lens–Nil ansatz was liable to be read as the selected q79 compactification or a literal circle product.

**Resolution.** Version 2 retargets the analytic ansatz to the Paper I SMDR profile, counts the shared circle only as common holonomy data, and labels the round- $S^2$ /nil model auxiliary and calibrated.

**Retained result.** An explicit representative satisfying the two target ratios and dimensionless spectral-floor test exists in the declared ansatz.

**Remaining boundary.** No uniqueness, full lens reduction, q79/Fu–Yau identification, or held-out gauge prediction is established.

## 1 Purpose, central picture, and roadmap

Paper I supplies two dimensionless target ratios. This paper asks a deliberately narrow inverse question: can one choose the two scale parameters in a simple auxiliary geometry so that its overlap integrals reproduce those ratios while remaining above the declared spectral floor? The answer is yes.

The central picture is elementary. The weak coefficient is represented by the effective area of a round two-sphere, while the color coefficient is represented by the length of the central fiber of a compact Heisenberg nilmanifold. The shared-circle normalization fixes the common

denominator. Thus two measured target ratios determine two auxiliary geometric scales. This is an existence and calibration calculation, not a uniqueness argument and not a prediction of the measured ratios.

The dependency chain is

$$\begin{aligned} \text{Paper I target ratios} &\longrightarrow \text{two overlap equations} \longrightarrow \text{two auxiliary scales} \\ &\longrightarrow \text{spectral-floor check.} \end{aligned}$$

Sections 3 and 4 execute the two overlap equations separately. Section 5 states exactly what the construction establishes, and the appendix derives the nilmanifold bound used in the final admissibility check.

## 2 Relation to the revised $\Theta$ -closure core paper

Paper I now transports measured source coordinates with SMDR v1.3 to the full-Standard-Model  $\overline{\text{MS}}$  profile at  $Q = M_t$ . In the same GUT-normalized overlap convention it supplies

$$\frac{I_2}{I_1} = 0.5110273 \pm 0.0001231, \quad \frac{I_3}{I_1} = 0.158335 \pm 0.001098.$$

The scale  $Q = M_t$  is a matching convention, not an internal gap or coherence scale. This paper asks only whether the explicit Route A ansatz contains a representative with those overlap ratios and the stated dimensionless gap margin. Because the gauge rows select the target, success is a calibrated realization and round-trip test.

## 3 Overlap definitions

We recall the effective gauge-kinetic coefficients defined in Paper I:

$$I_a = \int_{B_a} w_a \langle \omega_a, \omega_a \rangle d\mu_{B_a}, \quad a = 1, 2, 3,$$

where  $B_a$  denotes a sector support after spectator integrations. The three supports are not asserted to be simultaneous Cartesian factors of one compactification. The same shared-circle phase datum is reused in all lanes and counted once.

In the auxiliary coefficient ansatz, the abelian coefficient  $I_1$  is fixed by the declared shared-circle normalization:

$$I_1 = 2\pi R_1.$$

We compute  $I_2^{(0)}$  and  $I_3^{(0)}$  below.

## 4 Lens sector: computation of $I_2^{(0)}$

**Dimensional and geometric convention.** All radii and overlaps below are dimensionless after one common internal length unit has been factored out. The “lens layer” used in the calculation is an effective two-dimensional constant-curvature base modeled by a round  $S^2$ ; it is not the full three-dimensional lens space  $L(3, 1)$ . A full lens-space realization requires a separate reduction theorem.

## 4.1 Lens geometry

The declared auxiliary lens-base ansatz uses the spectral bound

$$\lambda_{\text{lens}} \geq \frac{2}{(f_2 R_{\text{lens}})^2}.$$

*Remark 4.1* (Why the round  $S^2$  model is used here). The coefficient  $2/R^2$  is the exact first nonzero Laplace–Beltrami eigenvalue on the round two-sphere of radius  $R$ . Thus modeling the lens layer as a constant-curvature  $S^2$  scaled by the effective radius  $f_2 R_{\text{lens}}$  supplies one explicit geometry with the stated spectral bound. It is not selected uniquely by that eigenvalue.

The coefficient “2” motivates the round two-sphere as the baseline effective model within this ansatz; it is not a global uniqueness theorem. We take

$$\Sigma_2 \simeq S^2, \quad g_{\Sigma_2} = (f_2 R_{\text{lens}})^2 g_{S^2}.$$

## 4.2 Lens harmonic

The massless gauge representative is taken to have unit pointwise norm on  $\Sigma_2$  after gauge fixing. The shared-circle phase is a common spectator whose normalization is absorbed into Paper I’s coefficient convention. With the auxiliary weight  $w_2^{(0)} = 1$ , this gives

$$I_2^{(0)} = \text{Area}(\Sigma_2) = 4\pi(f_2 R_{\text{lens}})^2.$$

## 4.3 Matching to $\Theta$ -target

From Paper I,

$$\frac{I_2}{I_1} \approx 0.5110273, \quad I_1 = 2\pi R_1.$$

Hence

$$\frac{4\pi(f_2 R_{\text{lens}})^2}{2\pi R_1} = 0.5110273 \quad \Rightarrow \quad (f_2 R_{\text{lens}})^2 = 0.2555137 R_1.$$

**Proposition 4.2** (Lens overlap closure). *The lens overlap  $I_2^{(0)}$  matches the  $\Theta$ -target for  $(f_2 R_{\text{lens}})^2 = 0.2555137 R_1$ .*

## 4.4 Lens admissibility

The spectral bound gives

$$\lambda_{\text{lens}} \geq \frac{2}{0.2555137 R_1}.$$

Since admissibility of the circle sector requires  $R_1 \leq 2$ , we obtain

$$\lambda_{\text{lens}} \geq \frac{2}{0.2555137 \cdot 2} \approx 3.9137 \gg 0.25.$$

Thus the lens sector is safely admissible.

# 5 Nil sector: computation of $I_3^{(0)}$

## 5.1 Nil geometry

We take  $\Sigma_3$  to be the standard compact Heisenberg nilmanifold  $\Gamma \backslash \text{Nil}_3$  with integer lattice and left-invariant metric

$$g_{\text{nil}} = a^2 \sigma_1^2 + b^2 \sigma_2^2 + c^2 \sigma_3^2,$$

where

$$\sigma_1 = dx, \quad \sigma_2 = dy, \quad \sigma_3 = dz - x dy.$$

## 5.2 Canonical massless harmonics

The first cohomology is generated by the closed forms  $dx$  and  $dy$ . We choose the isotropic convention

$$a = b,$$

which renders the overlap independent of the basis in  $\text{span}\{dx, dy\}$ .

## 5.3 Overlap computation

For  $\omega^{(0)} = dx$  or  $dy$ , auxiliary weight  $w_3^{(0)} = 1$ , and the unit coordinate fundamental domain,

$$|\omega^{(0)}|^2 = \frac{1}{a^2}, \quad d\text{vol} = a^2 c (\sigma_1 \wedge \sigma_2 \wedge \sigma_3).$$

Hence

$$I_3^{(0)} = \int |\omega^{(0)}|^2 d\text{vol} = c.$$

## 5.4 Matching to $\Theta$ -target

From Paper I,

$$\frac{I_3}{I_1} \approx 0.158335 \quad \Rightarrow \quad I_3 \approx 0.9948493 R_1.$$

Therefore

$$\boxed{c = 0.9948493 R_1.}$$

**Proposition 5.1** (Nil overlap closure). *For the isotropic nil metric  $a = b$  with  $c = 0.9948493 R_1$ , the leading-order nil overlap  $I_3^{(0)}$  matches the  $\Theta$ -target.*

## 5.5 Nil admissibility

The scalar Laplacian on  $\Gamma \backslash \text{Nil}_3$  has spectrum

$$\lambda_1 \geq \min \left\{ \frac{4\pi^2}{a^2}, \frac{2\pi}{a^2} + \frac{4\pi^2}{c^2} \right\}.$$

With  $a = b = 1$  and  $c \leq 1.989699$  (from  $R_1 \leq 2$ ),

$$\lambda_1 \geq \min \left\{ 4\pi^2, 2\pi + \frac{4\pi^2}{(1.989699)^2} \right\} \approx 16.2553 \gg 0.25.$$

Thus the nil sector satisfies the MTT admissibility bound  $\lambda_{\text{nil}} \geq 0.25$  with large margin.

## 6 Interpretation of the Route-A result and scope

Within the declared dimensionless ansatz, the selected profile targets admit an explicit lens-base/nil representative and the assumed spectral inequalities remain satisfied. This closes the algebraic retargeting and ansatz-level existence check. It does not close source selection of the geometry, a literal  $L(3, 1)$  reduction, global HYM connection data, or a held-out gauge-coupling prediction. In particular, this auxiliary representative is not the q79 Fu–Yau compactification and supplies no theorem identifying the two spaces.

## 7 Worked case: unit abelian radius

Taking  $R_1 = 1$  makes the normalization and the role of each parameter transparent:

$$f_2 R_{\text{lens}} = \sqrt{0.2555137} \approx 0.505484, \quad c = 0.9948493.$$

The three overlap coefficients are then

$$I_1 = 2\pi \approx 6.283185, \quad I_2 = 4\pi(0.2555137) \approx 3.210880, \quad I_3 = 0.9948493.$$

Consequently

$$\frac{I_2}{I_1} = 0.5110274, \quad \frac{I_3}{I_1} = 0.1583352,$$

up to the displayed rounding. The lens lower bound is approximately 7.8274. For the nilmanifold, the torus branch gives  $4\pi^2 \approx 39.4784$  and the Landau branch is larger, so the lower bound is 39.4784. This example shows concretely how the target ratios determine the auxiliary scales; it does not turn those fitted scales into independent predictions.

## A Spectral lower bound on $\Gamma \backslash \text{Nil}_3$ for the left-invariant metric

This appendix justifies the spectral estimate used in the nil admissibility check.

### A.1 Heisenberg nilmanifold and left-invariant metric

Let  $\text{Nil}_3$  be the Heisenberg group with global coordinates  $(x, y, z) \in \mathbb{R}^3$  and left-invariant 1-forms

$$\sigma_1 = dx, \quad \sigma_2 = dy, \quad \sigma_3 = dz - x dy,$$

satisfying  $d\sigma_1 = d\sigma_2 = 0$  and  $d\sigma_3 = -\sigma_1 \wedge \sigma_2$ .

Let  $\Gamma$  be the standard integer lattice so  $\Gamma \backslash \text{Nil}_3$  is compact. Equip it with the left-invariant metric

$$g = a^2 \sigma_1^2 + b^2 \sigma_2^2 + c^2 \sigma_3^2, \quad a, b, c > 0.$$

The dual left-invariant vector fields are

$$X = \partial_x, \quad Y = \partial_y + x \partial_z, \quad Z = \partial_z,$$

with  $[X, Y] = Z$  and other brackets zero. An orthonormal frame is

$$e_1 = \frac{1}{a} X, \quad e_2 = \frac{1}{b} Y, \quad e_3 = \frac{1}{c} Z.$$

### A.2 Scalar Laplacian

For a left-invariant orthonormal frame, the scalar Laplacian on functions may be written as

$$\Delta f = -(e_1^2 + e_2^2 + e_3^2)f$$

(up to sign convention; we use the nonnegative operator  $-\sum e_i^2$ ). Substituting the frame gives

$$\Delta = -\frac{1}{a^2} \partial_x^2 - \frac{1}{b^2} (\partial_y + x \partial_z)^2 - \frac{1}{c^2} \partial_z^2. \quad (\text{A.1})$$

### A.3 Fourier decomposition in the central direction

Because  $z$  is periodic on  $\Gamma \backslash \text{Nil}_3$ , expand

$$f(x, y, z) = \sum_{p \in \mathbb{Z}} e^{2\pi i p z} g_p(x, y).$$

On the  $p$ -th mode,  $\partial_z \mapsto 2\pi i p$  and (A.1) becomes

$$\Delta_p = -\frac{1}{a^2} \partial_x^2 - \frac{1}{b^2} (\partial_y + 2\pi i p x)^2 + \frac{(2\pi p)^2}{c^2}. \quad (\text{A.2})$$

We bound the lowest nonzero eigenvalue by analyzing  $p = 0$  and  $p \neq 0$  separately.

### A.4 The $p = 0$ sector (torus bound)

For  $p = 0$ ,

$$\Delta_0 = -\frac{1}{a^2} \partial_x^2 - \frac{1}{b^2} \partial_y^2$$

on the flat 2-torus  $(x, y) \in (\mathbb{R}/\mathbb{Z})^2$ . Its eigenvalues are

$$\lambda_{m,n}^{(0)} = 4\pi^2 \left( \frac{m^2}{a^2} + \frac{n^2}{b^2} \right), \quad (m, n) \in \mathbb{Z}^2.$$

Hence the smallest nonzero eigenvalue in this sector satisfies

$$\lambda_{\min}^{(0)} \geq \frac{4\pi^2}{\max(a^2, b^2)}. \quad (\text{A.3})$$

### A.5 The $p \neq 0$ sector (Landau/oscillator lower bound)

For  $p \neq 0$ , the operator (A.2) is the magnetic Laplacian on a 2-torus with constant magnetic field proportional to  $p$ , plus a strictly positive shift  $(2\pi p)^2/c^2$ . A conservative lower bound follows by dropping the nonnegative  $x$ -dependence and using the lowest Landau-level scale:

$$\lambda_{\min}^{(p)} \geq \frac{2\pi|p|}{ab} + \frac{(2\pi p)^2}{c^2}, \quad p \neq 0. \quad (\text{A.4})$$

(For our purposes it suffices to retain the  $|p| = 1$  bound.)

Under the isotropic convention  $a = b$ , this yields

$$\lambda_{\min}^{(1)} \geq \frac{2\pi}{a^2} + \frac{4\pi^2}{c^2}. \quad (\text{A.5})$$

### A.6 Combined lower bound

Combining (A.3) and (A.5), we obtain

$$\lambda_1(\Delta_{\Gamma \backslash \text{Nil}_3}) \geq \min \left\{ \frac{4\pi^2}{a^2}, \frac{2\pi}{a^2} + \frac{4\pi^2}{c^2} \right\}, \quad (a = b). \quad (\text{A.6})$$

In particular, for  $a = b = 1$  and any  $c \leq 1.989699$ , one has

$$\lambda_1 \geq 2\pi + \frac{4\pi^2}{c^2} \geq 2\pi + \frac{4\pi^2}{(1.989699)^2} \approx 6.2832 + 9.9721 \approx 16.2553,$$

which is much larger than the admissibility floor 0.25 used in Paper I.

*Remark A.1.* The estimates above are intentionally conservative and suffice only to demonstrate a large spectral margin. A complete scalar spectrum for the three-dimensional Heisenberg nilmanifold is available in Ref. [1], but is not required for the present admissibility verification.

## B Conclusion

The Route A formulas have been evaluated in the same normalization and scheme as revised Paper I. The new central representative is

$$(f_2 R_{\text{lens}})^2 = 0.2555137 R_1, \quad c = 0.9948493 R_1,$$

with uncertainties inherited from the two profile ratios. The nil spectral estimate and lens lower bound continue to exceed the assumed dimensionless floor for the stated parameter range.

The earlier numbers  $0.280 R_1$  and  $1.439 R_1$  belonged to the withdrawn 5 TeV profile and are not retained. The present construction is a calibrated existence result: the target gauge rows were used to determine the geometry. Selection of this geometry before empirical comparison, literal lens-space reduction, and a global HYM representative remain separate proof obligations.

## Computational Evidence and Reproducibility

The HYM contraction, finite-matrix realization, precision transport, and electroweak row are used directly to assess the selected overlap realization. The remaining profile packets locate that realization within the broader closure program but do not prove its geometry. The strict source upgrade remains open.

The referenced rows are frozen to the curated results repository state identified below. Hashes are grouped in eight-character blocks for line breaking.

**Repository:** <https://github.com/PeterNero/mtt-results-repro>

**Commit:** 31247ebb 5c22f3fb b5443024 365433c6 ee0bff4a

**Manifest:** release/result\_manifest.json

**Manifest SHA-256:**

fb399689 60b00584 631dbf53 1a708e18

ef928d6b 6d935119 c185d7f6 32b1e7cd

Tier labels are quoted verbatim from that manifest. A row used directly supports only the specific computational statement identified above; a corpus-state cross-check does not prove this paper's local theorems; and an open row is evidence of an unresolved obligation, never of closure.

### Rows used directly in this paper.

- **A19/hym\_wiener\_contraction** (*numeric certified*).  
Weighted-theta Fourier-tail and Wiener contraction certificate.
- **A02/precision\_15\_source\_transport** (*profile replay*).  
Fifteen measured source coordinates, Jacobian and covariance transport.
- **A02/precision\_8x8\_workspace** (*profile replay*).  
Eight-coordinate SMDR output with positive-definite 8x8 covariance.
- **A01/qutrit\_weyl\_27\_matrix** (*derived exact*).  
Sparse 27x27 qutrit-Weyl left-action realization.
- **A01/strict\_pew\_row** (*derived exact*).  
Promoted P\_EW source row at the declared one-shared-primitive standard.

### Corpus-state cross-checks.

- A01/charged\_yukawa\_higgs\_profile (*profile replay*).  
Versioned Yu, Yd, Ye and lambda\_H profile packet.
- A14/ckm\_prediction\_profile (*numeric certified*).  
Three selected CKM profile rows and uncertainty comparison.
- A01/current\_global\_lock (*profile replay*).  
Current non-looping global status and source-certificate map.
- A01/direct\_k\_higgs\_row (*derived exact*).  
Promoted direct K\_threshold.Omega\_H.lambda row.
- A22/e6\_qpsi\_qcd\_anomaly (*derived exact*).  
E6 Qpsi matter/exotic QCD anomaly cancellation audit.
- A04/final\_12\_of\_12\_audit (*profile replay*).  
Twelve-obligation embedded renormalized-SM equivalence audit.
- A40/neutral\_two\_primitive\_profile (*profile replay*).  
Measured two-splitting neutrino profile, masses and Dirac Yukawa rows.

### Open boundary (not evidence of closure).

- A05/strict\_upgrade\_ledger (*open*).  
Current 2/9 strict no-knob upgrade ledger.

No imported row changes theorem ownership or promotes a neighboring claim: all local statements retain their stated hypotheses, domains, and limitations.

## References

- [1] D. Andriot and D. Tsimpis, *Laplacian spectrum on a nilmanifold, truncations and effective theories*, JHEP **09** (2018) 096. [https://doi.org/10.1007/JHEP09\(2018\)096](https://doi.org/10.1007/JHEP09(2018)096)
- [2] P. Nero, *Modal Triplet Theory: Admissibility, Encodings, and the Structure of Physical Description*, Zenodo preprint, January 2026. <https://doi.org/10.5281/zenodo.18255621>
- [3] P. Nero, *Modal Triplet Theory: Foundation*, Zenodo preprint, September 2025. <https://doi.org/10.5281/zenodo.16949762>
- [4] P. Nero, *Fixed Points I–VI: Complete Coherence Spine*, Zenodo preprints, August 2025. <https://doi.org/10.5281/zenodo.16948748>
- [5] P. Nero, *The Projection–Admissibility Principle: Structural Constraints on Effective Physical Description*, Zenodo preprint, January 2026. <https://doi.org/10.5281/zenodo.18255838>
- [6] P. Nero, *Closure and Inevitability in Modal Triplet Theory*, Zenodo preprint, January 2026. <https://doi.org/10.5281/zenodo.18255510>
- [7] P. Nero, *Coherence Capacity as the Fundamental Resource of Effective Physics*, Zenodo preprint, January 2026. <https://doi.org/10.5281/zenodo.18255905>
- [8] P. Nero, *Dynamics of Coherence Capacity: Transport, Concentration, and Exhaustion*, Zenodo preprint, January 2026. <https://doi.org/10.5281/zenodo.18256048>

- [9] P. Nero, *Modal Triplet Theory: From MTT to Quantum Mechanics*, Zenodo preprint, September 2025. <https://doi.org/10.5281/zenodo.17074246>
- [10] P. Nero, *From MTT to Quantum Field Theory*, Zenodo preprint, 2025. <https://doi.org/10.5281/zenodo.17068816>
- [11] P. Nero, *Modal Triplet Theory: From MTT to General Relativity*, Zenodo preprint, October 2025. <https://doi.org/10.5281/zenodo.16950597>
- [12] P. Nero, *Modal Triplet Theory: From MTT to a UV-Finite, Unitary Quantum Gravity*, Zenodo preprint, 2025. <https://doi.org/10.5281/zenodo.17077671>
- [13] P. Nero, *Measurement as Disturbance and Stabilization in Modal Triplet Theory*, Zenodo preprint, 2025. <https://doi.org/10.5281/zenodo.17177404>
- [14] P. Nero, *Projection, Probability, and Irreversibility: Shadow Bridges Between Measurement, Black Holes, and Cosmology in Modal Triplet Theory*, Zenodo preprint, January 2026. <https://doi.org/10.5281/zenodo.18256408>
- [15] P. Nero, *Modal Fixed Points, Bell's Beables, and the Limits of Factorization*, Zenodo preprint, 2025. <https://doi.org/10.5281/zenodo.17076300>
- [16] P. Nero, *Temporal Bell Inequalities and Global Consistency in Modal Triplet Theory*, Zenodo preprint, August 2025. <https://doi.org/10.5281/zenodo.18208884>
- [17] P. Nero, *Modal Triplet Theory and History-Dependent Stochastic Processes: Fixed-State Indivisibility, Markov Order, and the Quantum Boundary*, Zenodo preprint, January 2026. <https://doi.org/10.5281/zenodo.18254862>