

Universal Perturbative Graph Structure:

A Typed Modal Diagrammatics and Its Quantum Boundary

Peter Nero

Version 3, September 2026

Abstract

Feynman graphs are used in quantum field theory, classical statistical field theory, stochastic dynamics, and effective descriptions of gravity. Their recurrence has a precise but limited explanation. Once one specifies a nondegenerate quadratic kernel, interaction tensors, a grading, and a contraction rule, Wick expansion organizes perturbative coefficients by graphs. The graph grammar is therefore not uniquely quantum. It also does not, by itself, select a physical propagator, Bose or Fermi statistics, causal locality, gauge reduction, counterterms, positivity, or unitarity.

This paper reformulates modal diagrammatics around that distinction. We prove a finite graded graph-expansion theorem and an exact projection theorem: if an even projector intertwines the quadratic operator, interaction tensors, grading, and declared identities, then every all-retained graph coefficient agrees with the coefficient computed in the projected theory. We also prove a nonselection theorem showing that the same graph combinatorics supports continuously many inequivalent weights and different statistics. Thus basin adjacency or coherence language alone cannot derive Feynman rules.

For Modal Triplet Theory, the result supplies a rigorous transfer target. The selected q79 twisted-Dirac construction already composes with standard CAR/AQFT machinery to give a free even local net. The present theorem explains how a selected upper action could transfer its perturbative graph data to a lower effective sector. It does not manufacture that action. The geometry-selected upper action, fixed-coupling interacting gauge-BRST C^* -completion, renormalization-group matching, and observable uncertainty packet remain open. The formulation retains the explanatory value of modal diagrammatics while making its mathematical and physical boundary explicit. A finite residual-and-metric jet supplies the complete repair-action graph datum, with exact transport under a compatible change of variables; its identification with physical Feynman data remains a separate action and state problem.

Version 3 revision note

Supersedes. Version 2; the earlier revision note and released identity are retained.

Reason. The residual-to-graph construction now supplies an upstream input to the transfer theorem. The graph formula also needed an explicit interaction sign and correctly typed field/dual projection.

Resolution. Explain the complete residual/metric jet, a polynomial example, and non-isometric coordinate transport. Distinguish cost pullback from reducing isometry and include the Euclidean vertex sign.

Retained result. Finite Wick organization, all-retained graph transfer, no-mixed-vertex factorization, and graph nonselection remain exact.

Open boundary. A positive repair cost does not select the physical signed action, causal inverse, statistics, state, or interacting completion.

Version 2 revision note

Supersedes. Version 1.0 (DOI [10.5281/zenodo.18330804](https://doi.org/10.5281/zenodo.18330804)).

Reason. Version 1 correctly emphasized that diagrammatic expansions are not exclusive to quantum theories, but it attributed too much to coherence and Gaussian combinatorics. In several places it treated a modal Hessian as a selected physical propagator, described projected diagrams as necessarily the standard Feynman rules, and discussed quantum behavior and gravity beyond what the supplied structure proved.

Resolution. Version 2 makes four substantive changes.

1. It replaces the broad origin claim by exact finite graded graph-expansion and projection theorems.
2. It distinguishes the quadratic inverse, Euclidean covariance, retarded/advanced Green operators, causal propagator, state two-point function, and Feynman kernel.
3. It proves that graph incidence data do not select numerical weights, statistics, gauge identities, counterterms, or a quantum theory.
4. It reconciles the paper with the current MTT ledger: the selected free q79 even-CAR net is closed at its declared tier, while the upper action and nonperturbative interacting completion remain open.

These are corrections to the central claim, not merely added caveats.

Retained result. The original observation that classical and quantum models can share the same perturbative graph grammar is retained and is now proved at the exact finite graded level.

Remaining boundary. A selected upper action, its complete projection intertwiners, and a fixed-coupling interacting gauge-BRST C^* -completion are not established here.

1 Why the same diagrams recur

A perturbative graph is a compact record of two operations:

1. insert multilinear interaction tensors;
2. contract their arguments pairwise with a chosen two-point kernel.

The record is useful because the number of pairings grows rapidly. Wick's theorem is the algebraic mechanism that turns those pairings into graph sums [15]. Nothing in that combinatorial statement alone decides whether the underlying system is a Euclidean random field, a Lorentzian quantum field, a response theory, or a finite algebraic model.

This observation motivates, but does not yet prove, the MTT interpretation. A fixed point can provide a background about which to expand. A projector can identify retained modes. Neither object determines the action, its quadratic inverse, or its higher derivatives.

Those data must be supplied or derived separately. The useful question is therefore not *whether coherence automatically creates Feynman rules*, but:

Under which explicit hypotheses does perturbative graph data pass from an upper description to a projected effective description?

The answer below is exact on a declared finite or regularized domain. It is intentionally not a claim about convergence of an infinite perturbation series or existence of an interacting continuum QFT.

2 The typed perturbative input

2.1 Finite graded field space

Let

$$E = E_{\bar{0}} \oplus E_{\bar{1}}$$

be a finite-dimensional real or complex super vector space. Even coordinates represent bosonic directions and odd coordinates represent fermionic directions. Fix:

1. an even nondegenerate quadratic operator $K : E \rightarrow E^*$;
2. its inverse contraction kernel $G = K^{-1} : E^* \rightarrow E$;
3. graded-symmetric interaction tensors

$$V_n \in (E^*)^{\otimes n}, \quad 3 \leq n \leq N;$$

4. a formal coupling parameter λ ;
5. an ordering and orientation convention for odd half-edges.

The formal action is

$$S(\phi) = \frac{1}{2} K(\phi, \phi) + \sum_{n=3}^N \frac{\lambda^{n-2}}{n!} V_n(\phi, \dots, \phi). \quad (2.1)$$

The powers of λ are conventional and may be replaced by independent formal couplings without changing the argument.

The word “typed” matters. The pair (K, V_n) does not yet define a physical QFT. It defines algebraic graph weights. A physical interpretation requires additional structures described in [section 6](#).

2.2 Graph weights

Let Γ be a finite graph whose vertices have valence between 3 and N . Assign $V_{\deg(v)}$ to every vertex and G to every internal edge. Contract matching tensor indices. For odd labels, include the Koszul sign $\epsilon(\Gamma)$ determined by the declared ordering. The resulting scalar or external tensor is

$$W_\Gamma(K, V, G) = \frac{(-1)^{|V(\Gamma)|} \epsilon(\Gamma)}{|\text{Aut } \Gamma|} \text{Contr}_\Gamma \left(\bigotimes_{v \in V(\Gamma)} V_{\deg(v)} \otimes \bigotimes_{e \in E_{\text{int}}(\Gamma)} G \right). \quad (2.2)$$

This formula exposes all of the information that a bare graph drawing hides: numerical kernels, vertex tensors, symmetry factors, and grading. The displayed sign is for the Euclidean convention $\exp(-S_{\text{int}})$ with $\hbar = 1$; coupling powers are understood. Other exponential

conventions replace that vertex factor explicitly. External insertions are labeled and graph automorphisms fix those labels. This formula first organizes unnormalized moments; division by the vacuum series gives normalized correlators and removes disconnected vacuum factors.

Definition 2.1 (Perturbative graph datum). A perturbative graph datum is the tuple

$$\mathfrak{D} = (E, \text{Gr } E, K, G, \{V_n\}, \epsilon, \mathcal{R}),$$

where $\mathcal{R}$ records any regularization or renormalization map needed to define singular contractions. In a finite nonsingular model, $\mathcal{R} = \text{Id}$.

3 The graph expansion theorem

Theorem 3.1 (Finite graded Wick graph expansion). *Let E , K , and V_n be as above. Treat the Gaussian expectation with covariance $G = K^{-1}$ and the interaction exponential as a formal power series. Every coefficient of every polynomial correlation function is a finite sum of the weights (2.2). The vertices are the tensors V_n , the internal edges are contractions by G , the automorphism denominator gives the symmetry factor, and the odd permutation convention gives the fermionic sign.*

Proof. Expand the interaction exponential order by order in λ . At a fixed order only finitely many products of the V_n occur. Apply the graded Wick rule to each polynomial Gaussian expectation. Every complete pairing identifies two tensor arguments and contracts them with G . Grouping pairings that differ only by relabeling yields an isomorphism class of graphs and the orbit-stabilizer factor $|\text{Aut } \Gamma|^{-1}$. Moving odd variables into the chosen pairing order produces the Koszul sign, while the interaction exponential contributes one minus sign per vertex. Conversely, every decorated graph determines such a pairing. Thus the two finite sums agree coefficient by coefficient. $\square$

Remark 3.2. The theorem is exact as a statement about formal coefficients. It does not assert convergence, reflection positivity, causal factorization, unitarity, or existence of a continuum measure.

3.1 What is universal

The universal content is now precise:

$$\begin{aligned} & \text{quadratic contraction} + \text{multilinear interactions} + \text{grading} \\ \implies & \text{graph-indexed formal coefficients.} \end{aligned}$$

Classical Gaussian measures and quasifree quantum states both instantiate this schema. Their physical meanings remain different because their algebras, kernels, state conditions, and localization rules differ.

4 Exact transfer through a finite projection

4.1 Compatible retained sector

Let $P : E \rightarrow E$ be an even idempotent and write $E_P = \text{im } P$. Let $P^* : E^* \rightarrow E^*$ be its dual. Because $K : E \rightarrow E^*$ is a quadratic form, the correctly typed block-diagonality condition is

$$KP = P^*K. \tag{4.1}$$

Then K is block diagonal on $E_P \oplus \ker P$, and

$$K_P = K|_{E_P}, \quad G_P = K_P^{-1} = PGP^*|_{E_P^*}.$$

Here E_P^* is embedded by the splitting determined by P . If a field metric identifies E with E^* and P is orthogonal for that metric, this condition becomes commutation of P with the Riesz operator of K . Define the retained vertex

$$V_n^P = V_n|_{E_P^{\otimes n}}.$$

Theorem 4.1 (All-retained graph transfer). *Under (4.1), let Γ be a decorated graph for which every half-edge label lies in E_P . Then*

$$W_\Gamma(K_P, V^P, G_P) = W_\Gamma(K, V, G)|_{E_P}.$$

The equality includes symmetry factors and graded signs.

Proof. At every retained vertex, V_n^P is exactly the restriction of V_n . On every retained internal edge, commutation and block diagonality give $G_P = PGP^*|_{E_P^*}$. Since P is even, it preserves parity and hence the Koszul sign. The underlying graph and its automorphism group are unchanged. Substitution into (2.2) proves the equality. $\square$

Corollary 4.2 (Closed retained subtheory). *If, in addition, every mixed interaction component vanishes,*

$$V_n(x_1, \dots, x_n) = 0 \quad \text{whenever retained and discarded arguments both occur,}$$

then the full formal perturbative expansion with retained external legs factorizes from the discarded sector. After normalization by vacuum terms, it equals the expansion computed solely from $(E_P, K_P, \{V_n^P\})$.

Proof. Condition (4.2) prevents a connected graph with retained external legs from containing a discarded internal label. Apply theorem 4.1 to every connected contribution. Purely discarded vacuum graphs factor and cancel in normalized correlators. $\square$

4.2 Why effective actions are needed

Without (4.2), the theorem does *not* say that simply deleting discarded modes gives the correct lower theory. Mixed vertices allow discarded internal lines to contribute to retained observables. Integrating them out produces an effective action

$$e^{-S_{\text{eff}}(\phi_P)} = \int_{\ker P} e^{-S(\phi_P + \chi)} d\chi$$

in a Euclidean finite model, or its appropriate formal/algebraic analogue. That effective action usually contains new nonlocal or higher-order vertices. Projection and marginalization are therefore different operations.

4.3 Transfer of identities

Proposition 4.3 (Intertwined identity transfer). *Let $\mathcal{A}$ and $\mathcal{A}_P$ be algebras of formal functionals, let $R : \mathcal{A} \rightarrow \mathcal{A}_P$ be the restriction induced by P , and let $\mathcal{W}, \mathcal{W}_P$ be linear identity operators. If*

$$R\mathcal{W} = \mathcal{W}_P R,$$

then $\mathcal{W}F = 0$ implies $\mathcal{W}_P(RF) = 0$.

Proof. Apply R to $WF = 0$ and use the intertwining equality. $\square$

This proposition is elementary, but it identifies the real obligation in gauge theory. Ward, BRST, or BV identities descend only after their operators and brackets are shown to intertwine. A mode count or a graph match does not establish this.

5 The nonselection theorem

Theorem 5.1 (Graph combinatorics does not select Feynman rules). *The incidence structure of perturbative graphs, even together with a list of allowed valences, does not determine:*

1. *propagator values;*
2. *interaction strengths;*
3. *bosonic or fermionic statistics;*
4. *causal or boundary prescriptions;*
5. *gauge-fixing, ghost, or BRST data;*
6. *renormalization extensions or counterterms.*

In particular, continuously many inequivalent perturbative graph data have the same underlying graph set.

Proof. Take the one-dimensional bosonic action

$$S_{k,g}(x) = \frac{k}{2}x^2 + \frac{g}{4!}x^4, \quad k > 0.$$

For every k and $g \neq 0$, the allowed connected graphs are the same four-valent graphs. A graph with I internal edges and V vertices has weight proportional to $k^{-I}g^V$. Varying k or g therefore changes all nontrivial weights without changing incidence data. Replacing the field by an odd multiplet changes exchange signs while the ungraded incidence graph can remain identical. Retarded, advanced, Euclidean, and Feynman prescriptions may solve equations associated with the same differential expression but obey different support or boundary conditions. Gauge fixing adds ghost and identity data, and renormalization chooses extensions of singular distributions. None is encoded by the bare graph set. $\square$

Corollary 5.2 (Basin adjacency is insufficient). *A basin graph, neighborhood relation, or coherence-capacity network cannot by itself derive physical Feynman rules. It must be supplemented by a selected quadratic operator and inverse prescription, interaction tensors, grading, gauge structure where applicable, and a renormalization rule.*

6 Which kernel is meant?

Version 1 used “propagator” too broadly. The following objects can be related, but they are not interchangeable.

Object	Defining input	What it does not automatically provide
Algebraic inverse K^{-1}	Nondegenerate finite quadratic form	Causality, positivity, or a quantum state
Euclidean covariance	Positive elliptic operator and measure	Lorentzian time ordering without reconstruction hypotheses
Retarded/advanced Green operator	Green-hyperbolic operator and support condition	A state two-point function
Causal propagator	Difference of retarded and advanced operators	Positive-frequency or Hadamard choice
Two-point function	Observable algebra and state	Time ordering or unique vacuum
Feynman kernel	Time ordering/state or boundary prescription	Nonperturbative completion

On curved spacetime, local covariant Wick products and time-ordered products require microlocal and renormalization conditions [1, 3, 4]. Epstein–Glaser renormalization shows that locality and causal factorization constrain the extension of distributions [2]. These are physical and analytic inputs, not consequences of graph drawing.

7 Gauge theories and renormalization

7.1 Gauge-fixed perturbation theory

For a gauge theory the naive Hessian is degenerate along gauge orbits. One must introduce gauge-fixing and ghost sectors, or work in a homological BV/BRST formulation. The interaction data then include graded fields, antifields, a bracket, and a differential. The quantum master equation or corresponding Ward identities control the consistency of the perturbative expansion [14].

Modal projection is compatible with such a construction only if the projection preserves the relevant complex and identities. Schematically,

$$PQ = QP, \quad P\{\cdot, \cdot\} = \{P\cdot, P\cdot\}_P, \quad RW = \mathcal{W}_P R.$$

These equations are proof obligations. Calling discarded directions “inadmissible” does not make them true.

7.2 Renormalized graph weights

In a continuum theory, products of propagators can be singular on coincident-point diagonals. A graph before extension may define a distribution only away from those diagonals. Renormalization extends it subject to locality, covariance, scaling, and symmetry requirements. Different allowed extensions correspond to finite renormalization freedoms. Thus the renormalization map $\mathcal{R}$ belongs in the perturbative graph datum.

The graph theorem remains useful: it organizes which distribution must be extended. It does not perform or uniquely select the extension.

8 Repair jets supply a finite graph datum

The frozen H4-T8 construction supplies an upstream arrow to this paper’s graph theorem [11]. Fix finite field and residual spaces, a background a_* , a smooth or formal residual $\Phi(a)$,

and a positive residual metric field $W(a)$. Set

$$S_{\text{rep}}(a) = \frac{1}{2} \langle \Phi(a), W(a) \Phi(a) \rangle.$$

Write $R_m = D^m \Phi(a_*)$ and $W_m = D^m W(a_*)$. Repeated Leibniz differentiation assigns each derivative to the left residual, metric, or right residual, so the entire vertex tensor is

$$V_n(v_1, \dots, v_n) = \frac{1}{2} \sum_{[n]=A \sqcup B \sqcup C} \langle R_{|A|}(v_A), W_{|B|}(v_B) R_{|C|}(v_C) \rangle.$$

The blocks are ordered by their factor roles and indices retain their original order; graded models insert the corresponding Koszul signs. Thus vertices are outputs of the joint residual-and-metric jet, not extra source coefficients. A residual without its metric derivatives is insufficient. At $R_0 = 0$, the Hessian form is $K = R_1^\dagger W_0 R_1$. At nonzero defect even constant W adds $\langle R_0, W_0 R_2(v, w) \rangle$, so the Gram formula is no longer complete.

8.1 A worked polynomial source

For $\Phi(x, y) = (x, y + y^2)$ and $W = I$ at zero,

$$S_{\text{rep}} = \frac{1}{2}x^2 + \frac{1}{2}y^2 + y^3 + \frac{1}{2}y^4, \quad K = I, \quad V_{3,yyy} = 6, \quad V_{4,yyyy} = 12.$$

All higher vertices vanish. In general a residual of degree r with constant metric has no vertices above degree $2r$. With the same scalar residual and $W(y) = 1 + y + y^2 > 0$, the cubic and quartic derivatives instead become 9 and 48, and the action has degree six. This is a metric-jet effect, not new independent interaction data.

In the unit-covariance Gaussian, $\langle y^4 \rangle = 3$ and $\langle y^6 \rangle = 15$. The unnormalized vacuum coefficients from one quartic vertex and two cubic vertices are consequently $-(12/4!)3 = -3/2$ and $(6/3!)^2 15/2 = 15/2$. These illustrate the exponential sign as well as the combinatorics.

8.2 Coordinate transport is not a physical isometry

If linear bijections T, U satisfy $\Phi'(Ta) = U\Phi(a)$ and $\langle Ur, W'(Ta)Us \rangle = \langle r, W(a)s \rangle$, then $S'_{\text{rep}}(Ta) = S_{\text{rep}}(a)$ and every vertex pulls back. For $T = \text{diag}(2, 3)$ in the example, the target Hessian form is $\text{diag}(1/4, 1/9)$, its contravariant Gaussian covariance is $\text{diag}(4, 9)$, and its cubic/quartic entries are $2/9, 4/27$. Edge and vertex factors cancel exactly in every contraction. Transporting the field metric also makes the Hessian Riesz operators and their functional calculi intertwine. With independently fixed physical metrics, a cost pullback alone does not prove that stronger statement.

Nor is a lower-dimensional cost restriction a reducing embedding. For $K = \begin{pmatrix} 1 & 1 \\ 1 & 2 \end{pmatrix}$ and $Tx = (x, 0)$, the restricted Hessian is $T^*KT = 1$, but $KT(1) = (1, 1) \neq T(1)$. The retained line leaks into its complement. The all-retained theorem therefore still needs (4.1), and the full retained theory still needs the no-mixed-vertex condition or an effective-action calculation.

8.3 What the source reduction does not quantize

On a specified kernel complement, the repair Hessian yields a finite Euclidean reduced inverse, zero-mode projector, and heat semigroup. These are sufficient for formal repair graphs. The joint q79 residual can organize integrability, HYM, balance, anomaly, and normalization lanes, but its actual endpoint and metric jets are not evaluated by H4-T8. In particular the signed-action comparison in the action paper [6] is still necessary: squaring an action

gradient changes both its Hessian and its vertices. Neither a causal/Feynman prescription nor a physical gauge/BRST state is obtained by differentiating a positive norm. The selected free-CAR and operational-QM results remain at their established tiers; this construction does not reopen them or complete the interacting theory.

9 What MTT contributes now

9.1 Fixed points and coherent sectors

MTT supplies a language of coherent fixed points, admissible sectors, and finite projections. At the level needed here, a fixed point identifies a background u_* , and a selected action S would yield

$$K = d^2 S[u_*], \quad V_n = d^n S[u_*].$$

If S , u_* , and the inverse prescription are selected from one source, the tensors in the graph datum are no longer arbitrary. If only the fixed point or carrier is known, they remain underdetermined.

This is the significance of the current upper-action blocker. The missing theorem is not another restatement of Wick expansion. It is a same-source construction of an upper differential/action whose Hessian, higher products, symmetries, and projections reproduce the accepted lower structures.

9.2 The selected free q79 CAR result

The companion QFT paper owns the current selected free-field theorem [10]. On the declared globally hyperbolic framed q79 representative, the selected twisted massless Dirac operator composes with standard Green-hyperbolic and CAR/AQFT machinery to produce an even local observable net with locality, covariance, the time-slice property, and a nonempty positive Hadamard state space.

The present paper does not re-prove that theorem. It uses it to mark the quantum boundary accurately:

selected Dirac source $\longrightarrow$ standard CAR/AQFT construction $\longrightarrow$ free local quantum net.

The CAR relations and Hadamard machinery are not inferred from modal graphs.

9.3 The interacting boundary

Current MTT work also records a classical BV master action, a gauge-fixed Green-hyperbolic equicausal algebra, and a formal all-orders anomaly-free quantum master equation at the declared perturbative tier. Those results support a conditional perturbative graph construction. A formal jet at zero coupling does not select a unique fixed-nonzero-coupling C^* -completion. A geometry-selected nonperturbative gauge-BRST bridge or regulator/continuum limit, selected physical state, RG matching, uncertainty control, and observable comparison remain open.

9.4 Finite transferred operations and the physical endpoint

The cohesive source is a twisted perfect-object benchmark with integrable superconnection $\overline{E}$. Conjugation cancels its scalar overlap twist on endomorphisms, giving the ordinary differential $d = [\overline{E}, -]$. A chosen Hermitian metric gives total-space Hodge theory; it does not give a

uniformly bounded family of fiberwise Green operators through a cohomology-rank jump [13, 7]. Its Maurer–Cartan residual $da + a^2$, together with the linear gauge row, has positive normal Δ_1 at the exact background. The resulting heat semigroup is the tangent repair evolution, not the full nonlinear flow or a signed physical action [8]. A derived equivalence preserving formal deformations does not automatically preserve that metric, adjoint, or spectrum. This is why the repair-jet graph construction still requires its own declared metric and action interpretation.

Cohesive’s finite response hierarchy supplies a related but different structure from the scalar repair jets above. Its 144-dimensional Weyl DGA retracts onto a 48-dimensional response complex. A DGA morphism preserving the inclusion, projection, and homotopy transports every transferred m_n ; the proof moves that morphism through each decorated transfer tree. The finite translation and Fourier actions preserve those source identities, so their order-36 covariance acts at all arities [7, 5]. This is not an inference from checking a few vertices. Nor are the m_n already a signed physical action or measured couplings. An approximate or nonreducing comparison needs separate defect estimates and does not inherit exact Hodge or all-arity identities.

The endpoint-factorization theorem explains what physical promotion would require [12]. Geometry and signed action (GAS), spectral synthesis and its domain/contraction certificates (SYN), and a BV-compatible four-dimensional compactification (BV4) must share one source. Their symmetry and Galerkin/Feshbach rows are then derived from that source, not additional freely selected kernels. For example, fixing a retained quadratic block does not fix the complement contribution to its Schur complement. The frozen contract accepts zero of the three complete physical packets and zero of its seven endpoint rows. These are structured source obligations, not a three-parameter count. Their openness leaves both the finite graph theorem and the selected free CAR net intact.

10 Three illustrative cases

10.1 Euclidean scalar model

For a positive matrix K , the finite Gaussian reference measure is literal. A quartic interaction defines a normalized probability measure only when its full Boltzmann weight is integrable; a nonnegative quartic form is one sufficient condition. For an arbitrary quartic tensor the Wick coefficients still make sense formally, but positivity of K alone does not make the interacting integral finite. This example shows that the graph grammar is not uniquely quantum without equating a formal series with a convergent integral.

10.2 Free fermions

For an odd field, the Gaussian integral is algebraic in Grassmann variables, and contractions carry signs. The ungraded graph shape alone cannot recover those signs. One needs the parity and ordering data. In a physical CAR theory one additionally needs the CAR algebra, localization, and a state.

10.3 Gauge fields

For gauge fields, the uncorrected Hessian has a kernel along gauge directions. A propagator appears only after gauge-fixing or homological reduction, and ghost vertices are part of the

graph datum. BRST or BV identities constrain admissible counterterms. This case makes especially clear why “inverse Hessian plus coherence” is insufficient.

11 Classical, quantum, and gravitational uses

11.1 Classical statistical fields

In classical statistical field theory, a positive covariance and interaction potential generate graph expansions of moments. This is a fully adequate explanation for the recurrence of the same combinatorial grammar. It does not imply that classical and quantum theories differ only in interpretation: noncommutative observables, interference phases, local commutation relations, and state conditions are additional mathematical structures.

11.2 Quantum field theory

In QFT, Wick expansion acts on a quasifree state or time-ordered product. The propagator prescription, statistics, local algebra, and renormalization conditions give the graphs their physical meaning. The current MTT free-CAR theorem provides such a free quantum source on its declared branch. Interacting QFT requires the remaining bridge.

11.3 Perturbative gravity

Expanding a specified gravitational action around a background also produces quadratic and higher tensors, so graph organization applies. This neither proves UV completion nor shows that spacetime quantization is unnecessary. It establishes only the same conditional perturbative grammar. Any stronger MTT gravity claim must derive the background, action, gauge complex, state, and ultraviolet control from the selected source.

12 Status and theorem ownership

Statement	Status	Owner or boundary
Finite graded Wick graph expansion	Exact here	Theorem 3.1
All-retained finite projection equality	Exact here	Theorem 4.1
Closed-sector factorization under no mixed vertices	Exact here	Theorem 4.2
Graph-combinatorics nonselection	Exact here	Theorem 5.1
Selected q79 free even-CAR net	Closed at declared tier	Companion MTT-to-QFT paper; imported CAR/AQFT theorems
Embedded renormalized-SM equivalence	Closed at profile tier	Current A04 result; not a derivation of quantization
Upper action and automorphism transfer	Open	Current blocker B.ACTION.01
Nonperturbative interacting gauge-BRST C^* bridge	Open	Current blocker B.QFT.02
No-knob Standard Model values	Open	Current strict-upgrade ledger

This ownership table prevents a common form of accidental duplication. The paper proves the finite graph and transfer statements. It cites the free-CAR result. It does not promote formal or profile evidence into an interacting-QFT theorem.

13 Validation program

A candidate upper-to-lower diagrammatic derivation should be accepted only after the following checks.

1. **Source check:** identify the action, background, grading, and all continuous inputs by path and hash.
2. **Quadratic check:** prove invertibility on the declared gauge-fixed or reduced domain and specify the inverse prescription.
3. **Vertex check:** compute every retained V_n from the same action, including overlap integrals and normalization conventions.
4. **Projection check:** verify $KP = P^*K$, the vertex intertwiners, and whether mixed vertices vanish or are integrated out.
5. **Statistics check:** preserve the $\mathbb{Z}_2$ grading and Koszul signs.
6. **Gauge check:** verify BRST/BV chain maps and Ward or QME identities.
7. **Renormalization check:** specify distribution extensions, scheme transport, and residual finite freedoms.
8. **Physical check:** establish local algebra, positivity/state conditions, uncertainty control, and observable comparison at the claimed tier.

Passing only the graph-shape check is not enough.

14 Discussion

14.1 What survives from the original idea

The central intuition survives in a stronger form. Feynman-style diagrams are not mysterious pictures added to quantum mechanics. They are the natural combinatorial language of perturbative contractions. MTT can use that universality because its fixed points and projections offer a candidate upstream organization of backgrounds and retained sectors.

The correction is equally important. “Universal graph grammar” is not “universal physical dynamics.” A graph does not know which kernel is causal, which state is positive, which field is fermionic, which gauge identity must hold, or which counterterm is allowed. Those facts reside in the typed data attached to the graph.

14.2 Why the projection theorem matters

[Theorem 4.1](#) gives a concrete target for future MTT work. If the selected upper action is found, one need not compare final amplitudes by analogy. One can verify the commuting projection at the level of the quadratic operator, every interaction tensor, grading, and identity operator. Equality of the retained graph coefficients then follows mechanically.

The theorem also identifies failure modes. If $KP \neq P^*K$, the lower propagator is not simply PGP^* . If mixed vertices survive, discarded modes generate effective interactions. If BRST/BV operators fail to intertwine, gauge identities need not descend. These are informative obstructions, not merely missing prose.

14.3 Best next theorem

The decisive upstream theorem is:

$$\begin{array}{c}
\text{selected q79 upper action and complex} \\
\Downarrow \\
K, \{V_n\}, Q_{\text{BRST}}, \text{state/renormalization data} \\
\Downarrow \quad \text{by certified intertwiners} \\
\text{accepted lower free and interacting structures.}
\end{array}$$

This would turn the graph datum from an input into a derived object. Until then, the present paper supplies the exact transfer mathematics and the boundary that any claimed derivation must cross.

15 Conclusion

The recurrence of Feynman graphs has a clean explanation. A quadratic contraction and multilinear interactions generate graph-indexed perturbative coefficients through graded Wick expansion. This grammar is shared by classical and quantum models. It is not, on its own, a quantization theorem.

We proved that compatible finite projection preserves every all-retained graph coefficient and, under absence of mixed vertices, the normalized retained perturbative expansion. We also proved that graph combinatorics cannot select numerical weights, statistics, causal prescriptions, gauge data, or renormalization. These statements give modal diagrammatics a precise role: it is an exact language for transferring a *selected* perturbative datum, not a substitute for selecting that datum.

For MTT, the free q79 even-CAR net is already available through a companion construction using standard CAR/AQFT machinery. The next frontier is the same-source upper action and its interacting gauge-BRST completion. A future success there can plug directly into the theorem proved here.

Computational Evidence and Reproducibility

The theorems proved here are finite algebraic statements and do not depend on numerical fitting. The curated repository <https://github.com/PeterNero/mtt-results-repro> is cited for current MTT status context. In particular, A04/final_12_of_12_audit records embedded renormalized-Standard-Model equivalence at the adopted profile tier, while A05/strict_upgrade_ledger records stronger no-knob obligations that remain open. Neither row proves [theorems 3.1, 4.1](#) and [5.1](#), and the open row is not evidence of closure.

The selected free-CAR theorem and the classical-to-quantum interface are documented in the companion records [\[10, 9\]](#). Exact release artifacts, source hashes, and the revision audit accompany this paper in the MTT papers repository.

References

- [1] Romeo Brunetti and Klaus Fredenhagen. Microlocal analysis and interacting quantum field theories: Renormalization on physical backgrounds. *Communications in Mathematical Physics*, 208:623–661, 2000.

- [2] H. Epstein and V. Glaser. The role of locality in perturbation theory. *Annales de l'Institut Henri Poincaré, Section A*, 19(3):211–295, 1973.
- [3] Stefan Hollands and Robert M. Wald. Local wick polynomials and time ordered products of quantum fields in curved spacetime. *Communications in Mathematical Physics*, 223:289–326, 2001.
- [4] Stefan Hollands and Robert M. Wald. Existence of local covariant time ordered products of quantum fields in curved spacetime. *Communications in Mathematical Physics*, 231:309–345, 2002.
- [5] Peter Nero. All-arity contraction-morphism source promotion. https://github.com/PeterNero/mtt-results-repro/blob/f141a20ea23c5c3ff19cc2161c0e226e29ade8a7/release/results/causal_base_q79_all_arity_source_promotion_packet/artifact.json, 2026. Frozen source record; result causal_base_q79_all_arity_source_promotion_packet. Scope as stated in the text.
- [6] Peter Nero. Closure geometry and a regime-local ten-dimensional action ansatz, 2026. Version 5, sections on variational anchors and cyclic action; MTT papers repository.
- [7] Peter Nero. Cohesive Closure Repair and Its Hodge, Kernel, and Projection Shadows: From Nonlinear Defects to Tangent Semigroups, with the Physical-Action Boundary, 2026. Version 14 structural identities; contextual owner of the finite transfer and endpoint contracts.
- [8] Peter Nero. Cohesive Maurer–Cartan repair and conditional transform. https://github.com/PeterNero/mtt-results-repro/blob/f141a20ea23c5c3ff19cc2161c0e226e29ade8a7/release/results/q79_maurer_cartan_repair/artifact.json, 2026. Frozen source record; scope as stated in the text.
- [9] Peter Nero. From coherence-basin statistics to a local quantum-field interface: Commutative measures, the selected free car net, and the interacting boundary, 2026.
- [10] Peter Nero. Modal triplet theory and quantum field theory on curved spacetime: A selected free car net and the interacting reconstruction boundary, 2026.
- [11] Peter Nero. Preprojection repair-jet to perturbative-graph functor, H4-T8. Frozen finite/formal result, https://github.com/PeterNero/mtt-results-repro/blob/f141a20ea23c5c3ff19cc2161c0e226e29ade8a7/release/results/preprojection_repair_jet_graph_functor/artifact.json, 2026.
- [12] Peter Nero. Seven-row endpoint factorization through three source packets. https://github.com/PeterNero/mtt-results-repro/blob/f141a20ea23c5c3ff19cc2161c0e226e29ade8a7/release/results/causal_base_q79_seven_row_endpoint_factorization_packet/artifact.json, 2026. Frozen source record; result causal_base_q79_seven_row_endpoint_factorization_packet. Scope as stated in the text.
- [13] Peter Nero. Twisted cohesive superconnection and stratified Hodge package. <https://github.com/PeterNero/mtt-results-repro/blob/>

[f141a20ea23c5c3ff19cc2161c0e226e29ade8a7/release/results/q79_cohesive_superconnection/artifact.json](#), 2026. Frozen source record; scope as stated in the text.

- [14] Kasia Rejzner. *Perturbative Algebraic Quantum Field Theory: An Introduction for Mathematicians*. Springer, Cham, 2016.
- [15] G. C. Wick. The evaluation of the collision matrix. *Physical Review*, 80:268–272, 1950.