

Modal Triplet Theory: Perturbative Coherent-Sector Quantum Gravity and the Heterotic UV-Completion Boundary

Peter Nero

Version 7
September 2026

Abstract

This paper separates three levels of the current Modal Triplet Theory (MTT) quantum-gravity program. First, an exact finite-carrier result supplies internal transverse-traceless support, while a separate controlled reduction recovers the four-dimensional Einstein action only under declared higher-dimensional action, truncation, gap, and normalization hypotheses. Second, once the resulting local Einstein–matter effective action, Wilson coefficients, state, gauge fixing, and scheme are supplied, standard background-field and BV/BRST methods give fixed-order perturbative quantum-GR equivalence. This is an imported parity construction, not a derivation of the quantum measure or Wilson coefficients from MTT. The two-derivative theory is not ultraviolet complete: its counterterm order grows with loop number and the two-loop pure-gravity divergence is nonzero. Third, the selected compatible ultraviolet route is the q79 Fu–Yau heterotic branch. A conditional inheritance theorem gives absence of local ultraviolet divergences at each fixed genus and multiplicity if an exact modular, anomaly-free heterotic worldsheet theory with factorization, GSO projection, and BV vertices is supplied. The current q79 worldsheet contract has five of twelve rows available, two partial, and five open. Hidden projective holomorphic existence and existential HYM are established at their declared scope. The selected visible bundle, explicit common chamber and connections, complete Green–Schwarz representative, exact worldsheet SCFT, string-field realization, infrared completion, all-genus control, and nonperturbative definition remain open. The resulting claim is therefore low-energy quantum-gravity parity plus a sharply specified heterotic completion program, not completed ultraviolet quantum gravity.

Version 7 revision note

Supersedes. Version 6, with its released metadata and earlier note retained.

Reason. The worldsheet boundary needed the intervening spectral, Bott–Chern, current-algebra, and finite-field classifications, rather than blanket openings inherited from an older ledger.

Resolution. Explain all six frozen assigned records in their physical context, including the full Bott–Chern successor and established hidden HYM existence. Keep physical spectral support distinct from auxiliary covers.

Retained result. Finite TT support, fixed-order EFT parity, the selected heterotic route, and conditional fixed-genus inheritance retain their tiers.

Open boundary. The same-source visible endpoint, explicit common HYM chamber, positive balanced realization, full worldsheet and BV data, and all-genus or nonperturbative completion remain open.

Version 6 revision note

Supersedes. *Modal Triplet Theory: Perturbative Coherent-Sector Quantum Gravity and the Heterotic UV-Completion Boundary*, version 5.

Reason. Version 5 correctly discovered the Gaussian positivity no-go and loop-rank obstruction, but it repeated those results in several appendices, retained an unselected schedule as a derivation of the SPT endpoint, and sometimes described imported perturbative QFT machinery as though it were selected by MTT.

Resolution. This edition assigns the SPT mathematics to the companion Euclidean TT paper, states the low-energy result at its exact fixed-order parity tier, presents all twelve q79 worldsheet gates, and separates fixed-genus ultraviolet inheritance from all-genus and nonperturbative completion.

Retained result. The current chain supports an exact internal TT carrier, conditional classical GR reduction, fixed-order quantum-GR EFT parity, selection of the q79 heterotic route as the primary compatible UV candidate, and a conditional fixed-genus inheritance theorem.

Remaining boundary. MTT has not yet selected the physical higher-derivative Wilson values, complete Lorentzian quantum state and measure, physical nonpullback heterotic bundles, exact q79 worldsheet theory, or nonperturbative ultraviolet completion.

Contents

Version 6 revision note	2
1 Purpose, Orientation, and Claim Tiers	3
2 The Low-Energy Coherent-Sector Tier	3
2.1 What MTT supplies before quantization	3
2.2 Fixed-order quantum-GR parity	4
2.3 Why fixed-order predictivity is not UV completion	5
2.4 Finite transfer and the interacting source dependency	5
3 Why SPT Does Not Close the Ultraviolet Problem	6
4 The Selected q79 Heterotic Route	7
4.1 Why this is the primary compatible candidate	7
4.2 The physical spectral support and line problem	7
4.3 Poincare descent acts on the spectral object	8
4.4 Bott–Chern exactness removes a source condition, not positivity	8
4.5 The left-current completion is not an aggregate monad split	8
4.6 Geometric consumer results and their remaining targets	9
4.7 Conditional fixed-genus inheritance	9
4.8 The twelve-row worldsheet contract	10
4.9 What the finite-field mirror classification actually excludes	11
5 Branch Orientation, Conjugation, and Double Traversal	11

6	What Has Been Achieved and What Would Close the Program	11
6.1	Achievements at their declared tiers	11
6.2	The shortest honest closure sequence	12
7	Conclusion	12

1 Purpose, Orientation, and Claim Tiers

In plain language, the present construction has a low-energy bridge and a candidate high-energy bridge. The low-energy bridge says that if MTT supplies the same effective gravitational data used by quantum GR, then the standard quantization machinery yields the same fixed-order observables. The high-energy bridge says that if the selected q79 geometry can be completed into an exact heterotic worldsheet theory, it inherits the string worldsheet’s fixed-genus ultraviolet behavior. Neither conditional bridge manufactures its missing input.

The logical tiers are reconciled with the frozen mixed QG ledger [1]. Its old paper-disposition and open-payload fields are historical checkpoints, not instructions to reopen later scoped results. The logical tiers are:

Tier	Current status	Meaning
Finite internal TT carrier	exact on the declared finite branch	Internal helicity-two support, not yet a normalized Lorentzian graviton
Four-dimensional Einstein reduction	conditional	Requires an higher-dimensional Einstein sector, controlled truncation, and stabilized non-Einstein modes
Interacting quantum-GR EFT	fixed-order parity	Standard QFT structure is imported after action, Wilson data, state, and scheme are declared
SPT Gaussian filter	conditional Euclidean model	Useful graph regulator; not selected physical propagation
q79 heterotic route	selected compatible route	The route is selected, but its physical worldsheet object is incomplete
Fixed-genus string UV inheritance	conditional theorem	Applies only after the exact worldsheet hypotheses are met
All-genus or nonperturbative QG	open	No convergence or constructive definition has been supplied

The word *parity* is important. It means equality to an already defined target theory after matching all target inputs at a declared order. It is a valid equivalence statement, but it is not a no-input prediction or a derivation of those inputs.

2 The Low-Energy Coherent-Sector Tier

2.1 What MTT supplies before quantization

The finite proto-spinor/GR program proves an exact internal TT support identity on the selected finite carrier. It identifies the intended helicity-two internal block and removes an

ambiguity that was formerly left as an ansatz. It does not by itself provide:

- a Lorentzian principal symbol and physical massless pole;
- the Newton or Planck normalization;
- universal conserved coupling to the full stress tensor;
- the causal state or asymptotic prescription; or
- the interacting higher-derivative Wilson coefficients.

The companion paper *Controlled Coherent Reduction to Four-Dimensional Einstein Gravity* supplies a separate analytic bridge. On a $4 + 6$ -dimensional geometry, it gives a contraction/Schur–Feshbach estimate for eliminating heavy modes, provided that the higher-dimensional Einstein sector, gap, small coupling, truncation, and zero-mode audit are already available. The shared circle is counted once as common holonomy data and is not identified with compact physical time.

The finite source is stronger than a mode count. The frozen ledger also records the normalized S_3 Reynolds mismatch Hessian: on the finite diagonal/edge strain blocks, $h_{DE} = 0$ and $h_{DD} = h_{EE} = \kappa_e > 0$. This exact finite operator does not wait for a continuum Fourier–Mukai/HYM realization. Conversely, the flat root-stack sheet-symbol connection cannot equal an entire nonzero-Chern physical HYM connection. A covariant family of quarter-turn related branches is not by itself an invariant one-branch Hessian. These distinctions preserve the finite theorem without promoting it to a continuum naturality theorem [1].

At the two-derivative source tier the ledger permits an explicit discrete physical-realization premise A_{QG} . With that premise and the declared coframe/metric-descent hypotheses, the torsion construction gives the TEGR/Einstein action up to normalization, cosmological term, and boundary terms. The premise is not derived by uniqueness of an equilibrium inside a chosen basin. Neither a quadratic TT Hessian nor the exact TEGR boundary identity selects all nonlinear physical action data. The selected causal representative additionally declares its binary causal boundary mark A_{causal} ; it does not identify the internal shared phase circle with time. This is the conditional branch/coframe interface, not selection of Newton normalization, the vacuum state, or ultraviolet completion [1].

2.2 Fixed-order quantum-GR parity

Let $S_{\text{EFT}}^{(N)}$ denote the local diffeomorphism-invariant Einstein–matter effective action through a declared loop and derivative order N . It includes the Einstein term, cosmological term, matter sector, and the finite list of local higher-curvature operators required at that order. Let σ denote the causal state, gauge fixing, scale, and renormalization scheme. The current parity construction is summarized by

$$\mathcal{O}_{\text{MTT}}^{(N)} = \text{Readout} \circ \text{Green}_\sigma \circ Q_{\text{GR,EFT}}^{(N)} \circ E_{q79}. \quad (2.1)$$

Here E_{q79} embeds the declared coherent source data into the effective field variables, $Q_{\text{GR,EFT}}^{(N)}$ is the standard fixed-order background-field/BV quantization, and the final maps construct the selected observables.

Proposition 2.1 (Fixed-order parity statement). *If E_{q79} supplies the same renormalized action $S_{\text{EFT}}^{(N)}$, Wilson coefficients, gauge fixing, state, scale, and scheme as quantum GR EFT through order N , then the renormalized generating functionals and gauge-invariant observables obtained from (2.1) agree through order N .*

Proof. After the stated data are identified, both constructions apply the same renormalized functional and the same observable maps. Equality follows by composition. The proposition proves a typed embedding/equivalence result; it does not select the matched data. $\square$

What this means. The proposition checks that the MTT carrier can host the standard fixed-order quantum-GR calculation without changing its predictions. Its scientific content is compatibility and typed transport. Predictive improvement would require MTT to emit some of the matched Wilson values, state data, or normalization before the target calculation is imported.

On a globally hyperbolic background with a Hadamard state, local covariant time-ordered products and perturbative observables can be constructed using Epstein–Glaser or perturbative algebraic QFT methods [2, 3, 4, 5]. The BV induction restores the quantum master equation order by order when the relevant local ghost-number-one anomaly class vanishes [6, 7, 8]. These are standard imported theorems. The current MTT source does not yet derive the full measure, Hadamard state, anomaly cohomology, or fixed-nonzero-coupling completion.

2.3 Why fixed-order predictivity is not UV completion

For a connected Einstein graph with L loops, V two-derivative vertices, and I internal lines, the superficial derivative order is

$$D = 4L + 2V - 2I.$$

Using $L = I - V + 1$ gives

$$D = 2L + 2. \tag{2.2}$$

For example, tree, one-loop, and two-loop graphs require operators of superficial derivative order two, four, and six respectively. The Riemann-cubed counterterm at two loops is therefore the first explicit worked case showing why the two-derivative parameter set cannot remain closed. At every fixed loop order only finitely many local counterterms are needed, which is precisely the effective-field-theory notion of predictivity [9]. Across all loop orders, however, the derivative order is unbounded. Pure gravity is on-shell finite at one loop for zero cosmological constant in the standard sense, but it has a nonzero two-loop Riemann-cubed divergence [10, 11].

Consequently, the two parameters of the two-derivative action do not define an all-scale interacting quantum theory. The current MTT parity theorem allows a finite set of declared Wilson values at any chosen order; it does not emit their complete tower.

2.4 Finite transfer and the interacting source dependency

The cohesive benchmark already carries an integrable twisted superconnection whose scalar overlap twist cancels in the endomorphism dg algebra. A chosen Hermitian metric supplies total-space Hodge theory, but not a uniformly bounded fiberwise Green family across rank jumps [12]. Its gauge-fixed Maurer–Cartan squared residual has Hodge tangent normal and tangent heat evolution [13]. These are exact structural statements, not a physical metric choice or Lorentzian signed action; a derived equivalence alone does not transport adjoints or spectra.

The all-arity transfer theorem owned by Cohesive explains a structural simplification relevant to perturbative vertices. A source DGA map preserving the differential, product,

inclusion, projection, and homotopy transports every transferred operation. The selected 144-to-48 finite contraction has this property under its order-36 covariance action [14, 15]. This is exact finite algebra and a general conditional transport theorem, not a continuum HYM action or an all-orders quantum-gravity amplitude construction. Exact Hodge transport further needs a unitary reducing map on the declared closed domains; cost pullback alone does not supply it.

The CBF.T12 endpoint contract groups seven physical rows into geometry and signed action (GAS), spectral synthesis (SYN), and BV-compatible four-dimensional compactification (BV4) [14, 16]. Symmetry transport and the Galerkin/Feshbach calculation follow from compatible GAS and SYN data. The count is three informational source types, not three scalars; no complete physical packet is accepted in the frozen record. Cotangent repair and a positive normal do not replace the field-only action, its real slice, or its charged compactification.

The companion quantum-mechanics source account owns a complementary family dependency [17]. A finite projection cannot recover discarded smooth connection directions. Given the selected physical HYM/action source, common domains, and ellipticity, however, the internal Dirac/Hessian family, gapped-chart Riesz projectors, Kato transport, and determinant line follow from standard analytic constructions. The determinant line is not a full chiral-measure current; crossing and sector gluing and uniform fixed-coupling estimates remain separate. Hidden existential HYM and the operational free CAR/QM results stay closed, while the selected common physical source and nonperturbative interacting theory remain open.

3 Why SPT Does Not Close the Ultraviolet Problem

The companion paper *An SPT-Filtered Euclidean TT Model and Its Conditional Perturbative Properties* owns the SPT theorems. Its conclusions can be summarized without repeating their proofs:

1. an internal spectral gap controls internal inversion but does not produce an external Gaussian multiplier;
2. a positive proper-time endpoint is an additional filter or schedule assumption;
3. Gaussian-damped lines control all L loop directions exactly when their loop-momentum matrix C_Γ has rank $C_\Gamma = L$, equivalently when unfiltered edges form a forest; and
4. a nonzero positive Kallen–Lehmann/Stieltjes propagator cannot have permanent Gaussian decay in the same Euclidean spectral variable.

These results leave SPT useful as a Euclidean regulator or exploratory effective filter. They withdraw the former route

internal gap $\implies$ physical Gaussian propagator $\implies$ OS positivity, causality, and unitarity.

Each arrow needs an independent theorem, and the first arrow is false in general. The exact finite TT eigenvalue recorded by the internal carrier is dimensionless until a physical units-and-normalization map is supplied; it is not automatically the Planck cutoff.

4 The Selected q79 Heterotic Route

4.1 Why this is the primary compatible candidate

The q79 program selects a Fu–Yau-type heterotic route rather than the SPT filter as its primary compatible ultraviolet candidate. Fu–Yau geometry provides established existence machinery for non-Kähler heterotic compactifications with torsion under specific bundle, stability, and anomaly conditions [18]. Torsion linear sigma models provide a related worldsheet framework [19]. These external theorems do not say that every topological MTT packet is automatically a physical heterotic vacuum.

The current q79 records include:

- a time-oriented discrete branch label $q = 79$, with conjugate 369 modulo 448;
- a degree-two K3 incidence GLSM base and rank-one Fu–Yau source data;
- curvature-level Green–Schwarz/Bianchi information;
- finite gerbe, torsion-phase, and modular-orbit data;
- a smooth topological visible $SU(3)$ candidate with $c_2 = 9u$ and $c_3 = \pm 6$; and
- partial twisted-spectral and modular-character calculations.

Topological existence and Hodge admissibility do not yet produce the physical nonpullback holomorphic bundle, a common positive HYM chamber, connections, or the complete local worldsheet anomaly data. Those are the physical endpoint obligations, not optional polish.

4.2 The physical spectral support and line problem

The visible spectral cutset is an exact reduction of a physical endpoint problem, not a completed endpoint [20]. Let H be the degree-two K3 polarization, D_0 the zero section of the dual elliptic factor, and u, Ω the declared degree-four and degree-six source classes. The target $\text{ch}(V_3) = 3 - 9u + 3\Omega$ has rank-one spectral support

$$C_{\text{phys}} \in |9H + 3D_0|.$$

The older $H + 3D_0$ cover cannot realize that character: changing its line does not change the codimension-one Chern row. The later auxiliary pencil $3H + 3D_0$ is likewise not this physical support. Its useful local calculations cannot be imported as physical value rows merely by matching the degree-three covering rank.

Smooth finite-flat degree-three members exist on a nonempty open subset. If a particular member carries the required twisted locally free line whose twist cancels the inverse Poincaré kernel, the inverse transform is a finite-flat pushforward of an ordinary line. Rank-three local freeness and vanishing higher direct images then follow; they need not be imposed again. The actual line and admissible member, however, have not been selected by this implication.

The determinant requirement is a twisted norm equation $\text{Nm}_\alpha(L) = \mathcal{O}(27H)$, since $\det p_* \mathcal{O}_C = \mathcal{O}(-27H)$. The integral baseline $\ell_0 = 2D_0|_C + 3p^*H$ has that norm. Writing $\ell = \ell_0 + q$, the remaining Grothendieck–Riemann–Roch conditions are the trace-free Prym requirements

$$\text{Nm}(q) = \mathcal{O}, \quad j_* q = 12H^2 - 9\gamma x, \quad j_*(q^2) = -66H^2 D_0.$$

Here $j : C \hookrightarrow K3 \times E^*$ is the inclusion, while γx denotes the mixed source class in the frozen transform’s convention. These equations specify the required line data; they do not construct it. The inherited ambient line alone would require fractional coefficients to yield

$c_3 = 6$, so that restricted ansatz is excluded. It does not exclude all Picard/Prym or twisted constructions.

No global rank exclusion is inferred from a fixed-fiber Picard point. The retained CBF T69 correction withdraws that inference [21]; T70–T73 establish local coefficient observability, not the global Picard/BHT derivative. The local/global distinction remains part of the physical line problem.

4.3 Poincare descent acts on the spectral object

The normalized Poincare kernel $\mathcal{P}$ obeys

$$(\text{id} \times t_g)^* \mathcal{P} \simeq \mathcal{P} \otimes p^* L_g, \quad t_g^* \Psi \simeq \Psi(- \otimes L_g).$$

For nonzero $g \in E[3]$, the double-dual degree-zero line L_g is nontrivial. The bare kernel therefore has no scalar-projective translation linearization. The exact categorical intertwiner instead puts the qutrit multiplier on isomorphisms $\phi_g : F \otimes L_g \rightarrow F$ of the spectral object [22]. With $\sigma((a, b), (c, d)) = \omega^{-cb}$, $\omega^3 = 1$, the finite cocycle has the required associativity. A length-three fiber object transforms to rank three at degree zero, and three such summands to rank nine. These are fiberwise categorical constructions, not a new selection of the global visible line or of physical connection coefficients.

4.4 Bott–Chern exactness removes a source condition, not positivity

The invariant computation has a 21-dimensional $(2, 2)$ space, a 20-dimensional closed subspace, and a one-dimensional $\partial\bar{\partial}$ image. Its 19-dimensional Bott–Chern quotient injects into de Rham cohomology [23]. Thus a total invariant de Rham-exact Chern–Weil difference is already $\partial\bar{\partial}$ -exact; separate exact formulas for the visible and hidden quadratic representatives are not additional Aepli premises.

The later full Fourier-homotopy result strengthens this statement [24]. On the holomorphic elliptic action, a nonzero character (m, n) has contraction eigenvalue product $-(m^2 + n^2)/4$. Its reciprocal gives a smooth summable homotopy, so every nonzero Fourier mode of a Bott–Chern cocycle is exact. Averaging handles the invariant zero mode. Consequently the full $(2, 2)$ map is injective, and gauge-connection equivariance is no longer a separate premise for this anomaly-potential conclusion.

This supplies a potential once the actual holomorphic source pair exists and its total topological class cancels. It does not prove that the resulting Hermitian form is positive and conformally balanced, or select the integral order-three string refinement. The hidden projective holomorphic object and existential HYM connection are already established [25]: a finite subobject bound M gives a stable-factor chamber for $t > 1 + 3M$, with the qutrit orbit sharing the bound. The numerical value of M , an explicit common visible–hidden chamber, and the connection arrays remain different requirements. Hidden existence is not reopened by the older frozen invariant packet.

4.5 The left-current completion is not an aggregate monad split

The current-algebra construction uses the ordered local visible/hidden rank-three/rank-nine carriers in different E_8 factors [26]. Their complementary projectors are $P_3 = \text{diag}(I_3, 0_9)$ and $P_9 = \text{diag}(0_3, I_9)$. They are not extracted from the aggregate rank-twelve Picard monad: that monad supplies total local anomaly and $c_2 = 20$ data, but its parity restriction excludes the physical odd $9 + 11$ split.

The left-current complement has an explicit minimal presentation by four neutral complex left-moving Fermis with $E = J = 0$ and unit kinetic metric. It contributes $\Delta c_L = 4$, $\Delta c_R = 0$, and the declared left-current and gravitational anomaly shifts -4 . Permuting their four optional q79 line labels changes only the fermionization basis: the kinetic action and the selected lattice have the same permutation symmetry. There is no extra fitted line assignment. Bosonization replaces the same Fermi determinants; it does not add a second copy of them. This closes the smooth-projective free left-current construction, not the interacting holomorphic E/J source, right-moving spin/GSO data, or an exact infrared $(0, 2)$ SCFT.

4.6 Geometric consumer results and their remaining targets

The global Fitting result owned by Cohesive concerns an auxiliary twisted Hartshorne–Serre derived object [14, 27]. Its local module isomorphisms and invertible transition lines preserve minor ideals, so the Fitting ideals descend as ordinary ideal sheaves even before literal overlap matrices have been executed. The local determinant $(rt)^3$ matches the global class $3H + 3D_0$. This does not trivialize the gerbe or identify that auxiliary class with the physical $9H + 3D_0$ support. Literal chain overlaps, a common physical HYM realization, and the continuum-to-finite intertwiner are stronger data. The later hidden existential HYM theorem is retained rather than overwritten by the older packet’s status field.

The same owner’s large-gauge result fixes the kernel of the selected b_{K3} map, not the number of physical source coordinates [28]. Its kernel is zero and its source remains $\mathbb{R}^2$, immersed injectively with dense nonclosed image in the compact transcendental T^{20} . Period normalization and kinetic scales are not supplied by rank zero, and the shared-circle action intertwiner is a separate target.

The eta9 execution results, owned by the flux/compactification account, have another scope. Gate 1 completes all thirty original-Jacobian groups and 225 selected support columns, notwithstanding its older readiness header [29]. It does not compute Gate 2, a marked integral meridian, the 248-coordinate period readout, or β_C . The selected transported three-cycle subsystem has Gram matrix $-2I_3$, determinant -8 , and rank three. B96 makes every affine coefficient vector on that subsystem a rational coboundary, so this particular detector is exhausted [30]. It is not an exclusion of the global affine class or of a larger meridian subsystem. These exact local/finite conclusions do not need to be restarted, but they do not supply the physical Deligne/HYM endpoint or ultraviolet quantum gravity.

Two later flux-owned results sharpen this execution boundary. The complete same-source local connection chart exists on $0 \leq u \leq 2^{-96}$, where u is a dimensionless marked connector, not physical time [31]. Positive width is an achieved local result; useful-width continuation, the global frame and path, integral readout, and β_C are not thereby computed. Separately, B89 is rejected by its certified same-source affine Deligne obstruction [32]: a mod-two covector annihilates $M - I$ but pairs to one with the affine return, obstructing an integral preimage. This decides that candidate, not a replacement endpoint, and does not establish exact order two of the integral class. Neither result is a reason to restart the closed local computations or to promote the physical HYM/UV frontier.

4.7 Conditional fixed-genus inheritance

Theorem 4.1 (Heterotic fixed-genus UV inheritance). *Assume the selected q79 background is completed to an exact anomaly-free modular heterotic $(0, 2)$ worldsheet theory with a tachyon-free GSO projection, correct factorization, and quantum-BV string vertices. Then*

each fixed-genus, fixed-multiplicity on-shell amplitude has no local point-particle ultraviolet divergence. Possible degeneration-boundary singularities are governed by physical factorization channels and require the separate tadpole, vacuum-shift, and infrared prescriptions.

Reason for the inheritance. Worldsheet moduli are integrated over a modular fundamental domain rather than over an unrestricted point-particle proper-time region. Short-distance operator collisions are controlled by the worldsheet OPE and BRST factorization, while modular equivalence removes duplicate ultraviolet regions. Degenerations of the Riemann surface represent long tubes and physical on-shell propagation, so their divergences are infrared or tadpole questions rather than local ultraviolet counterterms. This is the standard perturbative string framework applied conditionally to the q79 background [33, 34, 35]. $\square$

The theorem is coefficientwise in genus and multiplicity. It does not prove convergence of the genus expansion, existence of a preferred vacuum, or a nonperturbative definition.

4.8 The twelve-row worldsheet contract

The present contract is not a vague list of future work. It has twelve named rows, with five available, two partial, and five open:

Row	Required object	Current status
W1	Time-oriented q79 target branch	Available
W2	Fu–Yau Mukai charge and Green–Schwarz/Bianchi sector	Available
W3	Visible curvature-level Green–Schwarz cancellation	Available at curvature tier
W4	Critical heterotic central-charge balance	Available universally
W5	q79 low-energy GR and quantum-EFT limit	Available at declared parity tier
W6	Global Deligne gerbe and Freed–Witten condition on the full visible cycle set	Open; finite representative only
W7	All-orders-in- α' q79 target background	Open; current background is first-order
W8	Exact q79 heterotic $(0, 2)$ SCFT or all beta functions	Partial; base GLSM, local anomaly, and topological candidate exist, but the physical bundle and IR SCFT do not
W9	q79 modular-invariant GSO partition function and factorization	Partial; finite torsion module and seven-seed induction exist, analytic characters and GSO remain open
W10	q79-specific string-field vertices and BV master action	Open; standard framework exists but is not instantiated on q79
W11	Tadpole, vacuum-shift, infrared, and soft-state completion	Open
W12	All-genus convergence or a nonperturbative definition	Open

Rows W8 and W9 are genuine advances, but partial rows do not count as available rows. The current authoritative total remains 5/12, with two additional partials. Standard heterotic string-field BV machinery is known [36]; importing its existence is not the same as building the q79 vertices.

4.9 What the finite-field mirror classification actually excludes

The D10/D6 mirror-zero-zero packet is a separate algebraic source screen over $\mathbb{F}_{101}$ [37]. Each of two parent charts (indices 5 and 6) has 35 disjoint top-degree strata, indexed by $0 \leq h \leq 6$ and $0 \leq y \leq 4$. Each chart has 29 exact unit ideals and six open strata. In both charts the open degree pairs are

$$(4, 4), (5, 2), (5, 3), (5, 4), (6, 3), (6, 4).$$

Thus 58 of 70 strata are excluded exactly and 12 remain open; neither parent chart is excluded. A unit ideal means that the equations have no common zero over the algebraic closure of this finite field in the specified stratum.

The proof-direction rule matters. If a larger tail-support locus is empty, its source sublocus is empty. A positive-dimensional tail superset says nothing about existence in the source sublocus. Recursive covers require every branch, and partial Laurent calculations require the complete isomorphic quotient, before a source unit can be concluded. Timeouts and missing outputs give no mathematical exclusion. The totals are a hash-bound classification, not a proof of all physical mirror constraints, a characteristic-zero no-go, or worldsheet completion. Reading the row ledger and checking its counts is also not independent replay of its underlying Groebner certificates.

5 Branch Orientation, Conjugation, and Double Traversal

The q79 and 369 labels form a conjugate pair modulo 448:

$$79 + 369 \equiv 0 \pmod{448}.$$

Current MTT results select the retarded $q79/F/m1$ representative at the orientation level. They do not prove uniqueness of the global carrier measure, derive the initial state, or require the conjugate branch to be well-behaved.

A Schwinger–Keldysh or closed-time-path description doubles fields along forward and backward contour legs. Given a normalized density operator and unitary evolution, equal sources obey $Z[J, J] = 1$ [38, 39]. This identity expresses cancellation on a closed contour; it does not select the arrow of time or prove that a literal double traversal of the MTT circle returns the universe to a state with no space, time, or gravity.

The shared MTT circle is therefore treated as common phase or holonomy data. Physical time is a noncompact Lorentzian ordering variable, possibly related to a lift of phase data only after an additional theorem.

6 What Has Been Achieved and What Would Close the Program

6.1 Achievements at their declared tiers

- The internal TT sector is selected exactly on the finite carrier.

- A controlled analytic reduction to four-dimensional Einstein gravity is available under explicit hypotheses.
- Quantum GR EFT parity is closed at each fixed declared order after matching the action, Wilson data, state, gauge fixing, scale, and scheme.
- The finite internal carrier and the old SPT shortcut are excluded as automatic all-scale ultraviolet regulators.
- The q79 Fu–Yau heterotic route is selected as the primary compatible UV route.
- Fixed-genus UV inheritance is proved conditionally on the exact worldsheet contract.
- The worldsheet frontier is counted explicitly as five available, two partial, and five open rows.

6.2 The shortest honest closure sequence

The remaining work has a dependency order:

1. **Physical Hull–Strominger endpoints.** Retain the established hidden projective holomorphic object and existential HYM theorem; construct the selected visible bundle, an explicit common HYM chamber and connections, and the positive balanced/integral Bianchi realization.
2. **Complete W6–W9.** Promote the global gerbe, all-order background, exact $(0,2)$ SCFT, analytic modular characters, GSO projection, and factorization on that same source.
3. **Instantiate W10–W11.** Build q79-specific string-field/BV vertices and the tadpole, vacuum-shift, infrared, and soft-state completion.
4. **Connect low and high energy.** Derive the four-dimensional Wilson coefficients, Newton normalization, universal stress response, and selected state from the same compactification.
5. **Nonperturbative exit.** Establish all-genus control or another well-defined non-perturbative completion, with a positive physical observable sector and comparison map.

The first item controls the rest: without the physical bundles, worldsheet rows cannot all refer to one selected background. The final item cannot be inferred merely by adding more fixed-genus calculations.

7 Conclusion

The current MTT quantum-gravity result is substantial but tiered. It supplies an exact internal TT carrier, a conditional Einstein reduction, and fixed-order quantum-GR EFT parity. It also identifies the q79 Fu–Yau heterotic construction as the primary compatible ultraviolet route and proves the correct conditional fixed-genus inheritance statement.

It does not yet supply a UV-complete theory. The former Gaussian shortcut is a conditional Euclidean model, the low-energy quantum theory imports standard EFT quantization after matching its inputs, and the q79 worldsheet contract remains 5/12 with two partial rows. The path forward is therefore not to rename these tiers as closure. It is to construct one physical visible–hidden Hull–Strominger source, complete its worldsheet and BV contracts, derive the low-energy normalization and Wilson data from that same source, and then solve the all-genus or nonperturbative problem.

References

- [1] Peter Nero. QG source status and proof interfaces. https://github.com/PeterNero/mtt-results-repro/blob/f141a20ea23c5c3ff19cc2161c0e226e29ade8a7/release/results/qg_current_status/artifact.json, 2026. Frozen source record; result qg_current_status. Scope as stated in the text.
- [2] R. Brunetti and K. Fredenhagen. Microlocal analysis and interacting quantum field theories: Renormalization on curved spacetimes. *Communications in Mathematical Physics*, 208:623–661, 2000.
- [3] Stefan Hollands and Robert M. Wald. Local wick polynomials and time ordered products of quantum fields in curved spacetime. *Communications in Mathematical Physics*, 223:289–326, 2001.
- [4] Stefan Hollands and Robert M. Wald. Existence of local covariant time ordered products of quantum fields in curved spacetime. *Communications in Mathematical Physics*, 231:309–345, 2002.
- [5] Kasia Rejzner. *Perturbative Algebraic Quantum Field Theory: An Introduction for Mathematicians*. Mathematical Physics Studies. Springer, 2016.
- [6] Igor A. Batalin and Grigori A. Vilkovisky. Gauge algebra and quantization. *Physics Letters B*, 102(1):27–31, 1981.
- [7] Igor A. Batalin and Grigori A. Vilkovisky. Quantization of gauge theories with linearly dependent generators. *Physical Review D*, 28:2567–2582, 1983.
- [8] Glenn Barnich, Friedemann Brandt, and Marc Henneaux. Local brst cohomology in gauge theories. *Physics Reports*, 338(5):439–569, 2000.
- [9] John F. Donoghue. General relativity as an effective field theory: The leading quantum corrections. *Physical Review D*, 50:3874–3888, 1994.
- [10] Gerard 't Hooft and Martinus Veltman. One-loop divergencies in the theory of gravitation. *Annales de l'Institut Henri Poincaré A*, 20:69–94, 1974.
- [11] Marc H. Goroff and Augusto Sagnotti. Quantum gravity at two loops. *Physics Letters B*, 160:81–86, 1985.
- [12] Peter Nero. Twisted cohesive superconnection and stratified Hodge package. https://github.com/PeterNero/mtt-results-repro/blob/f141a20ea23c5c3ff19cc2161c0e226e29ade8a7/release/results/q79_cohesive_superconnection/artifact.json, 2026. Frozen source record; scope as stated in the text.
- [13] Peter Nero. Cohesive Maurer–Cartan repair and conditional transform. https://github.com/PeterNero/mtt-results-repro/blob/f141a20ea23c5c3ff19cc2161c0e226e29ade8a7/release/results/q79_maurer_cartan_repair/artifact.json, 2026. Frozen source record; scope as stated in the text.

- [14] Peter Nero. Cohesive Closure Repair and Its Hodge, Kernel, and Projection Shadows: From Nonlinear Defects to Tangent Semigroups, with the Physical-Action Boundary, 2026. Version 14 structural identities; contextual owner of the finite transfer and endpoint contracts.
- [15] Peter Nero. All-arity contraction-morphism source promotion. https://github.com/PeterNero/mtt-results-repro/blob/f141a20ea23c5c3ff19cc2161c0e226e29ade8a7/release/results/causal_base_q79_all_arity_source_promotion_packet/artifact.json, 2026. Frozen source record; result causal_base_q79_all_arity_source_promotion_packet. Scope as stated in the text.
- [16] Peter Nero. Seven-row endpoint factorization through three source packets. https://github.com/PeterNero/mtt-results-repro/blob/f141a20ea23c5c3ff19cc2161c0e226e29ade8a7/release/results/causal_base_q79_seven_row_endpoint_factorization_packet/artifact.json, 2026. Frozen source record; result causal_base_q79_seven_row_endpoint_factorization_packet. Scope as stated in the text.
- [17] Peter Nero. Physical-family source dependency and analytic completion. https://github.com/PeterNero/mtt-results-repro/blob/f141a20ea23c5c3ff19cc2161c0e226e29ade8a7/release/results/q79_family_source_cutset/artifact.json, 2026. Frozen source record; result q79_family_source_cutset. Scope as stated in the text.
- [18] Ji-Xiang Fu and Shing-Tung Yau. The theory of superstring with flux on non-kahler manifolds and the complex monge–ampere equation. *Journal of Differential Geometry*, 78(3):369–428, 2008.
- [19] Allan Adams, Morten Ernebjerg, and Joshua M. Lapan. Linear models for flux vacua. *Advances in Theoretical and Mathematical Physics*, 12:817–852, 2008.
- [20] Peter Nero. Physical visible spectral support and source cut-set. https://github.com/PeterNero/mtt-results-repro/blob/f141a20ea23c5c3ff19cc2161c0e226e29ade8a7/release/results/q79_visible_spectral_cutset/artifact.json, 2026. Frozen source record; result q79_visible_spectral_cutset. Scope as stated in the text.
- [21] Peter Nero. Correction: fixed-fiber Picard data do not decide global spectral ranks. https://github.com/PeterNero/mtt-results-repro/blob/f141a20ea23c5c3ff19cc2161c0e226e29ade8a7/release/results/q79_eta9_framed_member_spectral_rank_exclusion/artifact.json, 2026. Frozen source record; result q79_eta9_framed_member_spectral_rank_exclusion. Scope as stated in the text.
- [22] Peter Nero. Poincare–Heisenberg object-level intertwiner. https://github.com/PeterNero/mtt-results-repro/blob/f141a20ea23c5c3ff19cc2161c0e226e29ade8a7/release/results/q79_poincare_heisenberg_intertwiner/artifact.json, 2026. Frozen source record; result q79_poincare_heisenberg_intertwiner. Scope as stated in the text.

- [23] Peter Nero. Invariant Bott–Chern injectivity and source reduction. https://github.com/PeterNero/mtt-results-repro/blob/f141a20ea23c5c3ff19cc2161c0e226e29ade8a7/release/results/q79_bott_chern_reduction/artifact.json, 2026. Frozen source record; result q79_bott_chern_reduction. Scope as stated in the text.
- [24] Peter Nero. Full Bott–Chern injectivity. https://github.com/PeterNero/mtt-results-repro/blob/f141a20ea23c5c3ff19cc2161c0e226e29ade8a7/release/results/unified_source_bott_chern_injectivity/artifact.json, 2026. Frozen source record; result unified_source_bott_chern_injectivity. Scope as stated in the text.
- [25] Peter Nero. Hidden projective source and existential HYM theorem. https://github.com/PeterNero/mtt-results-repro/blob/f141a20ea23c5c3ff19cc2161c0e226e29ade8a7/release/results/unified_source_hidden_projective_hym/artifact.json, 2026. Frozen source record; result unified_source_hidden_projective_hym. Scope as stated in the text.
- [26] Peter Nero. Minimal free left-current Fermi completion. https://github.com/PeterNero/mtt-results-repro/blob/f141a20ea23c5c3ff19cc2161c0e226e29ade8a7/release/results/q79_minimal_fermi_completion/artifact.json, 2026. Frozen source record; result q79_minimal_fermi_completion. Scope as stated in the text.
- [27] Peter Nero. Global Fitting descent and auxiliary spectral-divisor match. https://github.com/PeterNero/mtt-results-repro/blob/f141a20ea23c5c3ff19cc2161c0e226e29ade8a7/release/results/q79_global_fitting_descent/artifact.json, 2026. Frozen source record; result q79_global_fitting_descent. Scope as stated in the text.
- [28] Peter Nero. Rank-zero large-gauge kernel on the selected K3. https://github.com/PeterNero/mtt-results-repro/blob/f141a20ea23c5c3ff19cc2161c0e226e29ade8a7/release/results/q79_bk3_rank_zero_kernel/artifact.json, 2026. Frozen source record; result q79_bk3_rank_zero_kernel. Scope as stated in the text.
- [29] Peter Nero. Original-Jacobian Gate 1 campaign. https://github.com/PeterNero/mtt-results-repro/blob/f141a20ea23c5c3ff19cc2161c0e226e29ade8a7/release/results/eta9_gate1_campaign/artifact.json, 2026. Frozen source record; result eta9_gate1_campaign. Scope as stated in the text.
- [30] Peter Nero. Three-cycle B96 non-detection. https://github.com/PeterNero/mtt-results-repro/blob/f141a20ea23c5c3ff19cc2161c0e226e29ade8a7/release/results/eta9_three_cycle_b96_nondetection/artifact.json, 2026. Frozen source record; result eta9_three_cycle_b96_nondetection. Scope as stated in the text.
- [31] Peter Nero. Local eta9 source-tube consolidation. https://github.com/PeterNero/mtt-results-repro/blob/f141a20ea23c5c3ff19cc2161c0e226e29ade8a7/release/results/q79_eta9_local_source_tube_consolidation/artifact.json, 2026. Frozen source record; scope as stated in the text.
- [32] Peter Nero. Same-source B89 Deligne obstruction and candidate rejection. <https://github.com/PeterNero/mtt-results-repro/blob/>

- [f141a20ea23c5c3ff19cc2161c0e226e29ade8a7/release/results/q79_eta9_b89_same_source_rejection/artifact.json](https://github.com/PeterNero/mtt-results-repro/blob/f141a20ea23c5c3ff19cc2161c0e226e29ade8a7/release/results/q79_eta9_b89_same_source_rejection/artifact.json), 2026. Frozen source record; scope as stated in the text.
- [33] Joseph Polchinski. *String Theory, Vol. 1: An Introduction to the Bosonic String*. Cambridge University Press, 1998.
 - [34] Ashoke Sen. Superstring perturbation theory revisited. *Journal of High Energy Physics*, 2015(11):008, 2015.
 - [35] Ashoke Sen. Ultraviolet and infrared divergences in superstring theory, 2015. KIAS Newsletter invited article.
 - [36] Ashoke Sen. Bv master action for heterotic and type ii string field theories. *Journal of High Energy Physics*, 2016(2):087, 2016.
 - [37] Peter Nero. Mirror zero-zero finite-field stratum classification. https://github.com/PeterNero/mtt-results-repro/blob/f141a20ea23c5c3ff19cc2161c0e226e29ade8a7/release/results/q79_mirror_zero_zero_final_classification/artifact.json, 2026. Frozen source record; result q79_mirror_zero_zero_final_classification. Scope as stated in the text.
 - [38] Felix M. Haehl, R. Loganayagam, and Mukund Rangamani. Schwinger–keldysh formalism i: Brst symmetries and superspace. *Journal of High Energy Physics*, 2017(6):069, 2017.
 - [39] Michael Crossley, Paolo Glorioso, and Hong Liu. Effective field theory of dissipative fluids. *Journal of High Energy Physics*, 2017(9):095, 2017.