

World-in-World Genesis: Local Comparison Geometry and a Globalization Program

Peter Nero

September 2026
Version 6

Abstract

We give the precise mathematical content of the world-in-world proposal and separate it from its physical interpretation. The central object is a comparison field $Q_{\text{WW}} \in \Gamma(\text{Hom}(TP, TI))$ between two oriented rank-three bundles. Its nine local components split, at a nonsingular background, into three orientation and six strain directions. A selected flag resolves the strain sector into scalar, diagonal-shape, and shear sectors of dimensions $1 + 2 + 3$. Therefore $1 + 3 \times 3 = (1 + 3) + (1 + 2 + 3) = 4 + 6$ is an exact component identity. It is not ordinary manifold-dimension multiplication and does not derive a ten-dimensional spacetime, Lorentzian signature, gravity, matter, or quantization. We state the missing globalization theorem to the selected q79 Fu-Yau carrier, distinguish the shared phase circle from physical time, and formulate the additional gates needed for each proposed physical emergence. The current comparison uses a curved connection compiler and a nonlinear six-coordinate spectral-strain shadow, not the excluded linear root-space embedding. The full relative-phase quotient and its support boundary remain distinct from that shadow. The resulting paper is a rigorous local geometry theorem plus a falsifiable research program.

Revision note for version 6

Supersedes. Version 5; its release identity and revision note are retained.

Reason. The globalization program needed the later curved and nonlinear owner results, rather than an undifferentiated request for a rank-six map.

Resolution. Imports projective naturality, the curved connection compiler and support-stratified strain comparison through scoped owner references. Distinguishes identity-coordinate orthogonality from general polar variations and complexifies the real carrier for a complex-linear comparison.

Retained. The comparison-field transformation law, local $1 + 2 + 3$ projectors, component count and independent physical gates.

Open boundary. Physical endpoint, reduced Green, symmetry lift and action selection are not consequences of the local normal form. This is an unreleased authoring revision with no new physical source selection.

Revision note for this edition

Supersedes. *World-in-World Genesis: A Proto-Geometric Origin of Time, Gravity, Matter, and Quantization*, version 4.

Reason. A three-dimensional base and three-dimensional fiber were multiplied into nine manifold dimensions, and the resulting count was used to claim automatic emergence of spacetime and several physical sectors.

Resolution. Version 5 replaces that construction by $Q_{\text{WW}} \in \Gamma(\text{Hom}(TP, TI))$, proves the local polar and 1+2+3 strain decompositions, and states separate globalization/physical gates.

Retained result. The exact component identity $1 + 3 \times 3 = (1 + 3) + (1 + 2 + 3) = 4 + 6$ survives as local comparison geometry.

Remaining boundary. The q79 bundle-and-connection intertwiner, Lorentzian completion, gravity, matter, and quantization are not yet generated by this identity.

Contents

Revision note for version 6	1
Revision note for this edition	1
1 How to Read World-in-World Geometry	3
1.1 A concrete strain split	3
1.2 The logical pipeline	3
2 Claim discipline	3
3 The local comparison field	4
4 Polar and strain decomposition	4
5 What the ten-component identity says	5
6 Shared circle and physical time	6
7 Selected q79 global carrier	6
8 The globalization theorem	6
8.1 The curved target and the replacement spectral map	7
9 Physical emergence gates	7
9.1 Time	8
9.2 Gravity	8
9.3 Matter	8
9.4 Quantization	8
9.5 Mass and scalar relaxation	8
10 Relation to current calculations	8
11 Conclusion	8

1 How to Read World-in-World Geometry

The closest standard mathematical picture is the deformation gradient in continuum mechanics. A three-dimensional body can carry a 3×3 linear map comparing two local frames without becoming nine-dimensional. The nine numbers describe how one three-dimensional fiber is compared with another at the same base point. Polar decomposition then separates a local rotation from a positive strain.

World-in-world geometry uses this comparison structure as a candidate pre-projection language. The local linear algebra is exact. Its proposed relation to a four-dimensional Lorentzian base and six-dimensional q79 internal carrier is a separate globalization problem.

1.1 A concrete strain split

For example, consider the symmetric matrix

$$S = \begin{pmatrix} 2 & 4 & 5 \\ 4 & -1 & 6 \\ 5 & 6 & 2 \end{pmatrix}.$$

Its trace is 3, so the scalar part is

$$P_1 S = I_3.$$

Relative to the displayed ordered frame, the traceless diagonal and off-diagonal parts are

$$P_2 S = \begin{pmatrix} 1 & 0 & 0 \\ 0 & -2 & 0 \\ 0 & 0 & 1 \end{pmatrix}, \quad P_3 S = \begin{pmatrix} 0 & 4 & 5 \\ 4 & 0 & 6 \\ 5 & 6 & 0 \end{pmatrix}.$$

They add back to S and are mutually orthogonal in the Frobenius inner product. The spaces from which these pieces are drawn have dimensions 1, 2, and 3. The numerical entries are not those dimensions, and the three subspaces are not three separate manifolds.

1.2 The logical pipeline

The paper advances through four stages:

1. define one local comparison field and prove its orientation/strain decomposition;
2. select a flag and obtain the local $1 + 2 + 3$ strain profile;
3. compare that profile with the independently selected q79 rank- $1 + 2 + 3$ carrier; and
4. construct the typed curved and nonlinear comparison, with its metrics, domains and effective operators, before treating the two as one physical construction. A stronger linear identification has additional isometry tests.

Only the first two stages are local consequences of Q_{WW} . The third provides a nontrivial structural clue. The fourth is the proposed globalization theorem and remains the decisive bridge.

2 Claim discipline

This paper has three layers:

Proved local geometry. Bundle dimensions, polar decomposition, and the orthogonal component splittings stated below.

Selected downstream data. The q79 trace-split carrier and Fu–Yau branch as recorded by their independent theorem packets.

Research program. The proposed identification of local strain sectors with q79 vertical operators and, after that, with physical spacetime fields.

No statement in the third layer is promoted by suggestive equality of dimensions. A bridge must identify maps, transition functions, connections, actions, and observables.

3 The local comparison field

Let B be a smooth manifold and let $TP, TI \rightarrow B$ be oriented Euclidean rank-three bundles. The labels “outer” and “inner” describe the roles of the two fibers; they do not create separate universes or coordinate factors.

Definition 3.1 (World-in-world field). A world-in-world field is

$$Q_{\text{WW}} \in \Gamma(\text{Hom}(TP, TI)).$$

Under changes of oriented orthonormal frames $g_P, g_I : B \rightarrow SO(3)$, its local matrix transforms as

$$Q_{\text{WW}} \mapsto g_I Q_{\text{WW}} g_P^{-1}.$$

This transformation law is essential: the nine entries are not nine scalar fields independent of frame choice.

Theorem 3.2 (Bundle dimension versus comparison components). *If $\dim B = 3$, the total space of either rank-three vector bundle has dimension six. The bundle $\text{Hom}(TP, TI)$ has rank nine, so a section has nine local fiber components.*

Proof. Local bundle charts have the forms $U \times \mathbb{R}^3$ and $U \times \text{Mat}(3, \mathbb{R})$, respectively. Dimensions are $3 + 3 = 6$ and ranks are $3 \cdot 3 = 9$. $\square$

Thus the original intuition survives only as “three-dimensional outer space plus a 3×3 comparison field.” It does not survive as three independent three-dimensional manifolds.

4 Polar and strain decomposition

Assume $Q_{\text{WW}}(b) \in GL^+(3, \mathbb{R})$ on an admissible domain. Polar decomposition is unique:

$$Q_{\text{WW}} = RU, \quad R \in SO(3), \quad U = (Q_{\text{WW}}^T Q_{\text{WW}})^{1/2} > 0.$$

The logarithmic strain is

$$S = \log U \in \text{Sym}(3, \mathbb{R}).$$

Theorem 4.1 (Orientation–strain normal form). *In identity-normalized matrix coordinates, the tangent component space has the orthogonal Frobenius decomposition*

$$\text{Mat}(3, \mathbb{R}) = \mathfrak{so}(3) \oplus \text{Sym}(3, \mathbb{R}), \quad 9 = 3 + 6.$$

Proof. Every matrix X decomposes uniquely as $X = (X - X^T)/2 + (X + X^T)/2$. Skew and symmetric matrices are orthogonal for $\langle X, Y \rangle_F = \text{tr}(X^T Y)$ and have dimensions three and six. $\square$

The polar-coordinate dimensions remain $3 + 6$ at every invertible background, but their images in the ambient tangent need not be orthogonal. At $Q = U = \text{diag}(2, 1, 1)$, let $\Omega_{12} = -1$, $\Omega_{21} = 1$ and let $\delta U_{12} = \delta U_{21} = 1$, with other entries zero. Then $\langle \Omega U, \delta U \rangle_F = 1$. Thus the proof establishes the skew/symmetric split, not an orthogonal polar tangent splitting at every background. Costs and metrics in polar coordinates must be pulled back; the $1 + 2 + 3$ projectors below act on the symmetric strain coordinate itself.

Choose an orthonormal flag, equivalently a preferred ordered frame modulo the declared residual symmetry. For $S \in \text{Sym}(3)$ define

$$P_1 S = \frac{\text{tr } S}{3} I_3, \quad P_2 S = \text{diag}(S) - P_1 S, \quad P_3 S = S - \text{diag}(S).$$

Theorem 4.2 (Selected $1 + 2 + 3$ strain split). *The maps P_1, P_2, P_3 are mutually orthogonal projectors with ranks 1, 2, 3, and*

$$S = P_1 S + P_2 S + P_3 S, \quad \|S\|_F^2 = \sum_{j=1}^3 \|P_j S\|_F^2.$$

Proof. $P_1 S$ is scalar diagonal, $P_2 S$ is traceless diagonal, and $P_3 S$ has zero diagonal. These subspaces are pairwise Frobenius-orthogonal, their dimensions sum to six, and the displayed formulas follow directly. $\square$

The $1 + 2 + 3$ split is flag-dependent. Without a selected flag, the canonical $SO(3)$ -equivariant split is $1 + 5$ into trace and traceless symmetric tensors. The flag is therefore source data that must either be selected by q79 geometry or declared as an assumption.

5 What the ten-component identity says

If one additional scalar u records an ordering coordinate, the local data have the count

$$\underbrace{1}_u + \underbrace{9}_{Q_{\text{WW}}} = \underbrace{(1+3)}_{u \text{ and orientation}} + \underbrace{(1+2+3)}_{\text{strain}} = 4 + 6.$$

Proposition 5.1 (No automatic spacetime theorem). *The preceding equality does not imply a splitting $TM_{10} \cong TY_4 \oplus TX_6$, does not select a Lorentzian metric on a four-dimensional factor, and does not identify the strain bundle with a compact internal tangent bundle.*

Proof. A component count contains no transition functions for TM_{10} , no integrable distributions, no metric of Lorentzian inertia, and no compactness or topology data. Each is independent structure. $\square$

The identity is nevertheless useful: it gives a precise local carrier whose four orientation/order and six strain components can be compared with a separately supplied $4 + 6$ physical realization.

In other words, the count is a compatibility test for a proposed map, not a construction of that map. A mismatch would rule out a simple component-preserving identification. A match allows the comparison to proceed to topology, connection, and operator tests, where genuinely global obstructions can appear.

6 Shared circle and physical time

Let $L_{\text{shared}} \rightarrow X_6$ be a Hermitian line bundle with unitary connection. Its circle fibers encode common phase or holonomy. Reusing this line in all three rank lanes does not add three circle coordinates and does not create a seventh internal dimension.

Physical time in the canonical completion is a noncompact ordering variable on a Lorentzian base, represented by a propagator $U(t_2, t_1)$. A phase map

$$\mathbb{R} \longrightarrow S^1, \quad t \longmapsto e^{i\omega t},$$

can synchronize phase with time after t and ω are supplied, but it is many-to-one and cannot make compact phase identical to physical time. An arrow of time further requires an asymmetric dynamical law or boundary condition.

7 Selected q79 global carrier

For the selected degree-three cover $\pi : C \rightarrow B$, define

$$\mathcal{A} = \pi_* \mathcal{O}_C, \quad \mathcal{A}_0 = \ker \text{Tr}, \quad \mathcal{A} = \mathcal{O} \oplus \mathcal{A}_0.$$

The selected common carrier is

$$\mathcal{H}_{\text{q79}} = L_{\text{shared}} \otimes (\mathcal{O} \oplus \mathcal{A}_0 \oplus \mathcal{A}), \quad \text{rank} = 1 + 2 + 3 = 6.$$

The trace split is global and intrinsic to the cover. It is not a global ordered-sheet flag: connected transitive monodromy prevents such an ordering. The signed sheet representation has a local binary-dihedral spin lift, but strict global Spin closure still requires all branch-complement relator signs or the equivalent obstruction class.

The q79 Fu–Yau branch is the current selected compactification candidate. The auxiliary space $L(3, 1) \times \text{Nil}_3$ is not homeomorphic to it and is used only as an effective model or operator mnemonic. Literal $S^1 \times \text{Lens} \times \text{Nil}$ and literal manifold nesting are not premises of this paper.

8 The globalization theorem

The rank-six local strain bundle is real, while the trace-split bundle is complex. A complex-linear comparison first complexifies the source; an alternative real comparison needs a compatible real form of the target. In the former convention, the proposed stronger linear map is

$$\mathfrak{J} : (\mathbb{R}I_3 \oplus \mathcal{D}_0 \oplus \mathcal{O}) \otimes_{\mathbb{R}} \mathbb{C} \longrightarrow L_{\text{shared}} \otimes (\mathcal{O} \oplus \mathcal{A}_0 \oplus \mathcal{A}).$$

Conjecture 8.1 (Stronger linear world-in-world/q79 identification). On the selected q79 Fu–Yau branch there is a source-selected flag and an isometric bundle isomorphism $\mathfrak{J}$ preserving the $1 + 2 + 3$ filtration such that

$$\mathfrak{J} \nabla^{\text{WW}} = \nabla^{\text{HYM}} \mathfrak{J}, \quad \mathfrak{J} H_{\text{WW}} = H_{\text{q79}} \mathfrak{J},$$

and likewise for the retarded and overlap kernels used in the numerical source packets.

The conjecture has four independent tests:

1. transition functions agree on all overlaps;

2. the selected flag is monodromy-compatible rather than an ordered-sheet choice;
3. the two inner products and connections are intertwined; and
4. the resulting operator coefficients equal the source coefficients used downstream, with a declared normalization and error certificate.

Rank matching proves none of these clauses. The scalar-field convention makes the comparison well-typed, but the operator problem is not finite merely because the bundle rank is finite.

The conjecture is therefore falsifiable in several independent ways. The bundles may have incompatible characteristic data, monodromy may obstruct the flag, the connections may fail to be conjugate, or the transported Hessian may disagree with the downstream source operator. Failure of any one test would end this proposed identification without undoing the local orientation/strain theorems.

8.1 The curved target and the replacement spectral map

The QM owner now supplies the appropriate curved operator language [1, 2, 3]. A smooth finite matrix projector $p(x)$ describes a bundle over a base, not a truncation to finitely many base modes. Its Čech compiler retains the supplied connection using (p, Γ) : the Grassmann connection alone generally is not the physical connection. Unitary naturality preserves the connection, curvature, metric, domains and transported spectral windows; it does not make arbitrary raw Fourier windows invariant. The auxiliary Lens/Nil model and the trace-split CLN filtration therefore do not specify the global q79 topology. A selected external flag must not be confused with a parallel splitting of an irreducible physical HYM bundle.

The old phase-stable linear rank-six root-space route is excluded by the S_3 representation mismatch. It cannot implement the preceding stronger conjecture. The surviving spectral map is nonlinear:

$$\Phi(X) = (d_0, d_1, d_2, |z_{12}|^2, |z_{20}|^2, |z_{01}|^2), \quad X \in \text{Herm}(3).$$

It keeps diagonal values and edge intensities, discarding the triangle phase of $z_{12}z_{20}z_{01}$ [4]. The full relative-phase quotient is generically seven-dimensional; the six-coordinate image is its strain shadow. Neither the discarded phase nor the shared circle is time. Given a positive source Hessian's reduced Green G_Q , minimization at fixed linearized strain $J = D\Phi$ gives $(JG_QJ^*)^{-1}$ where the reduced covariance is invertible. This shorted Hessian does not assert an isometric embedding. For TT-only strain, project J first, before inverting its covariance.

The later global result treats all eight edge-support strata [5]. At a zero edge the first derivative vanishes while the quadratic normal jet is $|\delta z_e|^2$. Its normal strain cost is degree one, so an ordinary full six-dimensional boundary Hessian is excluded. Tangential shorting still applies on each fixed stratum. A rank-six surjective derivative or the required local C_4 derivative J_{DE} forces the regular all-nonzero-edge stratum; regularity is no longer an independent physical input under that contract. The physical endpoint, Green and same-source $S_3 \times C_4$ lift remain to be supplied. J_{DE} is a tangent operation, not a global automorphism of the nonnegative intensity cone. These results sharpen the globalization program without identifying the real logarithmic strain with the spectral intensities by fiat.

9 Physical emergence gates

The original narrative can now be expressed as separate targets.

9.1 Time

A noncompact parameter, physical propagator, and arrow-producing condition are required. Bookkeeping order or phase winding alone is insufficient.

9.2 Gravity

A Lorentzian metric or tetrad, connection, action or equations, constraints, and a stress tensor are required. The local polar factor R does not derive the Einstein–Hilbert action.

9.3 Matter

A spin bundle, finite representation data, Dirac operator, chirality and real structure, and a particle/observable interpretation are required. The local $1 + 2 + 3$ strain split alone does not select Standard Model representations.

9.4 Quantization

A complex state space, observable algebra, dynamics, probability functional, and gauge and constraint treatment are required. Discrete topology or a compact resolvent can produce discrete spectra but not the Born rule by itself. This local nonimplication does not reopen the canonical operational QM construction. Measurement is an ordinary physical interaction; this paper’s component count adds neither an apparatus model nor an outcome-selection law.

9.5 Mass and scalar relaxation

Mass requires a canonically normalized pole or dispersion relation. A Higgs mode requires a selected scalar representation and couplings. Positivity of a six-dimensional strain Hessian does not imply a unique radial Higgs.

10 Relation to current calculations

The finite 27×27 matrix, charged Yukawa profile, CKM, electroweak, threshold, and precision packets close embedded renormalized-SM equivalence at the declared one-shared-physical-primitive/profile standard. They constitute a nontrivial downstream realization. Standard BRST/Faddeev–Popov quantization and measured renormalized profile data are admitted at that standard; strict no-knob selection remains an upgrade. These computations do not retroactively prove the world-in-world globalization conjecture.

11 Conclusion

World-in-world geometry has a clean surviving core. It is a rank-three comparison field with an exact orientation/strain decomposition and a selected $1 + 2 + 3$ local normal form. Its physical promise is concentrated into one testable globalization problem rather than many automatic-emergence claims. The excluded linear root-space implementation does not falsify the local linear algebra or the nonlinear spectral-strain replacement. Physical identification now requires the latter’s selected endpoint and operator comparison, or a separately established stronger linear realization.

Computational Evidence and Reproducibility

The earlier capsule below records its own source snapshot. Its quoted status labels are not current obligation counts; the present curved and nonlinear imports are bound separately by the frozen bibliography citations. The q=79, finite-matrix, HYM, and internal TT rows give concrete targets for the proposed globalization map. They do not prove the missing intertwiner from local comparison strain to the physical q=79 vertical geometry. Standard Model profile rows are context, and the strict upgrade remains open.

The referenced rows are frozen to the curated results repository state identified below. Hashes are grouped in eight-character blocks for line breaking.

Repository: <https://github.com/PeterNero/mtt-results-repro>
Commit: 31247ebb 5c22f3fb b5443024 365433c6 ee0bff4a
Manifest: release/result_manifest.json
Manifest SHA-256:
fb399689 60b00584 631dbf53 1a708e18
ef928d6b 6d935119 c185d7f6 32b1e7cd

Tier labels are quoted verbatim from that manifest. A row used directly supports only the specific computational statement identified above; a corpus-state cross-check does not prove this paper's local theorems; and an open row is evidence of an unresolved obligation, never of closure.

Rows used directly in this paper.

- A13/gr_tt_support (*derived exact*).
Exact-branch internal TT support certificate; physical normalization remains open.
- A19/hym_wiener_contraction (*numeric certified*).
Weighted-theta Fourier-tail and Wiener contraction certificate.
- A11/q79_exact_audit (*derived exact*).
Executable q=79 exact-branch audit.
- A11/q79_exact_theorem (*derived exact*).
CRT q=79 theorem on the selected exact branch.
- A01/qutrit_weyl_27_matrix (*derived exact*).
Sparse 27x27 qutrit-Weyl left-action realization.

Corpus-state cross-checks.

- A01/charged_yukawa_higgs_profile (*profile replay*).
Versioned Yu, Yd, Ye and lambda_H profile packet.
- A14/ckm_prediction_profile (*numeric certified*).
Three selected CKM profile rows and uncertainty comparison.
- A01/current_global_lock (*profile replay*).
Current non-looping global status and source-certificate map.
- A01/direct_k_higgs_row (*derived exact*).
Promoted direct K_threshold.Omega_H.lambda row.
- A22/e6_qpsi_qcd_anomaly (*derived exact*).
E6 Qpsi matter/exotic QCD anomaly cancellation audit.
- A04/final_12_of_12_audit (*profile replay*).
Twelve-obligation embedded renormalized-SM equivalence audit.

- A40/neutral_two_primitive_profile (*profile replay*).
Measured two-splitting neutrino profile, masses and Dirac Yukawa rows.
- A02/precision_15_source_transport (*profile replay*).
Fifteen measured source coordinates, Jacobian and covariance transport.
- A02/precision_8x8_workspace (*profile replay*).
Eight-coordinate SMDR output with positive-definite 8x8 covariance.
- A01/strict_pew_row (*derived exact*).
Promoted P_EW source row at the declared one-shared-primitive standard.

Open boundary (not evidence of closure).

- A05/strict_upgrade_ledger (*open*).
Current 2/9 strict no-knob upgrade ledger.

No imported row changes theorem ownership or promotes a neighboring claim: all local statements retain their stated hypotheses, domains, and limitations.

References

- [1] P. Nero, *Modal Triplet Theory: From MTT to Quantum Mechanics*, projective HYM, Čech compiler, spectral-strain and support-stratification sections, current owner manuscript, 2026.
- [2] P. Nero, [Covariant projective-module HYM naturality](#), frozen result, 2026.
- [3] P. Nero, [Explicit curved connection compiler](#), frozen result, 2026.
- [4] P. Nero, [Intrinsic spectral-strain quotient and shorted Hessian](#), frozen result, 2026.
- [5] P. Nero, [Global support-stratified strain and boundary-Hessian exclusion](#), frozen result, 2026.
- [6] S. Helgason, *Differential Geometry, Lie Groups, and Symmetric Spaces*, Academic Press, 1978.
- [7] J.-X. Fu and S.-T. Yau, *The theory of superstring with flux on non-Kähler manifolds and the complex Monge–Ampère equation*, J. Differential Geom. 78 (2008).
- [8] P. Nero, *MTT Foundational Geometry Reconciliation*, internal theorem and verification packet, 2026.