

Closure Geometry and a Regime-Local Ten-Dimensional Action Ansatz

Corrected fourth edition

Peter Nero

July 2026

Abstract

We formulate a ten-dimensional effective action compatible with closure-strain and q79 carrier data. The metric, bundle $M_{10} \rightarrow Y_4$, compact fiber, field representations, and derivative expansion are declared realization inputs. The action is an ansatz on a specified regime, not the most general action and not a derivation of gravity from projection. We give the conditions under which a closure Hessian contributes to canonically normalized pole masses, a rank-one alignment projector yields one Higgs doublet, an internal operator has discrete spectrum, and a four-dimensional mode truncation is consistent. Curvature is defined through explicit connections rather than inferred from nonuniform strain. The selected q79 Fu–Yau branch is the compactification candidate, while Lens–Nil remains auxiliary. Existing finite-matrix and Standard-Model profile calculations are incorporated at their declared embedded-renormalized-SM tier; their profile inputs and imported BRST quantization are not reclassified as no-knob consequences of this action.

Revision note for this edition

Supersedes. *Closure Geometry and Unified Dynamics: A Ten-Dimensional Action for Mass, Scalar Relaxation, Quantization, and Curvature*, version 3.

Reason. The metric and Einstein–Hilbert term were imported while the action was described as derived and most general; closure cost, Hessian positivity, nil language, and nonuniform strain were also promoted directly to mass, one Higgs, quantization, and curvature.

Resolution. Version 4 declares a regime-local EFT ansatz, lists omitted operators, and supplies separate pole-mass, alignment-projector, compact-resolvent, curvature, and consistent-truncation gates.

Retained result. The action remains a useful conditional synthesis and can host the currently certified finite-SM profile branch.

Remaining boundary. A same-source q79 action, normalized zero modes, gauge-fixed Hessians, reduction error, and strict value selection remain to be constructed.

Contents

Revision note for this edition	1
1 Status and regime	2

2	Fields and typed geometry	2
3	Regime-local action ansatz	3
3.1	Omitted operators	3
4	Closure Hessian and physical masses	3
5	Scalar alignment and the Higgs gate	4
6	Discrete spectra and quantization	4
7	Curvature and integrability	5
8	q79 Fu–Yau specialization	5
9	Four-dimensional reduction	5
10	Gauge fixing, anomalies, and observables	6
11	Relation to the current numerical closure	6
12	Scoped action theorem	6
13	Conclusion	6

1 Status and regime

Fix an energy interval $E \ll \Lambda_{10}$ and an admissible field neighborhood $\mathcal{U}$ on which a local derivative expansion is valid. The word “minimal” below means that only a displayed subset of operators is retained for a stated calculation. It does not mean uniqueness under all symmetries.

The action requires as input:

1. a ten-dimensional smooth bundle $\pi : M_{10} \rightarrow Y_4$ with compact six-dimensional fiber X_x ;
2. a slab-local Lorentzian metric g_{10} with one time direction and a globally hyperbolic four-dimensional physical base;
3. gauge and spin/ Spin^c bundles with compatible connections;
4. the selected q79 Fu–Yau geometry and HYM data, or another explicitly declared compactification;
5. a field content and symmetry group; and
6. a power-counting rule and cutoff controlling omitted operators.

The component identity $1 + 3 \times 3 = 4 + 6$ motivates a local comparison carrier but supplies none of these global inputs.

2 Fields and typed geometry

Let $Q_{\text{WW}} \in \Gamma(\text{Hom}(TP, TI))$ be the rank-three comparison field and let

$$S = \frac{1}{2} \log(Q_{\text{WW}}^T Q_{\text{WW}})$$

be its local strain. A selected flag gives the strain projectors $P_{\text{sc}}, P_{\text{sh}}, P_{\text{nil}}$ of ranks 1, 2, 3.

For the q79 degree-three cover, the selected internal carrier is

$$\mathcal{H}_{\text{q79}} = L_{\text{shared}} \otimes (\mathcal{O} \oplus \mathcal{A}_0 \oplus \mathcal{A}), \quad \text{rank} = 1 + 2 + 3.$$

An action using S as the source of q79 fields must include or cite a global intertwiner $\mathfrak{I}_{\text{WW} \rightarrow \text{q79}}$ preserving metrics, connections, and the selected operators. This paper does not infer it from rank equality.

Let ϕ^A denote real scalar coordinates on the retained field manifold, A_M^a a gauge connection, Ψ a spinor in a declared representation, and H a selected complex scalar module when present.

3 Regime-local action ansatz

One two-derivative ansatz is

$$\begin{aligned} S_{10} = \int_{M_{10}} \sqrt{|g_{10}|} d^{10}x \left[\frac{1}{2\kappa_{10}^2} R_{10} - \frac{1}{4} k_{ab}(\phi) F_{MN}^a F^{bMN} - \frac{1}{2} G_{AB}(\phi) D_M \phi^A D^M \phi^B \right. \\ \left. - V(\phi) + i \bar{\Psi} \Gamma^M D_M \Psi - \bar{\Psi} \mathcal{M}(\phi) \Psi + \mathcal{L}_{H,B,\Phi} \right] + S_{\text{gf}} + S_{\text{gh}} + S_{\text{bdy}}. \end{aligned} \quad (3.1)$$

Here k_{ab} and G_{AB} must be positive on physical directions, $\mathcal{L}_{H,B,\Phi}$ records the selected flux/torsion/dilaton sector, and the gauge-fixing, ghost, and boundary terms are part of the definition of a quantized perturbative calculation.

Equation (3.1) imports the Einstein–Hilbert term and a Lorentzian metric. It therefore realizes gravity; it does not derive gravity from strain or projection.

3.1 Omitted operators

Unless forbidden by a stated symmetry, the effective action can also contain

$$R^2, \quad R_{MN} R^{MN}, \quad R_{MNPQ} R^{MNPQ}, \quad R F^2, \quad F^3, \quad F^4, \quad (D\phi)^4, \quad R(D\phi)^2, \quad \phi^n, \quad \bar{\Psi} \Gamma \Psi D\phi,$$

torsion and Chern–Simons terms, higher fermion operators, and higher derivatives. A truncation must bound their contribution by powers of E/Λ_{10} and the relevant curvature or field amplitudes.

Higher-curvature or higher-time-derivative terms can introduce extra degrees of freedom or ghosts if treated nonperturbatively. In an EFT treatment they are perturbative operators below the cutoff; any claimed fundamental completion requires a separate constraint/propagator analysis.

4 Closure Hessian and physical masses

Let $\varphi = 0$ be a stationary background for retained real fields and write the quadratic four-dimensional action after integrating the internal fiber as

$$S_4^{(2)} = \frac{1}{2} \int_{Y_4} \sqrt{|g_4|} dd^4x \left(Z_{AB} \partial_\mu \varphi^A \partial^\mu \varphi^B - M_{AB}^2 \varphi^A \varphi^B \right).$$

Theorem 4.1 (Pole-mass promotion). *Assume Z is positive definite, the quadratic operator is self-adjoint on its declared domain, and interactions admit the stated perturbative pole definition. At tree level the squared masses are the eigenvalues of*

$$Z^{-1/2}M^2Z^{-1/2}.$$

*A closure Hessian $H_{\mathcal{J}}$ contributes to physical mass only after a source theorem identifies $M^2 = C^*H_{\mathcal{J}}C$ for a normalized field map C .*

Proof. The field redefinition $\hat{\varphi} = Z^{1/2}\varphi$ canonically normalizes the kinetic term. The inverse propagator is then $p^2I - Z^{-1/2}M^2Z^{-1/2}$, whose zeros give the tree-level poles. The final statement follows because an unnormalized dimensionless cost is not the quadratic coefficient in the physical action until C and units are fixed. $\square$

Loop-corrected pole masses require the renormalized self-energy and a declared mass scheme. Eigenvalues of an internal Hessian are not automatically pole masses.

5 Scalar alignment and the Higgs gate

A positive Hessian on the six-dimensional strain sector proves local stability, not one scalar. Suppose instead that the selected finite fluctuation space V_{scal} carries the required gauge representation and there is a source-selected orthogonal projector

$$P_H : V_{\text{scal}} \rightarrow V_H, \quad \text{rank}_{\mathbb{R}} V_H = 4,$$

onto one complex weak doublet.

Proposition 5.1 (Conditional one-doublet reduction). *If P_H commutes with the quadratic gauge-fixed operator, the orthogonal complement is massive above the truncation gap or consistently removed, and the potential restricted to V_H has the Standard-Model symmetry-breaking form, then the retained scalar sector contains one Higgs doublet at that scale.*

The selected finite-algebra packet executes such a rank-four alignment projector on a raw rank-twelve real fluctuation space and removes eight real scalar directions at the declared profile tier. That is the relevant selection result. It must not be replaced by “the Hessian has a unique radial direction,” which is false without the projector and representation data.

6 Discrete spectra and quantization

Let D_X be a self-adjoint internal Dirac-type or Laplace-type operator on the compact fiber with elliptic boundary conditions when a boundary is present.

Theorem 6.1 (Internal spectral discreteness). *If $(D_X - i)^{-1}$ is compact, then the spectrum of D_X is discrete with finite-multiplicity eigenvalues accumulating only at infinity.*

This theorem justifies a Kaluza–Klein or finite spectral expansion. A nil group, nil boundary, or divergent cost does not by itself imply compact resolvent or isolated minima. Moreover, spectral discreteness is not a derivation of quantum probabilities. Quantization still requires a state space, observable algebra, dynamics, constraints, and probability rule.

7 Curvature and integrability

Let P_H now denote a projector defining a horizontal distribution in a declared bundle, and let ∇ be a connection. Its curvature is

$$F_{\nabla} = \nabla^2,$$

or locally $F = dA + A \wedge A$. Frobenius failure of a distribution is tested by the vertical part of $[X, Y]$ for horizontal vector fields X, Y .

The condition $\nabla S \neq 0$ says only that strain is nonparallel. It does not by itself imply nonintegrability, Riemann curvature, or the Einstein equations. Any curvature–strain coupling in (3.1) must be written with an explicit connection and varied to obtain its field equations.

8 q79 Fu–Yau specialization

The current selected global compactification candidate is the q79 Fu–Yau branch with its declared complex, flux, spectral-cover, and HYM data. Known Hull–Strominger/Fu–Yau mathematics supplies a legitimate class of heterotic backgrounds once anomaly cancellation and the relevant bundle conditions are satisfied.

The auxiliary geometry $L(3, 1) \times \text{Nil}_3$ can support model calculations but is not the same manifold: its cohomology already differs from the q79 branch. The action must choose one background and integrate over that background. Literal $S^1 \times \text{Lens} \times \text{Nil}$ and literal nesting are not used to define X_6 .

The shared circle is encoded by L_{shared} and its connection. It is counted once and is not a time coordinate.

9 Four-dimensional reduction

Let $\{\chi_n(y)\}$ be normalized internal modes and expand a field as

$$\Phi(x, y) = \sum_{n \in I_{\text{keep}}} \phi_n(x) \chi_n(y) + \sum_{r \in I_{\text{disc}}} \eta_r(x) \chi_r(y).$$

Setting all η_r to zero is a consistent truncation only if their exact Euler–Lagrange equations vanish on the retained ansatz:

$$\left. \frac{\delta S_{10}}{\delta \eta_r} \right|_{\eta=0} = 0 \quad \text{for every } r \in I_{\text{disc}}.$$

Theorem 9.1 (Conditional coherent reduction). *Assume the retained mode space is invariant under all nonlinear terms to the order considered, the discarded equations vanish on the retained ansatz, the discarded linear operator has gap $\lambda_{\text{gap}} > 0$, and the nonlinear source into discarded modes obeys $\|J_{\text{disc}}\| \leq \epsilon$. Then the leading eliminated field obeys*

$$\|\eta\| \leq \|L_{\text{disc}}^{-1}\| \epsilon \leq \frac{C\epsilon}{\lambda_{\text{gap}}},$$

and substituting it gives a controlled local four-dimensional effective action with the corresponding Schur–Feshbach correction.

Proof. Solve the discarded equation by the inverse on the gapped complement and use the stated resolvent bound. The effective correction follows by substitution or the Schur complement. Exact consistency is the special case $J_{\text{disc}} = 0$. $\square$

A spectral gap alone does not prove nonlinear invariance or exact truncation.

10 Gauge fixing, anomalies, and observables

A perturbative gauge theory built from (3.1) must include a common gauge-fixing, ghost, zero-mode, regulator, scale, and renormalization-scheme policy. BRST nilpotency and anomaly cancellation are equations to verify. Physical predictions require a functor from action parameters to renormalized local or scattering observables.

The current embedded-renormalized-SM closure imports standard BRST/Faddeev–Popov quantization at its declared profile tier. This is a valid reconstruction standard, but it is not a derivation of the BRST/path-integral or Born-record rules from MTT.

11 Relation to the current numerical closure

At the adopted one-shared-physical-primitive/profile standard, the calculation repositories lock the finite 27×27 matrix, the physical finite Dirac operator at profile tier, charged Yukawa magnitudes, CKM rows, electroweak and threshold rows, multi-loop precision transport, and the renormalized observable functor. The final audit reports embedded renormalized-SM equivalence at that stated standard.

This action paper neither demotes nor strengthens those certificates. It records what would be needed to derive the same data from one ten-dimensional source action:

1. the world-in-world/q79 bundle-and-connection intertwiner;
2. the selected normalized internal zero modes and overlap kernels;
3. one common gauge-fixed Hessian and transport policy;
4. the consistent-truncation or error theorem; and
5. independent source selection if strict no-knob closure is claimed.

12 Scoped action theorem

Theorem 12.1 (Regime-local action statement). *Given the geometric, representation, metric, gauge-fixing, and EFT inputs of this paper, action (3.1) defines a local covariant realization on the declared domain. Under compact-resolvent, alignment-projector, pole-normalization, and consistent-reduction hypotheses, it yields a discrete internal mode expansion, a selected one-doublet scalar sector, canonically defined tree-level masses, and a controlled four-dimensional effective action.*

The theorem is conditional. It does not establish uniqueness of the action, derive the metric or dimension, select the q79 vacuum, or prove the full Standard Model without the listed source and observable maps.

13 Conclusion

The corrected action is a useful synthesis ansatz, not a universal derivation. Its value is that every physical promotion now has a recognizable mathematical gate: global geometry, connection, compact resolvent, scalar projector, canonical pole normalization, gauge consistency, and controlled reduction. The next decisive construction is the same-source q79 intertwiner followed by evaluation of the action’s normalized internal rows.

References

- [1] S. Weinberg, *The Quantum Theory of Fields, Vol. II*, Cambridge University Press, 1996.
- [2] J.-X. Fu and S.-T. Yau, *The theory of superstring with flux on non-Kahler manifolds and the complex Monge–Ampere equation*, J. Differential Geom. 78 (2008).
- [3] P. Nero, *MTT Current True SM Closure Consolidated Ledger*, internal theorem and verification packet, 2026.