

Spectral Delta Peaks, Resonances, and Survivor-Basin Encodings

Lorentzian Limits, Exact Spectral Atoms, and the MTT Source Boundary

Peter Nero

July 2026, Version 1

Abstract

Dirac delta functions appear in spectral physics as sharp energy levels, stable-particle poles, density-of-states atoms, spectral functions, and exact normal-mode decompositions. In ordinary notation one writes terms such as

$$\rho(E) = \sum_n \delta(E - E_n), \quad A(\omega) = 2\pi Z \delta(\omega - \omega_0) + \cdots.$$

This paper develops the spectral sequel to the delta-projection program in Modal Triplet Theory (MTT). The core mathematical statement is standard but structurally decisive: Lorentzian/Breit–Wigner kernels form approximate identities, and an identified broadened resonance family recovers a delta in its zero-width or infinite-lifetime limit. Genuine pure-point spectrum remains exact and need not arise by broadening.

We prove the weak convergence of Lorentzian kernels to the Dirac delta, derive the same kernel from exponential decay in time, and state the corresponding spectral-measure smearing theorem. We then classify sharp spectral atoms, resonances, finite lifetimes, LSZ poles, and density-of-states peaks. The MTT survivor-basin reading is conditional: a selected stability or leakage operator must produce the width before it can be interpreted as an intrinsic basin parameter.

Version 1 Revision Note

Supersedes The unversioned April 2026 manuscript on spectral deltas and resonances.

Reason The Lorentzian and spectral-measure results were correct, but the text did not always keep genuine pure-point spectrum distinct from a broadened-family limit, and it could identify a width with an MTT basin before a source map was supplied.

Resolution This version preserves exact spectral atoms, restricts the infinite-lifetime statement to identified resonance families, and makes the MTT survivor-basin reading conditional on a selected stability operator.

Retained result Lorentzian approximate-identity convergence, exponential-decay line shape, resolvent smearing, and weak convergence of spectral measures remain intact.

Remaining boundary MTT must derive the width-producing operator and distinguish intrinsic, environmental, and instrumental broadening.

Contents

1 Purpose and claim discipline	2
1.1 Non-claims	3

2	Sharp spectral atoms and finite-width peaks	3
3	Lorentzian approximate identity theorem	3
4	Finite lifetime produces Lorentzian width	4
5	Spectral-measure smearing theorem	5
6	Resolvent representation	5
7	Lifetime, width, and complex poles	6
8	Stable particles, LSZ poles, and resonance peaks	6
9	Kallen–Lehmann spectral representation	7
10	Finite-volume spectral atoms versus physical lifetime	7
11	MTT interpretation: survivor-basin idealization	7
12	Relation to the OU floor	8
13	Scope of proof and physical claim	9
14	Consequences for the delta-projection program	9
15	Conclusion	9

1 Purpose and claim discipline

The preceding papers in this sequence treated Dirac deltas as singular shadows of admissible projection in several settings: identity kernels, Green functions, gauge fixing, measurement, contact interactions, and scattering support. The present paper treats the spectral case.

The target object is the familiar spectral delta:

$$\delta(E - E_0).$$

It appears in:

- density of states;
- spectral functions;
- stable-particle poles;
- normal-mode decompositions;
- Fermi’s golden rule;
- Kallen–Lehmann-type spectral representations;
- condensed-matter quasiparticle peaks;
- resonance approximations.

The central claim is:

For an identified broadened resonance family, the sharp delta is its zero-width or infinite-lifetime limit.

1.1 Non-claims

We do not claim:

1. that self-adjoint operators cannot have genuine pure point spectrum;
2. that bound-state eigenvalues are artifacts;
3. that resonance theory is replaced by MTT language;
4. that all spectral broadening has one microscopic cause;
5. that finite lifetime is the only source of spectral width.

The narrower statement is that whenever an effective physical description writes a stable spectral contribution as a Dirac delta, MTT should ask which admissible survivor basin, lifetime scale, coherence width, detector resolution, or projection kernel has been idealized away.

2 Sharp spectral atoms and finite-width peaks

A normalized exact spectral atom at energy E_0 is written

$$\rho(E) = \delta(E - E_0).$$

If the mode has finite lifetime or finite resolution, the sharp atom is replaced by a broadened peak. The most common model is the Lorentzian or Breit–Wigner kernel

$$L_\Gamma(E - E_0) = \frac{1}{\pi} \frac{\Gamma/2}{(E - E_0)^2 + (\Gamma/2)^2}, \quad \Gamma > 0.$$

It is normalized:

$$\int_{\mathbb{R}} L_\Gamma(E - E_0) dE = 1.$$

As $\Gamma \downarrow 0$, it converges to $\delta(E - E_0)$ in the sense of distributions.

One possible MTT interpretation, requiring a sector source map, is:

$$\Gamma > 0 \quad \leftrightarrow \quad \text{finite survivor-basin width or finite lifetime,}$$

while

$$\Gamma = 0 \quad \leftrightarrow \quad \text{idealized infinitely stable survivor mode.}$$

The arrows are not identities established by the Lorentzian theorem. The theorem supplies the analytic encoding; MTT geometry must still derive which damping or leakage operator produces Γ .

3 Lorentzian approximate identity theorem

Definition 3.1 (Lorentzian kernel). For $\Gamma > 0$, define

$$L_\Gamma(x) = \frac{1}{\pi} \frac{\Gamma/2}{x^2 + (\Gamma/2)^2}.$$

Theorem 3.2 (Lorentzian kernels converge to the Dirac delta). *For every test function $f \in C_c^\infty(\mathbb{R})$,*

$$\lim_{\Gamma \downarrow 0} \int_{\mathbb{R}} L_\Gamma(E - E_0) f(E) dE = f(E_0).$$

Equivalently,

$$L_\Gamma(E - E_0) \rightarrow \delta(E - E_0)$$

in the sense of distributions.

Proof. Let $a = \Gamma/2$. Then

$$L_\Gamma(x) = \frac{1}{\pi} \frac{a}{x^2 + a^2}.$$

The functions L_Γ are nonnegative and normalized. Fix $\epsilon > 0$. Since f is continuous at E_0 , choose $\eta > 0$ so that

$$|f(E) - f(E_0)| < \epsilon$$

whenever $|E - E_0| < \eta$.

Then

$$\left| \int L_\Gamma(E - E_0) f(E) dE - f(E_0) \right| \leq I_1 + I_2,$$

where

$$I_1 = \int_{|E - E_0| < \eta} L_\Gamma(E - E_0) |f(E) - f(E_0)| dE \leq \epsilon,$$

and

$$I_2 = \int_{|E - E_0| \geq \eta} L_\Gamma(E - E_0) |f(E) - f(E_0)| dE.$$

Since f is compactly supported and bounded, $|f(E) - f(E_0)| \leq C_f$. Moreover the Lorentzian mass outside any fixed neighborhood tends to zero:

$$\int_{|x| \geq \eta} L_\Gamma(x) dx \rightarrow 0 \quad (\Gamma \downarrow 0).$$

Hence $I_2 \rightarrow 0$. Since ϵ was arbitrary, the result follows. $\square$

Corollary 3.3 (Breit–Wigner peak as regularized spectral delta). *A Breit–Wigner spectral contribution*

$$\rho_\Gamma(E) = Z L_\Gamma(E - E_0)$$

converges weakly to

$$Z \delta(E - E_0)$$

as $\Gamma \downarrow 0$.

4 Finite lifetime produces Lorentzian width

The Lorentzian is not merely a convenient regularization. It is the Fourier-domain signature of exponential decay.

Let a mode have time dependence

$$a(t) = \Theta(t) e^{-iE_0 t} e^{-\Gamma t/2}.$$

Its Fourier transform is, up to convention factors,

$$\hat{a}(E) = \int_0^\infty e^{iEt} e^{-iE_0 t} e^{-\Gamma t/2} dt = \frac{1}{\Gamma/2 - i(E - E_0)}.$$

Therefore

$$|\hat{a}(E)|^2 = \frac{1}{(E - E_0)^2 + (\Gamma/2)^2}.$$

After normalization, this is the Lorentzian $L_\Gamma(E - E_0)$.

Thus:

$$\boxed{\text{finite lifetime } \tau \sim \Gamma^{-1} \iff \text{finite spectral width } \Gamma.}$$

The spectral delta is recovered only when $\tau \rightarrow \infty$.

5 Spectral-measure smearing theorem

Let H be a self-adjoint operator on a Hilbert space $\mathcal{H}$, with spectral measure $P_H(dE)$. For a normalized vector $\psi \in \mathcal{H}$, define the scalar spectral measure

$$\mu_\psi(B) = \langle \psi, P_H(B)\psi \rangle.$$

A sharp spectral density is formal unless μ_ψ is absolutely continuous. But one can always form a Lorentzian-smeared spectral density:

$$\rho_{\psi,\Gamma}(E) = \int_{\mathbb{R}} L_\Gamma(E - E') \mu_\psi(dE').$$

Theorem 5.1 (Smeared spectral measures converge weakly). *For every bounded continuous test function f ,*

$$\lim_{\Gamma \downarrow 0} \int_{\mathbb{R}} f(E) \rho_{\psi,\Gamma}(E) dE = \int_{\mathbb{R}} f(E) \mu_\psi(dE).$$

Proof. By Fubini,

$$\int f(E) \rho_{\psi,\Gamma}(E) dE = \int \left(\int f(E) L_\Gamma(E - E') dE \right) \mu_\psi(dE').$$

The inner integral is $(L_\Gamma * f)(E')$, which converges to $f(E')$ pointwise as $\Gamma \downarrow 0$. Since f is bounded and L_Γ is normalized,

$$|(L_\Gamma * f)(E')| \leq \|f\|_\infty.$$

Dominated convergence with respect to the finite measure μ_ψ gives the result. $\square$

Remark 5.2. This theorem includes pure point, absolutely continuous, and singular continuous spectral measures. It does not say all spectra are physically broadened. It says Lorentzian broadening is a controlled approximation to any spectral measure and that delta atoms are recovered in the zero-width limit.

6 Resolvent representation

The Lorentzian also appears as the imaginary part of a resolvent:

$$L_\Gamma(E - E_0) = -\frac{1}{\pi} \text{Im} \frac{1}{E - E_0 + i\Gamma/2}.$$

More generally, for a self-adjoint H ,

$$\rho_{\psi,\Gamma}(E) = -\frac{1}{\pi} \text{Im} \left\langle \psi, (E - H + i\Gamma/2)^{-1} \psi \right\rangle.$$

Thus finite spectral width is equivalent to evaluating the resolvent off the real axis. The sharp spectral measure is recovered as the boundary value $\Gamma \downarrow 0$.

In physical language, Γ may encode finite lifetime, detector resolution, environment coupling, or damping. In a conditional MTT model it may encode the width of a survivor-basin representation. Those possibilities must not be conflated: instrumental convolution, environmental broadening, and an intrinsic pole width are different source mechanisms.

7 Lifetime, width, and complex poles

A finite resonance width is conventionally related to a finite lifetime by

$$\tau_{\text{life}} \sim \Gamma^{-1},$$

up to convention-dependent factors. The simplest way to see this is the exponentially decaying amplitude

$$a(t) = e^{-iE_0 t} e^{-\Gamma t/2} \mathbf{1}_{t \geq 0}.$$

Its Fourier transform has denominator

$$\hat{a}(E) \propto \frac{1}{E - E_0 + i\Gamma/2}.$$

Thus the resonance is represented by a pole displaced from the real axis:

$$E_{\text{pole}} = E_0 - i\Gamma/2.$$

The spectral peak on the real energy axis is the imaginary part of this off-axis pole contribution. A true spectral delta corresponds to the limiting case in which the pole approaches the real axis:

$$\Gamma \downarrow 0, \quad E_{\text{pole}} \rightarrow E_0.$$

Remark 7.1. This pole language is especially useful for separating three notions that are often blurred: an exact eigenvalue of a self-adjoint finite-volume operator, a stable infinite-volume particle pole, and an unstable resonance pole. All three may produce sharp or nearly sharp spectral features, but their mathematical status is different.

8 Stable particles, LSZ poles, and resonance peaks

In scattering theory, a stable one-particle state appears as a pole on the real mass shell or as a delta contribution to the spectral measure. A schematic spectral function has the form

$$A(\omega, \mathbf{p}) = 2\pi Z \delta(\omega - E_{\mathbf{p}}) + A_{\text{cont}}(\omega, \mathbf{p}).$$

For an unstable particle or quasiparticle, the sharp contribution is replaced by

$$A_{\Gamma}(\omega, \mathbf{p}) \approx 2\pi Z L_{\Gamma}(\omega - E_{\mathbf{p}}) + A_{\text{cont}}(\omega, \mathbf{p}).$$

For this resonance family, the delta peak is the infinite-lifetime limit:

$$\Gamma \downarrow 0 \implies A_{\Gamma} \rightarrow A.$$

Object	Finite-width form	Sharp limit
Stable bound state	narrow spectral atom	$\delta(E - E_n)$
Unstable particle	Breit–Wigner peak	pole approaches real axis
Quasiparticle	Lorentzian-like peak	infinite lifetime mode
Detector-limited line	resolution-broadened peak	ideal line spectrum
MTT survivor basin	finite basin width	zero-width survivor label

9 Kallen–Lehmann spectral representation

The same distinction appears in relativistic quantum field theory through the Kallen–Lehmann representation. For a scalar field, the two-point function can be written schematically as

$$\langle 0 | \phi(x) \phi(0) | 0 \rangle = \int_0^\infty d\mu^2 \rho(\mu^2) \Delta_+(x; \mu^2).$$

When the theory contains a stable one-particle state of mass m , the spectral density contains an atom:

$$\rho(\mu^2) = Z \delta(\mu^2 - m^2) + \rho_{\text{cont}}(\mu^2).$$

The coefficient Z is the pole residue or wave-function weight of the stable sector. If the would-be particle is unstable, the exact delta contribution is replaced by a broadened resonance contribution inside the continuum.

In the present terminology,

$$Z \delta(\mu^2 - m^2)$$

is the sharp idealization of an exactly stable survivor sector. A finite-width resonance is the corresponding finite-persistence version.

Remark 9.1. This section does not derive the Kallen–Lehmann representation. Its role is classificatory: it identifies one of the most important places where spectral delta functions enter QFT and shows that it fits the same delta-as-ideal-survivor pattern.

10 Finite-volume spectral atoms versus physical lifetime

A self-adjoint operator on a compact domain may have exact discrete eigenvalues. Its spectral measure may therefore contain genuine atoms even before any physical infinite-lifetime claim is made. These atoms are mathematical consequences of compactness, boundary conditions, or finite-volume quantization.

This paper distinguishes three cases:

Case	Meaning of the delta-like feature
Finite-volume eigenvalue	Exact atom of a compact/self-adjoint spectral problem.
Stable infinite-volume particle	Real-axis mass-shell pole with exact asymptotic persistence.
Unstable resonance	Finite-width peak from a pole off the real axis.

The MTT survivor-basin reading applies most directly to the second and third cases: the delta peak represents idealized exact persistence, while the resonance represents finite leakage or finite basin width. In finite volume, a delta may also reflect mathematical discreteness rather than physical infinite lifetime.

11 MTT interpretation: survivor-basin idealization

The triadic MTT carrier distinguishes bookkeeping, redundancy transport, and survivorship. Spectral peaks are naturally tied to survivorship: a mode that persists long enough to be tracked as a stable excitation appears as a spectral object.

Conditional on a source map identifying the resonance generator with an MTT stability operator, a finite-width resonance corresponds to a basin that is coherent but not infinitely stable. It has:

- a central energy or mass label;
- a finite damping or leakage rate;
- finite measurement resolution;
- finite admissibility width;
- possible coupling to continuum sectors.

A spectral delta can then encode an exact survivor:

$$\delta(E - E_0) \longleftrightarrow \text{zero-width encoding of an infinitely persistent survivor mode.}$$

This aligns with the fixed-point and measurement papers at the level of representation. It does not make exact pure-point spectrum approximate: a genuine eigenvalue remains an exact atom. The finite-width reading applies when a family of physical or instrumental broadenings has actually been identified.

12 Relation to the OU floor

In the MTT measurement and fixed-point stability papers, persistent disturbance balanced against damping produces an Ornstein–Uhlenbeck-type variance

$$\sigma^2 = \frac{\delta}{2\gamma}.$$

The proposed spectral analogue is:

$$\Gamma \sim \text{leakage/damping rate.}$$

If damping and disturbance leave a nonzero effective width, the spectral line does not become a true delta. Thus the exact delta limit requires not merely a central label, but removal of all effective broadening:

$$\Gamma \rightarrow 0, \quad \sigma \rightarrow 0, \quad \text{or resolution} \rightarrow 0.$$

This gives a diagnostic rather than a derivation:

When a spectral delta appears, ask which finite lifetime, basin width, or resolution scale has been suppressed.

13 Scope of proof and physical claim

Level	Status
Lorentzian approximate identity	Proved rigorously.
Finite lifetime to Lorentzian line shape	Standard Fourier calculation.
Smeared spectral measures	Proved as weak convergence.
Complex pole interpretation	Standard resonance-theory classification.
Kallen–Lehmann delta atom	Standard QFT spectral structure, used here classificatorily.
Resonance-family delta as infinite-lifetime limit	Standard physical interpretation.
Finite-volume spectral atoms	A distinct mathematical case, explicitly separated from lifetime claims.
MTT survivor-basin reading	Structural interpretation, not a new spectral theorem.
Exact derivation of widths from MTT carrier data	Not done here; belongs to execution-level work.

14 Consequences for the delta-projection program

This paper adds a new domain to the growing dictionary:

$$\delta(E - E_0) \rightsquigarrow L_\Gamma(E - E_0) \rightsquigarrow \text{finite survivor-basin spectral kernel.}$$

It shows that spectral deltas fit the same pattern as earlier cases:

- point-source deltas are zero-width source kernels;
- gauge-fixing deltas are zero-width section-selection kernels;
- measurement deltas are zero-width detector effects;
- contact deltas are zero-width overlap vertices;
- finite-time energy kernels converge to sharp energy deltas, while exact momentum-conservation support follows from translation symmetry;
- identified resonance families have infinite-lifetime delta limits;
- Kallen–Lehmann mass deltas are sharp stable-sector atoms.

15 Conclusion

Spectral delta peaks are among the most familiar idealizations in physics. They mark exact energy levels, stable-particle contributions, and perfectly sharp normal modes. The mathematical theory of self-adjoint operators allows genuine spectral atoms, and this paper does not deny that fact. Its point is different: in effective physical descriptions, a sharp spectral delta often encodes an idealization of infinite lifetime, zero width, perfect resolution, or exact survivor stability.

The Lorentzian/Breit–Wigner kernel makes this precise. It converges to the Dirac delta as its width tends to zero, and it arises directly from exponential decay in time. Smeared spectral measures converge weakly to the underlying spectral measure. Thus the delta peak is the singular endpoint of a controlled finite-width family.

In conditional MTT language, a broadened spectral family may be the finite-width encoding of a survivor basin. A genuine exact atom need not hide a width. The useful diagnostic question is therefore:

What finite lifetime, admissibility width, or survivor-basin structure has been collapsed into this sharp peak, if any?

References

- [1] H. Lehmann, “On properties of propagation functions and renormalization constants of quantized fields,” *Il Nuovo Cimento* **11** (1954), 342–357. [doi:10.1007/BF02783624](https://doi.org/10.1007/BF02783624).
- [2] M. Zworski, “Mathematical study of scattering resonances,” *Bulletin of Mathematical Sciences* **7** (2017), 1–85. [doi:10.1007/s13373-017-0099-4](https://doi.org/10.1007/s13373-017-0099-4).
- [3] M. J. Colbrook and A. C. Hansen, “Computing spectral measures and spectral types,” *Communications in Mathematical Physics* **384** (2021), 433–501. [doi:10.1007/s00220-021-04072-4](https://doi.org/10.1007/s00220-021-04072-4).