

Fixed Points III: Disturbance–Damping Balance and Stability

Peter Nero

September 2026

Version 6

Abstract

We develop disturbance–damping stability on the joint spectral decomposition of the FP–II internal operators. For each noncoherent joint mode α , the one-sided nonlinear damping margin is $\gamma_\alpha = d_\alpha - L_\alpha$. Deterministic force amplitude f_α and stochastic noise power q_α are distinct quantities. We prove the deterministic input-to-state floor $\limsup |a_\alpha| \leq f_\alpha/\gamma_\alpha$ and the stochastic second-moment floor $\limsup \mathbb{E}|a_\alpha|^2 \leq q_\alpha/(2\gamma_\alpha)$ when $\gamma_\alpha > 0$. Gaussian invariant laws and necessity of the sign condition are asserted only for exact Ornstein–Uhlenbeck dynamics or robust worst-case stability. Bundlewise results apply to $Q\Psi$ and use separate stochastic trace and deterministic weighted-series conditions. Finally, deterministic homogenization is made conditional on an enhanced functional central-limit theorem, tightness, and rough-path convergence; the symmetrized Green–Kubo tensor is $D = \int_0^\infty (R + R^*) ds$.

Version 6 Revision Note

Supersedes. Version 5, which remains the released edition. **Reason and resolution.** Separates the standalone analytic theorems from the physical source-selection problem. **Retained results.** The previous mathematical results and proofs are unchanged. **Boundary.** This is a contextual clarification in an unreleased revision, not a new physical-source theorem.

Revision note for version 5

Supersedes. *Fixed Points III: Disturbance–Damping Balance and Stability*, version 4.

Reason. The corrected deterministic and stochastic estimates in version 4 were sound, but their parallel notation could still obscure that force amplitude, noise power, invariant law, and homogenized noise are different objects.

Resolution. Version 5 adds an explicit argument map and worked interpretive discussion of the two disturbance lanes, their summability requirements, and the separate homogenization gate. It preserves theorem ownership and separates the public abstract from revision history and reproducibility metadata.

Retained result. The deterministic input-to-state floor, stochastic second-moment floor, OU restrictions, and enhanced invariance-principle conditions are unchanged.

Remaining boundary. Coherent-sector disturbances, general nonlinear invariant measures, and a physical stochastic source law remain unproved.

Revision note for version 4

Supersedes. *Fixed Points III: Disturbance–Damping Balance and Stability*, version 3.

Reason. Deterministic amplitudes and stochastic powers were conflated, nonlinear systems were assigned Gaussian/OU conclusions, and the Green–Kubo normalization and homogenization hypotheses were incomplete.

Resolution. Version 4 derives separate deterministic and stochastic floors on the joint modal spectrum, restricts Gaussian claims to exact OU dynamics, and states the enhanced invariance-principle and rough-path gates.

Retained result. Positive net modal damping still gives quantitative noncoherent stability and disturbance floors.

Remaining boundary. Coherent disturbances, nonlinear invariant laws, and physical stochastic interpretation require additional assumptions.

How to read this paper

FP–II separates the coherent sector $P\Psi$ from its damped complement $Q\Psi$. This paper asks what happens to the complement when it is continually disturbed. The answer depends on what kind of disturbance is meant, so the paper deliberately develops deterministic forcing and stochastic forcing in parallel without identifying their parameters.

The central picture in plain language. For each joint internal mode, damping removes amplitude at a rate d_α , while nonlinear feedback can return at most L_α in the one-sided energy estimate. Their difference $\gamma_\alpha = d_\alpha - L_\alpha$ is the available margin. A bounded deterministic push produces an amplitude floor proportional to f_α/γ_α . Brownian forcing deposits variance rather than a fixed amplitude and produces a second-moment floor proportional to $q_\alpha/(2\gamma_\alpha)$. The similar-looking denominators do not make f_α and q_α the same quantity.

Argument map. Sections 1–2 inherit the joint projector and build a non-double-counted spectral index for the overlapping vertical operators. Sections 3–4 derive the deterministic and stochastic modewise bounds. Section 5 identifies the limited settings in which positivity of the margin is also necessary and a Gaussian invariant law follows. Section 6 lifts individual mode bounds to the bundle by summability or covariance-trace conditions. Section 7 explains when rapidly mixing deterministic complement modes can instead appear as effective stochastic forcing of coherent variables. Sections 8–9 separate invariant measures from deterministic fixed points and turn the hypotheses into an execution checklist.

Scope boundary. Every modal result in the main stability chain concerns $Q\Psi$. It neither contracts coherent modes nor proves that a physical environment supplies Brownian noise. The homogenized diffusion is an emergent limit only after the enhanced functional-CLT, tightness, and rough-path hypotheses have been verified for the selected dynamics.

A physical application must independently select the damping generator, coherent projector, invariant domain, gap data, and forcing or noise law. The analytic estimates below consume those typed inputs; they do not emit them from MTT geometry.

Contents

Revision note for version 5	1
Revision note for version 4	2
How to read this paper	2
1 Scope and inherited framework	3
2 Joint modal decomposition	4
3 Two distinct disturbance classes	4
3.1 Deterministic forcing	4
3.2 Stochastic forcing	4
4 Modewise stability	5
5 Exact OU theorem and robust necessity	5
6 Bundlewise noncoherent stability	6
6.1 Stochastic trace condition	6
6.2 Deterministic weighted-series condition	6
6.3 Weyl-law check	6
7 Deterministic homogenization of coherent dynamics	6
8 Fixed points versus invariant measures	7
9 Examples and execution checklist	8
10 Conclusion	8
A Exact OU variance computation	8
B Uniform spectral gap for noncollapsing nil fibers	9

1 Scope and inherited framework

We use the corrected FP–I/II control framework. The compact internal space carries strongly commuting nonnegative self-adjoint operators A_1, A_2, A_3 on one common Hilbert space. Their joint harmonic projector is P , with $Q = I - P$. All stability results below concern $Q\Psi$ unless coherent forcing is explicitly introduced. The stabilization parameter is not physical Lorentzian time.

Deterministic fixed points of a time-step map and stochastic invariant measures are different objects. Existence of the former is inherited from FP–I/II under their invariant-set and compactness/condensing hypotheses. Existence of the latter requires a Markov/Feller and tightness argument stated separately below. For analytic-semigroup and dissipative-flow background, see [1, 6, 10]; for stochastic evolution and exact Ornstein–Uhlenbeck processes, see [9, 4]. The companion analytic framework is [8].

2 Joint modal decomposition

Strong commutation supplies a joint spectral calculus. In the compact pure-point case, choose a joint orthonormal basis $\{e_\alpha\}$ such that

$$A_n e_\alpha = \lambda_\alpha^{(n)} e_\alpha, \quad n = 1, 2, 3.$$

The multi-index α includes all three spectral labels and any multiplicity label. Define

$$d_\alpha := \sum_{n=1}^3 \kappa_n \lambda_\alpha^{(n)}, \quad \Lambda_\alpha := \sum_{n=1}^3 \lambda_\alpha^{(n)}.$$

Let $\mathcal{I}_Q$ contain the indices for which at least one $\lambda_\alpha^{(n)}$ is positive. Then

$$Q\Psi(t) = \sum_{\alpha \in \mathcal{I}_Q} a_\alpha(t) e_\alpha.$$

This joint index prevents multiple counting when the vertical structures overlap. It also applies to a genuinely nested realization after its inclusion maps have been supplied; rank labels alone do not establish nesting.

Assumption 2.1 (One-sided modal remainder bound). On the invariant set, the nonlinear modal remainder satisfies

$$\operatorname{Re}(\overline{a_\alpha} R_\alpha(a)) \leq L_\alpha |a_\alpha|^2, \quad \gamma_\alpha := d_\alpha - L_\alpha.$$

The remainder may couple modes; the displayed one-sided estimate, rather than an entrywise linearization identity, is the hypothesis used in the energy inequalities.

3 Two distinct disturbance classes

3.1 Deterministic forcing

For deterministic input $b_\alpha(t)$ assume

$$\|b_\alpha\|_{L^\infty(0,\infty)} \leq f_\alpha,$$

where f_α has units of amplitude per stabilization time. The mode equation is

$$\dot{a}_\alpha = -d_\alpha a_\alpha + R_\alpha(a) + b_\alpha(t). \quad (3.1)$$

3.2 Stochastic forcing

For the scalar real stochastic theorem let W_α be standard Brownian motions and let $q_\alpha \geq 0$ be noise powers. The mode equation is

$$da_\alpha = (-d_\alpha a_\alpha + R_\alpha(a)) dt + \sqrt{q_\alpha} dW_\alpha(t). \quad (3.2)$$

Correlated or infinite-dimensional noise is described by a positive covariance operator Q_ξ ; then the scalar q_α are replaced by its matrix entries and the bundlewise criterion is a weighted trace condition.

The deterministic amplitude f_α and stochastic power q_α are not interchangeable and are not represented by one common parameter.

4 Modewise stability

Theorem 4.1 (Deterministic input-to-state bound). *Assume Assumption 2.1 and $\gamma_\alpha > 0$. Every solution of Equation (3.1) obeys*

$$|a_\alpha(t)| \leq e^{-\gamma_\alpha t} |a_\alpha(0)| + \frac{f_\alpha}{\gamma_\alpha} (1 - e^{-\gamma_\alpha t}),$$

and hence

$$\limsup_{t \rightarrow \infty} |a_\alpha(t)| \leq \frac{f_\alpha}{\gamma_\alpha}.$$

Proof. The upper Dini derivative satisfies $D^+ |a_\alpha| \leq -\gamma_\alpha |a_\alpha| + f_\alpha$. Comparison with the scalar affine equation gives the result. $\square$

Theorem 4.2 (Stochastic second-moment bound). *Assume Assumption 2.1 and $\gamma_\alpha > 0$. Every solution of Equation (3.2) satisfies*

$$\frac{d}{dt} \mathbb{E} |a_\alpha|^2 \leq -2\gamma_\alpha \mathbb{E} |a_\alpha|^2 + q_\alpha,$$

and therefore

$$\mathbb{E} |a_\alpha(t)|^2 \leq e^{-2\gamma_\alpha t} \mathbb{E} |a_\alpha(0)|^2 + \frac{q_\alpha}{2\gamma_\alpha} (1 - e^{-2\gamma_\alpha t}).$$

Proof. Apply Itô's formula to $|a_\alpha|^2$, use the one-sided remainder bound, and take expectations. The martingale term has zero expectation and the quadratic variation contributes $q_\alpha dt$. $\square$

These two theorems are sufficient stability bounds for nonlinear dynamics. They do not assert that a nonlinear invariant law is Gaussian or that $\gamma_\alpha > 0$ is necessary for every particular forcing history.

What the two floors mean. Both estimates say that positive damping prevents indefinite accumulation in one mode, but they answer different questions. The deterministic theorem bounds every trajectory against a worst-case input amplitude. The stochastic theorem bounds an expectation after quadratic variation has injected power. Neither floor is automatically attained, and neither is a prediction until the mode, margin, and disturbance data have been obtained from the concrete operator model.

5 Exact OU theorem and robust necessity

Theorem 5.1 (Exact scalar OU classification). *For*

$$da = -\gamma a dt + \sqrt{q} dW_t, \quad q > 0,$$

there is a unique invariant probability law with finite variance if and only if $\gamma > 0$. It is Gaussian with mean zero and variance

$$\text{Var}(a) = \frac{q}{2\gamma}.$$

For $\gamma = 0$ the variance grows linearly, and for $\gamma < 0$ it grows exponentially.

Theorem 5.2 (Worst-case deterministic sign criterion). *The scalar family $\dot{a} = -da + R(a) + b(t)$ with $\text{Re}(\bar{a}R(a)) \leq L|a|^2$ is uniformly input-to-state stable for every bounded input and every admissible remainder only if $\gamma = d - L > 0$. For a specified forcing/remainder pair, $\gamma \leq 0$ does not by itself prove divergence because cancellation or a vanishing input may occur.*

Thus “if and only if” belongs to exact OU dynamics or a declared robust worst-case class, not to arbitrary nonlinear stochastic equations.

6 Bundlewise noncoherent stability

Under bounded geometry and the joint spectral calculus, assume

$$c_1 \sum_{\alpha \in \mathcal{I}_Q} (1 + \Lambda_\alpha) |a_\alpha|^2 \leq \|Q\Psi\|_{H_F^1}^2 \leq c_2 \sum_{\alpha \in \mathcal{I}_Q} (1 + \Lambda_\alpha) |a_\alpha|^2. \quad (6.1)$$

6.1 Stochastic trace condition

For independent exact OU modes define

$$\Sigma_q := \sum_{\alpha \in \mathcal{I}_Q} (1 + \Lambda_\alpha) \frac{q_\alpha}{2\gamma_\alpha}.$$

If every $\gamma_\alpha > 0$ and $\Sigma_q < \infty$, then

$$\limsup_{t \rightarrow \infty} \mathbb{E} \|Q\Psi(t)\|_{H_F^1}^2 \leq c_2 \Sigma_q.$$

For the independent exact OU product this condition is also necessary for a finite- H_F^1 second moment. For nonlinear dynamics it remains a sufficient bound unless additional lower estimates establish necessity. With correlated noise the correct replacement is the corresponding weighted covariance trace.

6.2 Deterministic weighted-series condition

Define

$$\Sigma_f := \sum_{\alpha \in \mathcal{I}_Q} (1 + \Lambda_\alpha) \frac{f_\alpha^2}{\gamma_\alpha^2}.$$

If every $\gamma_\alpha > 0$ and $\Sigma_f < \infty$, then

$$\limsup_{t \rightarrow \infty} \|Q\Psi(t)\|_{H_F^1}^2 \leq c_2 \Sigma_f.$$

This is an amplitude-series condition, not a stochastic trace.

6.3 Weyl-law check

If the joint counting function satisfies $N(\Lambda) \lesssim \Lambda^{d_{\text{eff}}/2}$ and $\gamma_\alpha \gtrsim 1 + \Lambda_\alpha$, then $q_\alpha \lesssim (1 + \Lambda_\alpha)^{-p}$ with $p > d_{\text{eff}}/2$ is sufficient for $\Sigma_q < \infty$. The exact exponent must be recomputed if the damping growth or joint multiplicities differ.

No statement in this section controls a disturbance acting directly in $P\Psi$. Such coherent forcing needs a separate base/coherent stability theorem.

7 Deterministic homogenization of coherent dynamics

This section addresses a different route to stochastic behavior. The underlying system can remain deterministic while rapidly mixing noncoherent modes drive slow coherent variables. Under a sufficiently strong limit theorem, the accumulated fast forcing converges to Brownian transport. Ordinary mixing is not enough: the second iterated integrals determine whether the limiting equation is the stated Stratonovich equation or carries an additional bracket drift.

Write $\Psi = X + Y$ with $X = P\Psi$ and $Y = Q\Psi$. For frozen x , let the fast flow for Y have invariant measure μ_x , and decompose

$$g(x, y) = \bar{g}(x) + G(x, y), \quad \bar{g}(x) = \int g(x, y) d\mu_x(y).$$

The correctly scaled slow-fast system is

$$\dot{X}_\varepsilon = f(X_\varepsilon) + \bar{g}(X_\varepsilon) + \varepsilon^{-1/2}G(X_\varepsilon, Y_\varepsilon), \quad (7.1)$$

$$\dot{Y}_\varepsilon = \varepsilon^{-1}L_{X_\varepsilon}(Y_\varepsilon). \quad (7.2)$$

For fixed x define the stationary covariance

$$R_x(s) = \int G(x, Y_s) \otimes G(x, Y_0) d\mu_x(Y_0).$$

Assumption 7.1 (Enhanced invariance principle). On the selected compact slow domain:

- (H1) the frozen fast flow has a unique invariant measure and sufficient mixing for an integrable covariance;
- (H2) the centered additive functionals satisfy a functional CLT uniformly in x , and their laws are tight in the required path topology;
- (H3) the lifted first and second iterated integrals converge to a geometric Brownian rough path, with zero area anomaly for the canonical Stratonovich statement below;
- (H4) the Poisson equation/corrector and its x -derivatives have the bounds needed to control slow dependence; and
- (H5) initial layers and exits from the compact slow domain are controlled on $[0, T]$.

For infinite-dimensional coherent spaces, the corresponding Hilbert-space tightness, trace, and rough-path assumptions must be supplied separately.

Theorem 7.2 (Conditional homogenized limit). *Under Assumption 7.1, X_ε converges in law on $C([0, T])$ to*

$$dX_t = (f(X_t) + \bar{g}(X_t)) dt + \sigma(X_t) \circ dW_t,$$

where

$$D(x) := \sigma(x)\sigma(x)^* = \int_0^\infty (R_x(s) + R_x(s)^*) ds. \quad (7.3)$$

If the enhanced limit has a nonzero area anomaly, an additional deterministic bracket drift must be included; if only an ordinary functional CLT is known, the Stratonovich conclusion is not established.

8 Fixed points versus invariant measures

For autonomous or τ -periodic deterministic forcing, the corrected FP-I/II Schauder or Darbo theorem applies after a deterministic invariant set is proved; the result is respectively a step-fixed or periodic point, with equilibrium promotion only under the FP-II Lyapunov condition.

For stochastic forcing, the solution defines a Markov semigroup. An invariant probability measure requires, for example, the Feller property, a Lyapunov moment bound, and tight Krylov–Bogoliubov averages. Uniqueness requires an additional irreducibility/coupling or contractivity argument. A stochastic invariant measure is not called a deterministic fixed point.

9 Examples and execution checklist

For a circle operator with eigenvalues m^2 and linear damping $\gamma_m = m^2 - L$, exact OU noise gives $\text{Var}(a_m) = q_m/[2(m^2 - L)]$, whereas bounded deterministic forcing gives $\limsup |a_m| \leq f_m/(m^2 - L)$. These are different floors.

For a compact Heisenberg nilmanifold in a noncollapsing left-invariant metric class, the first positive eigenvalue has a uniform lower bound. This supplies one ingredient in d_α but does not replace the joint-mode or summability checks.

For every concrete geometry:

1. construct the joint spectral index and multiplicities;
2. compute d_α and prove the one-sided constants L_α ;
3. list deterministic amplitudes f_α and stochastic covariance data q_α or Q_ξ separately;
4. check $\gamma_\alpha > 0$ and the appropriate Σ_f or covariance trace;
5. control coherent forcing independently; and
6. for homogenization, verify the enhanced invariance principle rather than only quoting mixing.

10 Conclusion

The damping-margin principle survives, but with precise scope. Positive γ_α gives deterministic input-to-state and stochastic second-moment bounds. Exact Gaussian invariant laws and sign necessity belong to exact OU or robust worst-case formulations. Bundlewise results concern $Q\Psi$ and require either a stochastic trace or deterministic weighted series. The homogenized Stratonovich equation is conditional on enhanced functional-CLT/rough-path data and uses the corrected Green–Kubo normalization. These distinctions prevent stochastic invariant measures from being conflated with deterministic fixed points.

The paper therefore contributes a translation layer between FP–II spectral damping and later statistical descriptions. Its robust content is the pair of modewise inequalities and their bundlewise summability conditions. Exact Gaussianity belongs only to the OU specialization, while emergent diffusion belongs only to the verified homogenization limit. Coherent disturbances, nonlinear invariant-law uniqueness, and a physical identification of the noise source remain separate model-specific tasks.

A Exact OU variance computation

For $da = -\gamma a dt + \sqrt{q} dW_t$,

$$a(t) = e^{-\gamma t} a(0) + \sqrt{q} \int_0^t e^{-\gamma(t-s)} dW_s.$$

Itô isometry gives

$$\mathbb{E} \left| \sqrt{q} \int_0^t e^{-\gamma(t-s)} dW_s \right|^2 = \frac{q}{2\gamma} (1 - e^{-2\gamma t})$$

when $\gamma > 0$.

B Uniform spectral gap for noncollapsing nil fibers

Proposition B.1. *Let $F = \Gamma \backslash \text{Nil}_3$ be fixed and let $\mathcal{G}_\epsilon$ be a compact class of left-invariant metrics satisfying $\epsilon I \leq G \leq \epsilon^{-1} I$. Then*

$$\inf_{g \in \mathcal{G}_\epsilon} \lambda_1(F, g) > 0.$$

The same bound is uniform for a smoothly base-dependent family remaining in $\mathcal{G}_\epsilon$.

Proof. The first positive eigenvalue varies continuously on this compact metric family and is positive for every compact connected fiber, so it attains a positive minimum; compare the spectral-geometric background in [3, 2, 7, 5]. $\square$

References

- [1] Herbert Amann. *Linear and Quasilinear Parabolic Problems, Vol. I: Abstract Linear Theory*. Birkhäuser, 1995.
- [2] Peter Buser. A note on the isoperimetric constant. *Annales Scientifiques de l'École Normale Supérieure*, 15(2):213–230, 1982.
- [3] Isaac Chavel. *Eigenvalues in Riemannian Geometry*, volume 115 of *Pure and Applied Mathematics*. Academic Press, 1984.
- [4] Crispin W. Gardiner. *Stochastic Methods: A Handbook for the Natural and Social Sciences*. Springer, Berlin, 4 edition, 2009.
- [5] Emmanuel Hebey. *Nonlinear Analysis on Manifolds: Sobolev Spaces and Inequalities*. American Mathematical Society, 2000.
- [6] Dan Henry. *Geometric Theory of Semilinear Parabolic Equations*. Springer, 1981.
- [7] Tosio Kato. *Perturbation Theory for Linear Operators*. Springer, 2 edition, 1976.
- [8] Peter Nero. Fixed points over multi-bundle manifolds: A general framework for emergent coherent structures. Companion paper; analytic foundations (bounded II, compact/condensing Φ_τ , existence/uniqueness, LS rates), 2025.
- [9] Giuseppe Da Prato and Jerzy Zabczyk. *Stochastic Equations in Infinite Dimensions*, volume 44 of *Encyclopedia of Mathematics and Its Applications*. Cambridge University Press, Cambridge, 1992.
- [10] Roger Temam. *Infinite-Dimensional Dynamical Systems in Mechanics and Physics*. Springer, 2 edition, 1997.

Computational Evidence and Reproducibility

The numerical and machine-verifiable claims used by this paper are archived in the curated repository, <https://github.com/PeterNero/mtt-results-repro>. The mapped authority/result identifiers are `no result rows mapped`. Claim tiers in that capsule distinguish exact derivation, certified numerics, profile replay, conditional results, and open obligations.