Conditional flow matching in acute human wounds: a three-donor audit of population reconstruction

Zeyu Fu[1,*]

[1]State Key Laboratory of Trauma and Chemical Poisoning,

Institute of Combined Injury, Chongqing Engineering Research Center for Nanomedicine,

College of Preventive Medicine, Army Medical University, Chongqing, China

*Correspondence: fuzeyu09@gmail.com; fuzeyu99@126.com

Supplementary methods and numerical results

Supplementary Note: Extended results

S0.1 Observed fibroblasts define the material available for temporal prediction

The physical research object is the recovered fibroblast population in separate healthy-skin wound biopsies, not a tracked fibroblast lineage. Three volunteers supplied intact skin and acute-wound biopsies at days 1, 7 and 30, yielding 12 specimens and 12,259 author-annotated fibroblasts (Figure 1a). The full-width schematic separates destructive sampling from two population-prediction questions: global missing-time reconstruction and held-donor transfer. The first permits a later biopsy and excludes day 7 in all donors; the second permits all times in the training donors but excludes the test donor's target. Neither depicts prospective source-only prognosis. A 7,002-gene input, with 5,383 covered genes, feeds frozen mean coordinates and training-only scaling before field fitting and numerical prediction. A common projection displays cells by collection time (Figure 1b) and donor (Figure 1c). Its first two principal components explain 43.5% and 10.6% of day-7-excluded training variance; nearly half remains outside this plane, so visual separation cannot replace evaluation in all 15 coordinates.

Matched raw-count profiles show the expression of matrix, inflammatory and contractile genes in each donor's four biopsies (Figure 1d). The gene-wise color scale compares a gene across samples and does not compare absolute expression between genes. Retained fibroblast counts differ across donor–time combinations (Figure 1e), so neither point density nor the number of cells provides independent temporal replication. Projected paths connect observed day-1 starting coordinates to model-predicted coordinates at the prespecified day-7 time (Figure 1f). These are numerical field trajectories; the data do not track the corresponding cells through the wound. Sampling counts are documented in Table S30.

Display-subset medians and interquartile crosses summarize time and donor structure in the same projection (Figure 1b–c). Focus windows retain full-range insets rather than removing the distant points. All saved predicted endpoints and a PC1 cumulative-distribution inset complement the paths (Figure 1f); the observed comparison uses the display subset and is descriptive, not a replacement for full-coordinate scoring or a confidence region.

S0.2 A held-out acute human time point can be reconstructed under the fixed protocol

Global timepoint holdout removes day 7 from every donor, whereas donor transfer removes one donor and retains all four times in the others. Both use the frozen representation and the training-only scaling in Equation (2). The product bridge in Equation (4), field objective in Equation (5) and initial-value prediction in Equation (7) specify a reproducible estimator, not an observed cell transition. This distinction determines whether the target time itself has been observed during training; the permitted source, target exclusions and fitting settings are recorded in Table S14.

A fixed-seed refit assessed numerical resolution of the RK4 update in Supplementary equation (S1) and sensitivity to evaluation sampling. Across 25, 50, 100 and 200 RK4 steps, the target energy distance remained 0.705395 and the improvement remained 49.863%; the largest discrepancy from the 200-step coordinates was below 6×10^{-7} in root-mean-square units (Table S27). Repeated evaluation at 250, 500 and 1,000 cells per distribution gave median improvements of 48.46%, 48.39% and 49.26%, respectively. Gain exceeded the same-count split-half reference in all 50 draws at each count (Table S28). These results assess one fitted field and the observed cells; the ranges across draws quantify evaluation sensitivity, not additional donor replication or confidence in clinical generalization.

Using the empirical weighting rule in Equation (17) with all cells and the same fixed 50-step prediction, equal cell weights gave a 49.40% improvement and equal donor mass gave 49.69%. The three donor-specific improvements under this pooled field were 47.15%, 46.92% and 40.07% (Table S29). This sensitivity changes the evaluation measure without refitting the field; it is distinct from the donor-transfer experiment and from the pooled capped-distance estimate.

The result is deliberately bounded. The donors are healthy volunteers, the wounds are acute experimental wounds and the target is one observed time point. The analysis does not contain a DFU patient trajectory or a time-to-closure endpoint. "Transfer" here means transfer across the three specified acute-wound donors under a held-out protocol; it does not mean clinical generalization.

Within-donor interpolation uses an observed later biopsy from the donor being reconstructed, which is permitted by the global timepoint-holdout task. It would not be available in prospective source-only prediction for a new donor. Conversely, the donor-transfer benchmark observes the target time in the training donors and asks a different question. Its method comparisons cannot be generalized to a globally unobserved time. The probability-conservation identity in Equation (8) concerns the constructed training paths, not identification of a biological trajectory. The observed reconstruction gains remain conditional on this representation, interval and three donors (Tables S15 and S23). Improvement over unchanged source alone does not establish incremental value over other permitted observations.

Using the empirical version (Equation (10)) of the energy discrepancy defined in Equation (9), the finalized held-out Wound7 analysis improved on the stay-still baseline by +49.9% under Equation (11) (Figure 2a). Dropping one donor at a time preserved improvement in 3/3 retained-donor sensitivity runs; these runs predict the retained donors, not the omitted donor. The held-out donor transfer analysis also passed in 3/3 folds, with a mean improvement of +75.9% (Figure 2b). In the geometry audit, the median same-donor temporal distance was 1.967 and the median same-time donor distance was 0.233, giving a time-to-donor ratio of 8.46 (descriptive 95% resampling range 6.02–13.11; Figure 2c). The mean displacement cosine was +0.948, with a descriptive resampling range of 0.943–0.954. Shared donors make these pair distances dependent, so the ranges do not have established donor-population coverage.

S0.3 Simple interpolation is competitive when an entire time point is missing

A retrospective extension tested whether improving over unchanged source requires a neural field. The additional comparisons hold out either day 7 or day 1 globally, keeping the frozen encoder, rank clock, training-only scaling and optimization budget fixed. For each task, the observed time immediately before the target supplies the source and the next observed time supplies an anchor. The source and anchor belong to the same donor; the held-time cells enter only the evaluation. Error is calculated using every available cell in all 15 coordinates within each donor, then averaged with equal donor weight. The donor-averaged error in Equation (19) differs from the earlier pooled and capped estimands. The two endpoint controls in Equation (18) therefore address model complexity under this separate, explicit scoring rule.

For withheld day 7, unchanged source had mean donor error 1.4725. Across three seeds, shared flow reduced it to 0.8110, while within-donor centroid translation reduced it further to 0.6674. An independent-endpoint linear bridge gave 0.9284 (Figure 3a). The corresponding improvements were 44.9%, 54.7% and 36.9%. That bridge is the one-segment population target of the product-coupling objective. The shared field's lower error is compatible with multi-segment fitting and finite capacity; it does not show that the objective targets a different interpolant, and translation remained lower. Withholding day 1 instead trained on intact skin, day 7 and day 30 and predicted from intact skin. The unchanged-source error was 1.3460; shared flow, centroid translation and the independent bridge gave 1.0571, 0.9293 and 1.0959, corresponding to improvements of 21.5%, 31.0% and 18.6% (Figure 3b). The complete seed-level errors appear in Table S31.

The centroid translation had lower seed-averaged error in all three donors for withheld day 7 and in two of three for withheld day 1 (Figure 3c). The flow error ranged from 0.8044 to 0.8237 across day-7 fits, and from 0.9484 to 1.1231 across day-1 fits (Table S31). Every fit improved over its unchanged source; Figure 3d displays these seed-specific relative improvements. These ranges characterize training variation in the existing cohort; they do not provide six independent biological replications. All donor-specific comparisons are retained in Table S32.

The added comparison supports temporal information in the measured distributions, while providing no evidence that a neural field is necessary for the two interpolation tasks tested here.

The location–scale predictor in Equation (20), which interpolates endpoint standard deviations as well as means, gave errors of 0.6982 for day 7 and 0.9034 for day 1, also lower than the shared-flow errors in both tasks (Figure 3a–b). Mean translation was the better of these two deterministic controls for day 7, whereas location–scale interpolation was better for day 1. This ordering is reported without selecting a new predictor for either target.

The distribution-shape diagnostics explain why a single aggregate error does not identify a uniform advantage. For day 7, flow had slightly lower centered energy distance than translation (0.1441 versus 0.1495), despite higher total error. For day 1, flow had higher centered error (0.1220 versus 0.0647), while the independent bridge also had higher centered error (0.1445; Figure 3e). The independent bridge's predicted-to-target variance ratios were 0.319 and 0.579 at days 7 and 1, indicating substantial contraction in this comparison (Table S33). These are separate geometry summaries, not a causal or additive error decomposition.

Matched 200-cell evaluation preserved the lower error of both deterministic endpoint controls in all 50 draws for each held time. Flow-minus-translation error ranged from 0.1132 to 0.1716 for day 7 and 0.0977 to 0.1545 for day 1; flow-minus-location–scale ranges were 0.0847–0.1405 and 0.1194–0.1763. Flow remained better than the independent bridge in every draw (Figure 3f; Table S34). The full-cell calculations reproduced all previous donor/seed errors to numerical precision. These results establish robustness of the observed ordering to the tested evaluation samples, conditional on three donors and the frozen fits.

S0.4 Time parameterization changes the held-out result

The historical single-seed, nearest-grid comparison gave linear days an improvement of −18.2% relative to unchanged source (Table S16). The revised paired audit instead evaluates 12 fields at the exact target time with identical random streams and evaluation rows within each seed. The day-based transforms in Equation (3) define separate fitted bridges, rather than transforming one fixed field by the chain rule in Equation (6). Mean donor energy distances were 0.8104, 2.1542, 0.8746 and 0.7725 for rank, linear days, square-root days and log days, respectively, versus an unchanged-source distance of 1.4746. The corresponding centroid translations gave 0.6667, 0.9775, 0.7476 and 0.6654, lower than the shared field under every clock on the donor-and-

seed average (Table S17; Figure 4a). Donor-specific flow-minus-centroid differences retain the heterogeneity behind this macro comparison (Figure 4b). These equal-donor, exact-time values have a different estimand and sampling cap from the historical pooled scan. Changing the coordinate changes bridges and optimization targets, so the comparison does not isolate interval compression or select a universal biological clock.

S0.5 The training-target distribution challenges the held-donor method ranking

The historical held-donor benchmark gave shared flow matching a mean energy distance of 0.3947 (approximately 0.395) and gains above the split-half reference in 3/3 folds. Mean displacement, nearest-donor displacement, entropic transport and the conditioned field also met that descriptive criterion. An added predictor instead uses only the training donors' Wound7 distributions, assigning equal donor mass and matching the held-source particle count. Its mean error was 0.1818, with a full range of 0.1479–0.2150 across 50 prediction-sampling draws (Figure 5a). The same historical target indices and scaling were used, so improvement over unchanged source does not establish incremental value from the excluded donor's source expression (Tables S18, S19 and S35).

The donor-specific results prevent a universal ranking: target-marginal mean errors were 0.1569, 0.1416 and 0.2469 for PWH26, PWH27 and PWH28, whereas shared-field errors were 0.7110, 0.2881 and 0.1850. The field therefore remained better in PWH28, including all 50 matched marginal draws. Those draws reuse the same three donors and quantify prediction-sampling sensitivity rather than population uncertainty.

The context model in Equation (13) improved all 6/6 training-donor comparisons (Figure 5b; Table S24) while giving a higher held-out mean than the shared field. Its greater fit to known contexts therefore supplies no evidence of a stable generalization advantage.

The donor-specific marginal comparison shows its full prediction-sampling range and the reversal in PWH28 (Figure 5c). Comparing the shared and conditioned fields in genuinely held donors (Figure 5d), rather than only training donors, prevents improved training fit from being read as generalization evidence.

The original neural and transport method comparison used one training seed and successive endpoint random draws. The entropic plan in Equation (14) is a separate comparator, not the coupling used to train the flow. The added target-marginal result is a retrospective baseline audit, and the historical fields were not saved as per-cell predictions for new paired target sampling. Neither result selects a model class for DFU patients; independent donors and matched repeated fits are required for that question.

S0.6 Donor-count comparisons require a common evaluation scale

The initial count curve changed its standardization when the training subset changed, so subtracting its absolute errors mixed information gain with a change of distance geometry. A controlled rerun retained subset-only training normalization but scored inverse-transformed predictions in one common reference per held donor. Across 27 fits, the hierarchical average in Equation (15) gave shared-field mean error of 0.5466 with one training donor and 0.3983 with two.

The same common geometry also showed lower error for mean displacement (0.5095 to 0.4186), the full equal-donor target marginal (0.2222 to 0.1676), and its matched-particle version (0.2332 to 0.1818). Unchanged-source error remained 1.4760 at both training sizes, checking that the reference metric did not change with k (Table S25). The full target-marginal decrease, however, is exactly the mixture effect in Equation (12); it cannot independently identify target-relevant information from adding a donor. The neural and mean-displacement comparisons involve different fitted predictors and are not algebraically identical to this target-marginal mixture.

Supplementary Fig. S1 shows all three seeds and the seed-0 line (0.5315 to 0.3854). Seed-specific reductions were 0.1462, 0.1558 and 0.1430 (Supplementary Fig. S1). The mean reduction was 0.1483, with all 3/3 donor blocks improving. The descriptive three-block percentile range [0.0732, 0.2727] is the smallest and largest of those three donor reductions (Table S20), not a calibrated 95% interval.

The donor-resolved curves retain the unequal reductions within the same three blocks (Supplementary Fig. S1). The separate 200-particle source audit distinguishes each individual alternative from their mixture (Supplementary Fig. S1); it is a sensitivity analysis of fixed fields, not another donor-count experiment.

The saved two-donor fields also allow the starting distribution to be changed without changing training, target cells or scoring scale. Averaged across seeds and donors, the actual held-source prediction had error 0.3983. Replacing its input with equal-donor draws of training Wound1 cells reduced mean error to 0.2450; the matched training Wound7 marginal, which does not use any source expression from the held donor, had error 0.1818 (Table S36). The donor effects were heterogeneous: real-source versus substituted-source errors were 0.7586 versus 0.2802 for PWH26, 0.2368 versus 0.2732 for PWH27, and 0.1994 versus 0.1816 for PWH28. Thus, PWH27 favored its own starting cells on average, while the other two did not. The substitution is a sensitivity analysis of the same fitted fields, not a comparison of nine independent donors; the 50 cell draws likewise supply no new biological replication. It does not demonstrate a consistent incremental value for the excluded donor's initial expression in this cohort, and it does not imply that such information is universally unhelpful.

The exact target-marginal reduction was 0.054601 across donors and matched one quarter of the training-target mutual discrepancy, with maximum numerical residual below 2.45×10^{-15} (Table S37). The additional 200-particle source audit gave macro errors of 0.4036 for real sources, 0.3075 when the two individual substituted-source errors were averaged, and 0.2527 for their sampled mixture. PWH28 exposes the distinction: real-source error was 0.2043, lower than either individual substitution (0.2503 and 0.2636), yet the mixed substitution was lower still at 0.1904. PWH27 also favored its real source over both individual alternatives, while PWH26 did not (Table S38). Thus, the earlier mixed-source comparison combines source replacement with a change

in the predicted mixture and cannot isolate individual source-state value. The exact identity does not decompose the gain from a jointly fitted two-donor field. All results remain conditional on the same three donors, fields, target cells and reference geometry.

Within a fixed donor and training subset, the mean training SD was 0.0701, or 3.07 times the mean split-half reference of 0.0228; the largest seed range was 0.3377 (Table S26). The aggregate direction persisted across seeds, while individual fits remained variable. The three-block range is descriptive because the held-donor folds share training donors. Two training sizes cannot determine a plateau or a required cohort size, and seeds, subsets and repeated marginal draws add no independent biological replication.

S0.7 Observable expression errors do not establish a flow advantage over target marginals

Raw-expression alignment and frozen re-encoding passed for all 12,259 cells, with maximum encoder-mean discrepancy 1.91×10^{-6}. On the exact saved targets and all 5,383 common-panel genes, shared-flow total variation was 0.2934 with one training donor and 0.2823 with two, versus 0.3529 for decoded persistence. The decoded mean-displacement control gave 0.2882 and 0.2825; decoded training-target marginals gave 0.2590 and 0.2546. Directly observed training-target marginals were lower still at 0.1279 and 0.1115, without using the excluded donor's source expression (Supplementary Fig. S1; Table S39). Flow improved over decoded persistence in all three donor blocks but over decoded target marginals in only one of three, and over observed target marginals in none. At two training donors its total variation was close to mean displacement, while its Jensen–Shannon divergence was higher (Supplementary Fig. S1), so these metrics do not establish a consistent neural advantage. The all-day-7 sensitivity retained the same aggregate interpretation. The observed-target decoder reference had mean total variation 0.2430 against actual expression (Table S40); because it uses the target itself, this is neither prediction performance nor an error floor. The results add a mean-expression evaluation within the existing cohort, not independent biological validation or evidence about unassayed genes.

S0.8 Finite-budget PCA diagnostics preserve the baseline concern

The separate train-PCA diagnostic completed all 48 flow fits and 12 baseline jobs across the same three donors. At both 120 and 600 steps, each of the four flows had higher within-donor mean energy than the fixed seed-0 target draw when averaging the three observed-source intervals and two flow seeds. At 600 steps, the flow-minus-draw differences ranged from approximately 0.145 to 0.184 for PWH26, 0.226 to 0.243 for PWH27 and 0.346 to 0.413 for PWH28 (Table S41). Individual intervals can favor flow; the within-donor means do not establish that every prediction loses or that a model class is universally inferior.

Longer budgets generally reduced unconditioned flow errors, but not every conditioned donor comparison improved. Mean shift and location–scale controls supplied no consistent flow advantage across donors and budgets. All 4,320 entropic-bridge training batches were retained; 20 failed the frozen two-sided marginal L1 tolerance of 0.001, with maximum row and column residuals of 0.002061 and 0.00001014 (Table S42). These are coupling diagnostics, not evidence of optimizer convergence or biological validity. Fold-specific PCA geometry, smaller architectures, finite budgets and asymmetric baseline sampling prevent substituting this experiment for a matched repeated comparison in the primary frozen coordinates. The 180 predictions and additional seeds do not increase the three-person biological sample.

S0.9 Three mouse arms expose model-arm heterogeneity

The all-arm audit gave different results under the same protocol, using the endpoint-distance diagnostic in Equation (16). NDB changed by −1.5% relative to stay-still and did not beat the baseline. PDB improved by +32.1%, but its POD7 target failed an endpoint-distance ordering check, warning against a simple interpolation interpretation without proving an off-manifold trajectory. GDB improved by +50.6% with a POD7 target satisfying that ordering check. In aggregate, 2/3 arms beat stay-still, 2/3 met the noise criterion and 2/3 satisfied the ordering check (Figure 6a). Each arm had one animal per time point, and NDB covered 59.7% of the frozen panel. These arm-level computations are useful for protocol auditing, but they do not establish a mouse biological trajectory, a cross-species equivalence or DFU transfer (Tables S22 and S21).

The paired baseline and predicted distances show the absolute error scale within each mouse arm (Figure 6b); relative improvement alone would hide these differences.

The complete arm–time cell counts expose unequal recovery (Figure 6c), while separate baseline, noise and endpoint-ordering outcomes retain every failed diagnostic (Figure 6d). Each count comes from one animal at that arm and time; neither more cells nor a passing computational check increases biological replication.

S0.10 Symmetric repeated evaluation preserves donor and information-set differences

All 107 jobs passed original-backend replay with zero maximum prediction discrepancy. The unchanged CPU replay tolerance nevertheless exposed cross-backend prediction discrepancies in some saved fields. That portability check remains failed rather than being certified by hash agreement or a looser tolerance. Original-CUDA-backend replay is distinguished from CPU portability, and neither replay identifies biological lineage or independently proves which observations generated a trained field. Numerical execution evidence and clinical validation consequently remain separate.

Three population-transport draws emitted Sinkhorn iteration warnings. A satisfied marginal check is not a proof of solver convergence.

The source-bound repeated experiment completed all 107 planned fitted jobs, retaining all permitted comparators and three neural seeds under a common architecture and 8,000-step optimizer budget. Missing-time ratios compare each method with mean translation inside the identical donor and draw. For day 7, mean log2 energy ratios were 0.367–0.666 for shared fields and 0.465–0.552 for context fields; all six donor–method means favored translation. For day 1, the shared-field PWH26 ratio was −0.0009,

close to parity, while the other five neural donor–method means favored translation (0.108–0.345). The location–scale control favored translation at day 7 but improved slightly over translation at day 1.

These results retain the near-parity and contrary comparisons rather than asserting universal neural inferiority (Figure 7a; Table S1).

Held-donor ratios instead use the observed training-target marginal, with day-1-to-day-7 results displayed separately and all adjacent intervals retained numerically (Figure 7b; Table S2). Shared-field log2 ratios were 2.372, 3.123 and −0.419 for PWH26, PWH27 and PWH28, respectively; context-field values were 2.518, 2.740 and −0.105. Thus, both neural donor means favored the marginal for PWH26 and PWH27, but favored flow for PWH28. For PWH28 the shared field favored flow in every seed's draw mean (−0.570 to −0.135), whereas context-field seed means crossed zero (−0.346 to 0.280). Fold-specific standardization is held fixed within each comparison; raw errors from different folds are not treated as one biological effect. No primary or secondary metric had an undefined paired ratio.

Source-replacement ratios compare the substituted and own source from the same field and target draw, retaining every donor, seed and substitute choice (Figure 7c). At day 7, shared-field means across seeds and individual substitute choices were −0.263, −0.965 and 1.502 for PWH26, PWH27 and PWH28; context-field means were 0.139, −1.007 and 1.322. Positive values favor the own source. Both neural families therefore favored the substituted source for PWH27 and the own source for PWH28, while PWH26 depended on the field. These fields exclude both the held donor and the substitute from fitting, which differs from replacing the source of a field trained on both remaining donors. No sampled mixture of the two substitutes is used in this comparison.

Fit costs count each fitted job once (Figure 7d). Shared and context neural jobs had median fit times of 9.64 and 9.45 seconds, respectively, with ranges of 8.49–15.77 and 8.77–18.71 seconds. These measurements exclude setup, prediction, evaluation and file operations; transport coupling constructed during prediction is not included in its comparator's fit time. They are not end-to-end efficiency rankings. Matched optimizer steps are not matched wall time. Fifty cell draws within each fit quantify computational sampling variability, not 50 new donors; the complete experiment still contains three healthy acute-wound donors.

Two warning draws were PWH27 day-0-to-day-1 draws with both returned marginals inside the fixed audit tolerance; the PWH26 day-7-to-day-30 draw had column L1 residual 1.061×10^{-6}, above the 10^{-6} bound, while its row residual was below 1.12×10^{-16}. These predictions remain in the comparison with their diagnostics; no warning or adverse draw was removed to improve a ratio.

S0.11 A cross-sectional mouse series admits one population for withheld-day reconstruction

The hashtag audit separates what GSE186527 can and cannot support (Table S3). All 5,353, 655 and 335 cells of the three unwounded and POD2 libraries have zero HTO counts, so their conditions cannot be assigned. The six later libraries have HTO counts in 431 to 892 cells at POD7 and POD14, but in only 218 and 217 cells at POD30, where the frozen rule gives 96 and 57 singlets. After gate assignment, the Lin− PBS and verteporfin populations at POD30 contain 34 and 27 cells, and the Lin+ verteporfin population contains 26 cells, all below the 50-cell floor (Table S4). The looser and stricter rules change these counts by at most eight cells and rescue none. Three of four arm and gate tasks therefore fail admission, including both fibroblast-enriched lineage-negative tasks. One population is admitted: the lineage-positive PBS arm, the non-fibroblast gate, with 264, 393 and 62 cells at POD7, POD14 and POD30 from four mice each.

For that population, POD14 reconstruction from POD7 with a POD30 anchor gave energy 0.441 for stay-still and 0.343 for the declared location–scale comparator (Table S5). Mean shift and quantile mapping gave 0.358 and 0.360, and the Gaussian geodesic gave 0.315. Three transport-based predictors were lower: OT displacement 0.248, the OT-coupled flow 0.235 (seed range 0.229 to 0.241) and the Schrödinger-bridge flow 0.241 (0.238 to 0.246), with natural-log ratios to location–scale of −0.326, −0.379 and −0.355. The independent-coupling flow gave 0.310 (0.287 to 0.332; natural-log ratio −0.103), and 0.284 under a log-days clock (natural-log ratio −0.180 against 0.340). The transport-coupled flows therefore exceeded the protocol's natural-log margin of 0.182, the natural logarithm of a 1.2-fold energy ratio and not the human target-marginal energy of the same displayed magnitude, but non-neural OT displacement reached the same range without a fitted field.

Scoring each prediction against the four held mice separately shows the scale of the nesting (Table S6). Mean per-mouse energies ranged from 0.85 to 1.02 across methods, with single values from 0.20 to 2.43. One held mouse differs strongly from the other three (energy 2.37 against their pooled cells), whereas the other three are 0.17 to 0.48 apart. Differences among methods are small beside this between-mouse spread. These values are descriptive: one population, four held mice, libraries confounded with day and arm, and a retrospective interpolation that uses the later day. Repeated scores do not add animals, identify trajectories or establish a population-level method ranking.

A second execution of the same nominal protocol and seeds saved reloadable fitted states for all 19 fits (Table S7). Its OT-coupled flow mean energy was 0.233 (seed range 0.223–0.240), versus the original 0.235 (0.229–0.241); the largest paired-fit energy change was 0.011673. Every newly saved prediction reproduced exactly under input-backed CUDA replay, with all population and held-mouse scores recomputed without predictor fitting. This verifies those saved states, not exact retraining of the original unsaved weights or additional biological replication; the small-comparator and no-neural-advantage interpretation is unchanged.

S0.12 Coupling and coordinates alter errors without establishing a consistent neural advantage

The additional human follow-up retained every declared fit. All 354 original fields and nine selected outer refits reproduced their full-source predictions exactly on CUDA; all 31 cached coordinate matrices also reproduced exactly from the input. A separate recomputation matched 2,700 deterministic-control evaluation rows to within 4.5×10^{-16}. These checks support the saved

computations, with the retrospective training-provenance and alpha-grid limits stated in Methods. The earlier CPU-portability failure remains unresolved.

For missing day 7, OT-coupled log2 energy ratios to translation were −0.026, 0.321 and 0.351 in PWH26, PWH27 and PWH28; entropic-bridge ratios were 0.016, 0.284 and 0.192 (Figure 8a; Table S8). Thus OT coupling favored flow in one donor, and neither coupled family favored flow in at least two donors. Withheld-day-1 OT ratios were −0.145, 0.183 and 0.122. Independent 40,000-step fits remained above translation in all three day-7 donors. These comparisons do not certify optimizer convergence or universal inferiority.

Across all five coordinate systems, the donor-mean day-7 ratio favored translation for both independent and OT fields (Figure 8b; Table S9). Individual donor signs nevertheless changed: the day-7 OT advantage in one donor under standardized coordinates disappeared under whitening and training PCA. Held-donor OT likewise favored one donor under standardized coordinates but none under centered raw or whitened coordinates. Repeated aggregate ordering does not make donor-specific conclusions invariant to the geometry.

Nested selection reduced held-donor day-1-to-day-7 errors compared with the default field, while leaving the comparison with the target marginal donor-specific (Figure 8c; Table S10). Selected-field log2 ratios to the marginal were 1.991, 0.825 and −0.471, versus default values 2.372, 3.123 and −0.419. All three selected seed means were positive in PWH26 and PWH27 and negative in PWH28. The bounded inner search improves this computational comparison without adding a donor or establishing population superiority.

The nested mixture in Equation (1) also retained contrary results (Figure 8d; Table S12). Day-1-to-day-7 independent-flow mixtures used weight 0.10 in every donor and gave log2 ratios of draw-averaged energy to the marginal of −0.029, 0.045 and −0.143. Selected mixtures therefore improved on the marginal in two donors and worsened it in one. These ratio-of-means values are a different estimand from the matched-draw means above. The geometry summaries alone do not determine whether source substitution helps or whether a held donor is outside the training population.

Training-cell decoding gave held-donor mean-expression total variation of 0.171 for independent flow, 0.149 for OT flow, 0.144 for location–scale and 0.113 for the target marginal (Figure 8e; Table S13). Decoding the observed target gave 0.084, showing substantial decoder mismatch even at observed coordinates. These errors assess mean gene proportions under an empirical decoder; they neither establish faithful cellwise expression nor replace the earlier topic-decoder audit. The displayed comparison contains all three held donors; the one-donor missing-time diagnostic is not promoted to a cohort result.

The target-informed alpha scan found a day-7 donor-mean minimum at alpha 0.55 for translation and 0.45 for OT flow (Figure 8f; Table S11). Their minimum errors were 0.700 and 0.817, compared with 0.705 and 0.819 at rank alpha 0.5. Changing the evaluation location of a fixed rank-trained field is distinct from the refitted-clock experiment. These oracle scans describe error surfaces after viewing the held target and do not select a validated temporal coordinate.

Supplementary Note: Extended discussion

Coupling variants and nested model selection qualify the interpretation of the default neural field. The completed outer refits reduced its errors, but selected fields still lost to the training target marginal in two of three held donors and improved in the third. Aggregate missing-day-7 comparisons favored translation across the five coordinate systems, while some donor-specific directions changed with geometry. These findings retain both favorable and adverse neural cases. The nested search uses the same small cohort, unequal configuration grids and a discovery-trained encoder; it therefore improves the computational audit without delivering an independent population ranking. Training-cell expression decoding and mixture shrinkage also address explicit alternative predictors rather than establishing biological dynamics. Their distinct decoders and ratio estimands prevent treating all numerical summaries as one pooled effect.

The symmetric repeated comparison confirms the baseline concern without making it universal. Missing-day-7 reconstruction favored mean translation in every neural donor–method mean, while held-donor day-7 reconstruction favored neural fields over the target marginal for PWH28 but not PWH26 or PWH27. Individual source replacements also had opposing donor-level directions under fields fitted without either compared donor. These results support reporting matched, donor-resolved comparisons rather than a pooled method rank. Complete same-backend replay reproduced every prediction, but failed CPU portability remains a numerical qualification, and additional draws or seeds do not expand the biological sample. A fitted population field brought day-1 cells closer to the observed day-7 distribution in the acute human wound cohort. Improvement occurred when day 7 was withheld globally and under a separate protocol that withheld each donor. These comparisons support reconstruction of the measured interval in three healthy volunteers, with different information available to the model in each task. Donor transfer can use day-7 observations from other people, whereas global timepoint holdout cannot. Keeping these tasks distinct is necessary when deciding what a successful prediction says about a future dataset or an individual patient [1, 2].

The global missing-time comparison changes the interpretation of model complexity. A within-donor mean translation had lower average error than the shared field for both withheld times, even though the field improved over an unchanged source. This shows why no-motion is an incomplete baseline for retrospective interpolation [3, 4]. The translation preserves the shape of the starting distribution and uses the observed later biopsy to estimate a displacement; its performance suggests that this simple construction captures enough of the measured change to be competitive under the chosen error metric. On one segment, an exact product-coupling solution targets the independent-bridge distribution rather than displacement interpolation. A shared finite-capacity field can depart from that bridge without thereby acquiring a designed advantage over translation; transport-coupled refits, whose objective is closer to displacement interpolation, still lost to translation in at least two withheld-day-7 donors. It

does not identify the biological process producing the shift, nor imply that all aspects of fibroblast heterogeneity are preserved correctly.

The donor-holdout task exposes a further baseline requirement. Because the target time is observed in training donors, their empirical target distributions can supply a prediction without using the excluded donor's starting expression. This target-marginal predictor had lower mean error than the historical shared-field estimate under matched prediction counts and the same target samples. In the separate three-seed saved-field audit, replacing the new donor's source with a mixture of training sources also lowered aggregate error without changing the model. That replacement changes both identity and mixture structure. At matched particle counts, PWH28 favored its real source over either individual training-donor substitution, yet favored the mixture over its real source; PWH27 favored its real source over both individual and mixed alternatives. These results prevent interpreting the aggregate mixture benefit as evidence that an individual's starting expression is universally unhelpful. Improvement over unchanged source remains insufficient to establish consistent incremental source-state value, but that value has not been isolated by these retrospective controls. Globally missing-time reconstruction excludes this target-marginal baseline because no training donor supplies that time. These distinct information sets must govern both baseline choice and the interpretation of apparent transfer [2, 5].

The observed-cell maps and biopsy expression profiles make the population being transported more concrete, but they do not establish a lineage or wound mechanism. A two-dimensional projection omits variation even when fitted solely to training cells [6]. Model curves remain one construction compatible with a specified training objective, and the data do not observe proliferation, migration, death or individual cell fates along those curves. Additional time-resolved biopsies from independent donors would test missing-time generalization; spatial measurements and lineage-informed experiments would address localization and the identity of connecting processes. A DFU cohort with authoritative clinical outcomes remains a separate requirement for prognostic claims.

The learned field is a model of distributions under a specified sampling construction. Independent endpoint cells from the same donor define linear paths and regression targets; those pairings are not observed cell lineages or an estimated minimum-cost coupling. At an ideal regression solution, probability conservation refers to that constructed path measure. It does not establish that an unobserved biological distribution follows the same path. Snapshot ambiguity and mathematical identification theory therefore remain relevant even when a held-out empirical distribution is predicted accurately [7, 8]. The result supports an empirical prediction within the evaluated protocol, while leaving the biological process connecting those observations underdetermined.

Additional biological measurements would constrain a different part of the problem. Lineage tracing relates ancestry to later cell states, whereas RNA velocity derives directional information from transcript processing under a kinetic model [9–11]. Probabilistic fate mapping and neural stochastic differential equations combine such information with further modeling assumptions [12, 13]. These methods were not evaluated here, and their published performance cannot be transferred to this wound cohort. Their relevance is that they identify possible sources of information beyond marginal expression distributions. Incorporating them would require compatible assays and a new validation design, rather than only changing the function used to fit the current endpoints.

Normalization to a probability distribution introduces another biological limit. Wound repair involves proliferation, loss, recruitment and matrix remodeling, with experimental evidence for coordinated fibroblast and macrophage responses [14, 15]. The present normalized distributions do not measure absolute tissue cell counts or distinguish those processes. A mass-preserving field can describe a change in a recovered state distribution without identifying whether it arose through movement of surviving cells, altered population growth or differential recovery during sampling. This explains why a computationally coherent flow can remain incomplete as a mechanism of tissue repair.

The time-axis comparison identifies a consequential modeling choice. The revised three-seed audit pairs the random streams and integrates to the exact target, addressing defects in the original single-seed comparison. Centroid translation still has lower mean error than the shared field for every tested clock. Changing the coordinate changes the specified bridges, target velocities and optimization weights; it is not simply a relabeling of time along one fitted field. Poor prediction under linear days and improvement under several alternatives show sensitivity within this experiment, but do not isolate interval compression as the cause or identify a universal biological clock. Selecting the transformation after inspecting the same target would make that target part of model selection. A later evaluation should separate selection from assessment, following the distinction emphasized in cross-validation studies [5, 16]. The scientific question is whether a fixed time rule transfers to independent observations, rather than whether one of several rules can fit the present interval.

Donor conditioning improved the fit to known donor contexts without improving the overall held-out mean relative to the shared field. Additional context can increase fitting capacity while also creating an extrapolation problem for an unfamiliar donor. A shared field already assigns different velocities at different latent positions, so different donor-average displacements do not establish that conditioning is necessary. The non-neural predictors also improved over unchanged source, showing that simple displacement rules capture part of the available structure. This makes them useful reference models when evaluating more flexible transport or response models [17]. With three test donors, the observed method ordering remains a description of this benchmark rather than an estimate of a stable population ranking.

The distance used for evaluation also determines the interpretation of improvement. Energy distance compares distributions through pairwise geometry and has a formal relationship to kernel discrepancies [18, 19]. Its value here depends on the frozen expression coordinates and training-only scaling. Centering the distributions exposed different shape behavior at the two withheld times, and independent endpoint interpolation contracted total dispersion. A deterministic rule that also interpolated endpoint standard deviations remained competitive. The tested shared field therefore offers no consistent distribution-shape advantage

across these two tasks. A lower aggregate discrepancy can coexist with errors in a rare subpopulation or in an individual gene that contributes little to that geometry. Representation benchmarking has shown why apparently successful integration must also be assessed for preservation of relevant biological variation [20]. The additional observed-expression audit evaluates average cellular proportions on the assayed part of the fixed panel. Flow improves over persistence, but not over the permitted training-target marginals on the aggregate primary metric; at two training donors it is close to mean displacement and the ordering depends on the expression metric. This evidence does not establish a consistent neural advantage. Substantial error remains even when decoding the observed target itself, which points to representation and reconstruction limits without defining a mathematical error floor. Common-panel conditioning and mean-profile evaluation leave unassayed genes, absolute counts, full cellular distributions, spatial tissue structure and clinical outcomes unvalidated.

The fitted and evaluated donor measures also differ. Cycling through donor–segment pairs approximately balances training contributions, whereas pooled evaluation weights cells according to yield. Equal-donor-mass evaluation of the same saved prediction preserved the improvement, and each donor separately improved under that pooled time-holdout field. This narrows the explanation that the favorable pooled result is driven only by unequal cell counts. It leaves other weighting choices unchanged, including scaling and fitting, and supplies no additional donor-transfer fold. The appropriate independent unit remains the person, not the recovered cell or the number of times a prediction is reevaluated [21].

Numerical integration and evaluation sampling were stable over the tested settings for one fixed field. Increasing Runge–Kutta resolution changed the coordinates negligibly, and repeated subsampling preserved improvement. The usual fourth-order convergence statement for the numerical integrator requires smoothness and does not itself certify this fitted field's prediction error [22]. These checks address numerical approximation and finite-cell evaluation, while training seeds revealed larger changes in some individual configurations. The shared-field error reduction with a second training donor persisted across seeds after fixing the scoring geometry within each held donor. Returning predictions to encoder coordinates before common-reference scoring removes the earlier metric change, but does not make every decreasing control curve independent evidence of information gain. For the target-marginal predictor, the reduction from the average single-donor error to the mixture error is fixed entirely by the distance between its two component distributions, without involving the test target. This exact identity does not decompose the improvement of a jointly trained neural field; it clarifies the baseline's estimand. Two training sizes still cannot identify a learning-curve plateau or a sufficient cohort size. The donor-block percentile summary uses bootstrap resampling [23], but overlapping training sets correlate the held-out comparisons, and three donor blocks provide little information about the tails of a population error distribution [24, 25]. Repeated computation therefore improves characterization of this experiment without replacing biological replication.

The finite-budget train-PCA diagnostic extends the baseline concern across multiple coupling variants and both budgets in all three donors. It remains a separate experiment: preprocessing, architecture, intervals and optimization settings differ from the primary frozen-coordinate analysis. Lower error after a larger budget does not make every fit converged, and retained marginal-tolerance failures remain relevant to numerical interpretation. Its single target-marginal draw left sampling stability incompletely matched to flow seeds. The completed frozen-coordinate comparison now uses matched repeated draws and three neural seeds, addressing that computational asymmetry without replacing the distinct PCA experiment or supplying a population ranking.

An external mouse series was admitted narrowly. Three of four arm and gate tasks failed the frozen 50-cell floor at the last time point, and the unwounded and POD2 libraries could not be assigned because no hashtag counts were released. In the one admitted population, the lineage-positive non-fibroblast PBS gate, optimal-transport displacement and the transport-coupled flows exceeded the declared natural-log error-reduction margin relative to location–scale. Both fibroblast-enriched lineage-negative gates failed the prespecified cell-count floor. The Gaussian and independent-coupling alternatives gave smaller reductions that did not reach that margin; translation and quantile mapping had higher errors than location–scale. Non-neural optimal-transport displacement reached the same error range as the transport-coupled flows, so the comparison does not establish a neural advantage. This agrees with the human audit's distinction between improving over persistence and adding value beyond permitted simple alternatives. It rests on four mice from one library per day, with day and arm confounded with batch and one held mouse that differs strongly from the others. It therefore neither ranks methods nor reconstructs a mouse trajectory, and it does not validate the human wound or diabetic-foot findings.

The mouse arms had different errors, one animal per time and incomplete panel coverage; time, animal and representation effects remain entangled. Neither acute human nor mouse wounds supply independently observed diabetic-foot-ulcer courses [26]. The contribution is a reproducible evaluation of conditional flow matching, not a new algorithm or reconstructed biological lineage. Permitted-information controls, common scoring coordinates and observed-expression comparisons show why improving over persistence does not establish added value over simple predictors. The mixture identity clarifies a baseline effect, not a biological mechanism. Independent donors under a frozen representation, clock and evaluation protocol are needed to assess generalization [2]; a clinical cohort with authoritative outcomes is separately required for prognosis.

The source-bound repeated comparison retains every donor/interval value. Its ratios compare methods inside the same fold and draw, rather than averaging raw distances across fold-specific coordinate scaling. All five metrics and individual source substitutions remain in the accompanying numerical data; the compact tables below display the complete primary energy comparisons.

Table S1. Complete missing-time paired energy comparisons against mean translation. Values are mean log2(method error/comparator error), averaging matched draws within fit and then training seeds within donor/interval. Negative values favor the listed method. Seed ranges span draw means, whereas draw ranges span the paired values across all fit seeds. Deterministic methods use one fit seed; neural fields use three. These computational ranges are not donor confidence intervals. No raw distance is averaged across fold-specific standardization.

Donor	Days	Method	Log2 ratio	Seed range	Paired draw range
26	0 to 1	Context CFM	0.108	0.027, 0.151	-0.059, 0.255
26	0 to 1	Shared CFM	-0.001	-0.087, 0.057	-0.184, 0.154
26	0 to 1	Persistence	0.372	0.372, 0.372	0.277, 0.473
26	0 to 1	Location–scale	-0.067	-0.067, -0.067	-0.100, -0.030
26	0 to 1	Mean shift	0.000	0.000, 0.000	0.000, 0.000
26	0 to 1	OT displacement	0.072	0.072, 0.072	-0.039, 0.218
27	0 to 1	Context CFM	0.229	0.218, 0.236	0.131, 0.309
27	0 to 1	Shared CFM	0.345	0.309, 0.397	0.237, 0.464
27	0 to 1	Persistence	0.499	0.499, 0.499	0.392, 0.599
27	0 to 1	Location–scale	-0.028	-0.028, -0.028	-0.064, -0.001
27	0 to 1	Mean shift	0.000	0.000, 0.000	0.000, 0.000
27	0 to 1	OT displacement	0.083	0.083, 0.083	0.010, 0.177
28	0 to 1	Context CFM	0.164	0.130, 0.213	0.027, 0.355
28	0 to 1	Shared CFM	0.240	0.217, 0.282	0.123, 0.433
28	0 to 1	Persistence	0.693	0.693, 0.693	0.595, 0.794
28	0 to 1	Location–scale	-0.018	-0.018, -0.018	-0.045, 0.015
28	0 to 1	Mean shift	0.000	0.000, 0.000	0.000, 0.000
28	0 to 1	OT displacement	0.053	0.053, 0.053	-0.068, 0.132
26	1 to 7	Context CFM	0.527	0.401, 0.666	0.329, 0.759
26	1 to 7	Shared CFM	0.367	0.284, 0.452	0.178, 0.531
26	1 to 7	Persistence	1.271	1.271, 1.271	1.154, 1.405
26	1 to 7	Location–scale	0.026	0.026, 0.026	0.010, 0.041
26	1 to 7	Mean shift	0.000	0.000, 0.000	0.000, 0.000
26	1 to 7	OT displacement	0.013	0.013, 0.013	-0.053, 0.094
27	1 to 7	Context CFM	0.465	0.387, 0.505	0.272, 0.637
27	1 to 7	Shared CFM	0.489	0.467, 0.514	0.339, 0.660
27	1 to 7	Persistence	1.162	1.162, 1.162	0.985, 1.384
27	1 to 7	Location–scale	0.105	0.105, 0.105	0.054, 0.152
27	1 to 7	Mean shift	0.000	0.000, 0.000	0.000, 0.000
27	1 to 7	OT displacement	0.257	0.257, 0.257	0.144, 0.338
28	1 to 7	Context CFM	0.552	0.428, 0.769	0.325, 0.915
28	1 to 7	Shared CFM	0.666	0.494, 0.860	0.392, 1.045
28	1 to 7	Persistence	0.790	0.790, 0.790	0.559, 1.007
28	1 to 7	Location–scale	0.056	0.056, 0.056	0.039, 0.067
28	1 to 7	Mean shift	0.000	0.000, 0.000	0.000, 0.000
28	1 to 7	OT displacement	0.389	0.389, 0.389	0.286, 0.572

Table S2. Complete held-donor paired energy comparisons against the target marginal. Values are mean log2(method error/comparator error), averaging matched draws within fit and then training seeds within donor/interval. Negative values favor the listed method. Seed ranges span draw means, whereas draw ranges span the paired values across all fit seeds. Deterministic methods use one fit seed; neural fields use three. These computational ranges are not donor confidence intervals. No raw distance is averaged across fold-specific standardization.

Donor	Days	Method	Log2 ratio	Seed range	Paired draw range
26	0 to 1	Context CFM	-0.338	-0.636, -0.129	-1.032, 0.237
26	0 to 1	Shared CFM	-0.489	-0.539, -0.451	-0.965, -0.061
26	0 to 1	Persistence	1.806	1.806, 1.806	1.518, 2.349
26	0 to 1	Location–scale	-0.357	-0.357, -0.357	-0.596, 0.122
26	0 to 1	Mean shift	-0.354	-0.354, -0.354	-0.574, 0.141
26	0 to 1	Population OT	1.186	1.186, 1.186	0.949, 1.570
26	0 to 1	Target marginal	0.000	0.000, 0.000	0.000, 0.000
27	0 to 1	Context CFM	0.145	-0.031, 0.430	-0.413, 0.744
27	0 to 1	Shared CFM	0.196	0.064, 0.331	-0.217, 0.700
27	0 to 1	Persistence	2.183	2.183, 2.183	1.725, 2.546
27	0 to 1	Location–scale	0.160	0.160, 0.160	-0.163, 0.447
27	0 to 1	Mean shift	0.214	0.214, 0.214	-0.113, 0.501
27	0 to 1	Population OT	1.304	1.304, 1.304	1.036, 1.544
27	0 to 1	Target marginal	0.000	0.000, 0.000	0.000, 0.000
28	0 to 1	Context CFM	1.502	0.997, 1.984	0.507, 2.798
28	0 to 1	Shared CFM	2.229	1.684, 2.678	1.241, 3.282
28	0 to 1	Persistence	2.616	2.616, 2.616	2.212, 3.100

Table S2 (continued)

Donor	Days	Method	Log2 ratio	Seed range	Paired draw range
28	0 to 1	Location–scale	-0.077	-0.077, -0.077	-0.580, 0.486
28	0 to 1	Mean shift	0.334	0.334, 0.334	-0.049, 0.901
28	0 to 1	Population OT	1.530	1.530, 1.530	1.096, 2.015
28	0 to 1	Target marginal	0.000	0.000, 0.000	0.000, 0.000
26	1 to 7	Context CFM	2.518	2.073, 2.848	1.329, 3.353
26	1 to 7	Shared CFM	2.372	2.342, 2.410	1.659, 2.934
26	1 to 7	Persistence	3.377	3.377, 3.377	2.761, 3.916
26	1 to 7	Location–scale	1.700	1.700, 1.700	1.077, 2.252
26	1 to 7	Mean shift	1.739	1.739, 1.739	1.108, 2.311
26	1 to 7	Population OT	2.049	2.049, 2.049	1.348, 2.665
26	1 to 7	Target marginal	0.000	0.000, 0.000	0.000, 0.000
27	1 to 7	Context CFM	2.740	2.611, 2.947	1.904, 3.737
27	1 to 7	Shared CFM	3.123	2.825, 3.375	2.278, 4.204
27	1 to 7	Persistence	3.282	3.282, 3.282	2.601, 3.832
27	1 to 7	Location–scale	1.218	1.218, 1.218	0.640, 1.951
27	1 to 7	Mean shift	1.288	1.288, 1.288	0.786, 1.964
27	1 to 7	Population OT	2.082	2.082, 2.082	1.394, 2.647
27	1 to 7	Target marginal	0.000	0.000, 0.000	0.000, 0.000
28	1 to 7	Context CFM	-0.105	-0.346, 0.280	-1.238, 0.831
28	1 to 7	Shared CFM	-0.419	-0.570, -0.135	-1.554, 0.298
28	1 to 7	Persistence	2.007	2.007, 2.007	1.274, 2.750
28	1 to 7	Location–scale	0.202	0.202, 0.202	-0.452, 0.620
28	1 to 7	Mean shift	0.420	0.420, 0.420	-0.231, 0.813
28	1 to 7	Population OT	1.585	1.585, 1.585	1.053, 2.090
28	1 to 7	Target marginal	0.000	0.000, 0.000	0.000, 0.000
26	7 to 30	Context CFM	0.121	-0.183, 0.311	-1.544, 1.088
26	7 to 30	Shared CFM	-0.006	-0.610, 0.520	-1.925, 1.153
26	7 to 30	Persistence	1.573	1.573, 1.573	0.437, 2.975
26	7 to 30	Location–scale	0.714	0.714, 0.714	-0.471, 1.634
26	7 to 30	Mean shift	0.803	0.803, 0.803	-0.510, 1.881
26	7 to 30	Population OT	1.114	1.114, 1.114	0.190, 2.232
26	7 to 30	Target marginal	0.000	0.000, 0.000	0.000, 0.000
27	7 to 30	Context CFM	1.248	0.848, 1.552	-0.457, 2.607
27	7 to 30	Shared CFM	0.600	0.250, 0.949	-0.655, 1.828
27	7 to 30	Persistence	3.648	3.648, 3.648	2.962, 4.433
27	7 to 30	Location–scale	1.103	1.103, 1.103	0.454, 1.846
27	7 to 30	Mean shift	1.502	1.502, 1.502	0.885, 2.224
27	7 to 30	Population OT	1.466	1.466, 1.466	0.274, 2.249
27	7 to 30	Target marginal	0.000	0.000, 0.000	0.000, 0.000
28	7 to 30	Context CFM	3.454	2.931, 3.779	1.645, 4.944
28	7 to 30	Shared CFM	3.543	2.995, 3.969	1.834, 5.233
28	7 to 30	Persistence	4.353	4.353, 4.353	3.295, 5.710
28	7 to 30	Location–scale	2.994	2.994, 2.994	1.882, 4.241
28	7 to 30	Mean shift	3.108	3.108, 3.108	2.043, 4.338
28	7 to 30	Population OT	2.784	2.784, 2.784	1.701, 4.173
28	7 to 30	Target marginal	0.000	0.000, 0.000	0.000, 0.000

Tables S3–S7 report the frozen-protocol external-mouse diagnostics and their repeated execution.

Table S3. Hashtag audit of the nine GSE186527 libraries. Day and arm for the six admitted libraries come from the library; mouse and gate come from the author-provided hashtag dictionary. Singlet, doublet and negative counts use the frozen primary rule (top count at least 20 and at least five times the second). The three unwounded/POD2 libraries contain no hashtag counts in any cell, so their conditions cannot be separated.

Accession	Condition	Cells	Any HTO	Singlet	Doublet	Negative	Admitted
GSM5429653	POD0/POD2 mixture	5353	0	–	–	–	No: zero HTO counts
GSM5429654	POD0/POD2 mixture	655	0	–	–	–	No: zero HTO counts
GSM5429655	POD0/POD2 mixture	335	0	–	–	–	No: zero HTO counts
GSM5429656	PBS POD7	589	501	399	65	125	Yes
GSM5429657	Verteporfin POD7	952	892	603	259	90	Yes
GSM5429658	PBS POD14	906	836	661	120	125	Yes
GSM5429659	Verteporfin POD14	470	431	294	97	79	Yes
GSM5429660	PBS POD30	474	218	96	76	302	Yes
GSM5429661	Verteporfin POD30	452	217	57	72	323	Yes

Table S4. Cells and mice per arm, day and sorting gate after the frozen primary hashtag rule, with the two sensitivity rules for comparison. Lin− is the fibroblast-enriched gate and Lin+ the remaining cells. A population is admitted at 50 cells and two mice. Mouse 5 is excluded from the Lin+ gate because the dictionary labels both of its tags Lin+ and the conflict is not resolved. No sensitivity rule rescues a population.

Gate	Arm	Day	Cells	Mice	Cells (loose)	Cells (strict)	Admitted
Lin-	PBS	POD7	131	4	133	126	Yes
Lin-	PBS	POD14	228	4	237	222	Yes
Lin-	PBS	POD30	34	2	34	33	No
Lin-	Verteporfin	POD7	205	4	214	197	Yes
Lin-	Verteporfin	POD14	143	4	152	133	Yes
Lin-	Verteporfin	POD30	27	2	35	26	No
Lin+	PBS	POD7	264	4	282	250	Yes
Lin+	PBS	POD14	393	4	399	375	Yes
Lin+	PBS	POD30	62	4	68	53	Yes
Lin+	Verteporfin	POD7	349	4	359	325	Yes
Lin+	Verteporfin	POD14	119	4	128	109	Yes
Lin+	Verteporfin	POD30	26	3	29	23	No

Table S5. Held-day reconstruction of the admitted lineage-positive PBS population, the non-fibroblast gate (source POD7, held POD14, later anchor POD30). Energy is the squared energy statistic in the 30-dimensional mouse-local PCA space fitted without POD14 cells; lower is better. Log ratio is the natural logarithm of energy divided by the location–scale energy. The last two columns score each prediction against the four held mice separately. One non-fibroblast population and four mice are a diagnostic, not a fibroblast or method ranking.

Clock	Method	Seeds	Energy	Range	Log ratio	Per-mouse mean	Per-mouse range
rank	identity	1	0.441	0.441, 0.441	0.250	1.017	0.408, 2.161
rank	mean shift	1	0.358	0.358, 0.358	0.042	0.962	0.325, 2.212
rank	location scale	1	0.343	0.343, 0.343	0.000	0.948	0.310, 2.198
rank	quantile map	1	0.360	0.360, 0.360	0.048	0.964	0.323, 2.210
rank	ot displacement	1	0.248	0.248, 0.248	-0.326	0.860	0.218, 2.143
rank	gaussian geodesic	1	0.315	0.315, 0.315	-0.084	0.922	0.281, 2.178
rank	cfm independent	3	0.310	0.287, 0.332	-0.103	0.976	0.276, 2.434
rank	cfm ot	3	0.235	0.229, 0.241	-0.379	0.854	0.202, 2.154
rank	cfm sb	3	0.241	0.238, 0.246	-0.355	0.858	0.207, 2.152
log days	location scale	1	0.340	0.340, 0.340	0.000	0.942	0.306, 2.185
log days	cfm independent	3	0.284	0.271, 0.292	-0.180	0.941	0.249, 2.358

Table S6. Descriptive scale references for the admitted population: energy between the pooled held day and the source or anchor day, and between each held mouse and the other held mice.

Reference	Mouse	Energy
held mouse vs other held mice	1	0.168
held mouse vs other held mice	2	0.481
held mouse vs other held mice	3	0.248
held mouse vs other held mice	4	2.375
source day vs held day	pooled	0.441
anchor day vs held day	pooled	0.610

Table S7. A second execution of the same nominal external protocol and seeds, without replacing the original results. Energy means and full seed ranges describe retraining variation, not new mice. The last column is the largest absolute paired-fit energy change. Input-backed CUDA replay of all 19 newly saved fitted states has zero prediction discrepancy; this is distinct from exact retraining reproduction of the original run, whose weights were not saved.

Clock	Method	Fits	Original mean (range)	Repeat mean (range)	Max fit change
rank	identity	1	0.441 (0.441, 0.441)	0.441 (0.441, 0.441)	0.000000
rank	mean shift	1	0.358 (0.358, 0.358)	0.358 (0.358, 0.358)	0.000000
rank	location scale	1	0.343 (0.343, 0.343)	0.343 (0.343, 0.343)	0.000000
rank	quantile map	1	0.360 (0.360, 0.360)	0.360 (0.360, 0.360)	0.000000
rank	ot displacement	1	0.248 (0.248, 0.248)	0.248 (0.248, 0.248)	0.000000
rank	gaussian geodesic	1	0.315 (0.315, 0.315)	0.315 (0.315, 0.315)	0.000000
rank	cfm independent	3	0.310 (0.287, 0.332)	0.310 (0.286, 0.332)	0.000080
rank	cfm ot	3	0.235 (0.229, 0.241)	0.233 (0.223, 0.240)	0.011673
rank	cfm sb	3	0.241 (0.238, 0.246)	0.241 (0.238, 0.246)	0.000019
log days	location scale	1	0.340 (0.340, 0.340)	0.340 (0.340, 0.340)	0.000000
log days	cfm independent	3	0.284 (0.271, 0.292)	0.284 (0.271, 0.292)	0.000039

The completed human follow-up separates matched-draw ratios, target-informed alpha diagnostics and nested mixture ratios of mean energies. The compact summaries retain all three donors and all declared outer seeds.

Table S8. Coupling and optimizer-budget follow-up under the original standardized frozen coordinates. Values are donor-specific means of matched-draw log2 energy ratios to mean translation; draws are averaged within fit, then three seeds within donor for fitted fields. Negative values favor the listed predictor. Exact OT interpolation samples from a full endpoint transport plan; the barycentric map uses its row means. Fifty draws reuse the same three donors. Longer training is an optimization diagnostic, not a convergence certificate.

Task	Predictor	PWH26	PWH27	PWH28
Missing day 7	Independent CFM	0.367	0.489	0.666
Missing day 7	Independent CFM, 40k	0.548	0.501	0.650
Missing day 7	OT-CFM	-0.026	0.321	0.351
Missing day 7	SB-CFM	0.016	0.284	0.192
Missing day 7	Exact OT interpolation	-0.014	0.255	0.367
Missing day 7	OT barycentric map	0.023	0.277	0.392
Missing day 7	Independent bridge	0.270	0.467	0.617
Missing day 7	Location–scale	0.026	0.105	0.056
Missing day 1	Independent CFM	-0.001	0.345	0.240
Missing day 1	Independent CFM, 40k	-0.008	0.308	0.184
Missing day 1	OT-CFM	-0.145	0.183	0.122
Missing day 1	SB-CFM	-0.106	0.170	0.114
Missing day 1	Exact OT interpolation	0.078	0.079	0.040
Missing day 1	OT barycentric map	0.094	0.101	0.041
Missing day 1	Independent bridge	0.255	0.246	0.188
Missing day 1	Location–scale	-0.067	-0.028	-0.018

Table S9. Representation sensitivity. Each row first averages the paired log2 ratios within donor, then gives the three-donor mean and count of donors below zero. Missing-time comparators are mean translation; held-donor comparators are training target marginals. Scaling, whitening and PCA preparation use permitted training cells only. Raw distances across these geometries are not commensurate. Sign agreement does not provide independent validation.

Coordinates	Task	Independent	Below 0	OT-CFM	Below 0
Standardized encoder	Missing day 7	0.508	0/3	0.215	1/3
Standardized encoder	Missing day 1	0.195	1/3	0.054	1/3
Standardized encoder	Held donor, day 1–7	1.692	1/3	0.956	1/3
Centered encoder	Missing day 7	0.378	0/3	0.121	1/3
Centered encoder	Missing day 1	0.146	1/3	0.114	1/3
Centered encoder	Held donor, day 1–7	1.285	1/3	1.500	0/3
Whitened encoder	Missing day 7	0.509	0/3	0.319	0/3
Whitened encoder	Missing day 1	0.316	0/3	0.163	0/3
Whitened encoder	Held donor, day 1–7	1.753	1/3	1.240	0/3
Sum-zero encoder	Missing day 7	0.509	0/3	0.221	1/3
Sum-zero encoder	Missing day 1	0.185	1/3	0.048	1/3
Sum-zero encoder	Held donor, day 1–7	1.719	1/3	0.955	1/3
Training PCA30	Missing day 7	0.231	0/3	0.099	0/3
Training PCA30	Missing day 1	0.215	0/3	0.097	0/3
Training PCA30	Held donor, day 1–7	0.910	1/3	0.995	1/3

Table S10. Nested model selection and fresh outer refits for held-donor day-1-to-day-7 prediction. Each held donor is excluded from both inner folds. The search contains 28 configurations with unequal grids across coupling families, scored with seed 0 on the two inner held donors. The selected configuration is refitted on both permitted donors with three seeds before outer evaluation. Last columns compare the default and selected fields with the same training target marginal using matched-draw log2 ratios. The bounded search does not equate optimization costs or exhaust all possible configurations.

Held donor	Selected family	Width	Steps	Rate	Default	Selected
PWH26	OT-CFM	256	2000	3e-04	2.372	1.991
PWH27	Independent CFM	64	20000	3e-04	3.123	0.825
PWH28	Independent CFM	64	8000	1e-03	-0.419	-0.471

Table S11. Target-informed alpha scan of fixed rank-clock fields and endpoint controls. Alpha spans 0 to 1 in increments of 0.05 along the permitted source-to-anchor segment. Each optimum uses the held target and is therefore an oracle diagnostic, not an independently chosen prediction time. Rank-alpha is 0.5; day-alpha is 6/29 for withheld day 7 and 1/7 for withheld day 1. Evaluating a rank-trained field at another alpha is not retraining under a different biological clock. Original saved alpha grids are retained without a new full-grid replay certificate.

Task	Predictor	Oracle alpha	Oracle ED	Rank ED	Day-alpha ED
Missing day 7	Mean translation	0.550	0.700	0.705	1.015
Missing day 7	Location–scale	0.550	0.734	0.736	1.031
Missing day 7	Independent CFM	0.450	0.986	1.000	1.126
Missing day 7	OT-CFM	0.450	0.817	0.819	1.047
Missing day 7	SB-CFM	0.500	0.790	0.790	1.063
Missing day 1	Mean translation	0.450	0.954	0.956	1.156
Missing day 1	Location–scale	0.500	0.932	0.932	1.151
Missing day 1	Independent CFM	0.550	1.108	1.108	1.225
Missing day 1	OT-CFM	0.450	0.998	1.004	1.175
Missing day 1	SB-CFM	0.450	0.998	1.007	1.169

Table S12. Nested mixture shrinkage for all three adjacent intervals and both transported-source families. Prediction mixes transported source with the training target marginal using weight w chosen from 0, 0.05, ..., 1 in the two inner donor folds, then applied to the outer held donor. Ratios are log2 of draw-averaged energies, then averaged over seeds; they differ from the mean of per-draw log ratios in the other follow-up tables. Zero weight recovers the marginal; negative ratios favor the selected mixture. Target-informed oracle weights are not used to make the selected predictions.

Transport	Held donor	Days	Selected w	Mixture/marginal
Independent CFM	PWH26	0–1	0.000	0.000
Independent CFM	PWH26	1–7	0.100	-0.029
Independent CFM	PWH26	7–30	0.100	-0.039
Independent CFM	PWH27	0–1	0.350	-0.066
Independent CFM	PWH27	1–7	0.100	0.045
Independent CFM	PWH27	7–30	0.000	0.000
Independent CFM	PWH28	0–1	0.450	0.849
Independent CFM	PWH28	1–7	0.100	-0.143
Independent CFM	PWH28	7–30	0.000	0.000
Mean translation	PWH26	0–1	0.200	-0.245
Mean translation	PWH26	1–7	0.300	0.192
Mean translation	PWH26	7–30	0.100	-0.092
Mean translation	PWH27	0–1	0.650	-0.033
Mean translation	PWH27	1–7	0.200	-0.196
Mean translation	PWH27	7–30	0.000	0.000
Mean translation	PWH28	0–1	0.550	-0.164
Mean translation	PWH28	1–7	0.150	-0.173
Mean translation	PWH28	7–30	0.250	0.822

Table S13. Held-donor expression evaluation by the mean profiles of 15 nearest permitted training cells. Each metric is averaged over fitted seeds within donor and then equally over the three donors. Expression uses per-cell proportions on 5,383 common genes. Fold-change Spearman correlations use 2,000 training-selected variable genes or a training-selected 50-gene signature. TV is total variation and JS uses natural logarithms. The decoded observed target is a decoder reference, not a predictor or a universal error lower bound. NA marks undefined fold-change comparisons for the unchanged observed source.

Predictor	TV	JS	HVG rho	Signature rho
Independent CFM	0.171	0.030	0.775	0.942
Independent CFM, 40k	0.196	0.041	0.743	0.942
OT-CFM	0.149	0.020	0.829	0.944
SB-CFM	0.150	0.020	0.825	0.942
Persistence	0.316	0.077	0.442	0.775
Location–scale	0.144	0.019	0.821	0.927

Table S13 (continued)

Predictor	TV	JS	HVG rho	Signature rho
Mean translation	0.148	0.019	0.822	0.918
Observed persistence	0.341	0.092	NA	NA
Target decoding reference	0.084	0.008	0.896	0.962
Training target marginal	0.113	0.013	0.858	0.942

Table S14 separates the missing-time and missing-donor questions. Training endpoints are paired by donor, while the individual cells drawn from those endpoints are independent. Holding out a sample in an already trained donor is not equivalent to holding out that donor. The representation is frozen before all temporal comparisons.

Table S14. Training and evaluation information for the six temporal protocols. Human data comprise three donors at Skin, Wound1, Wound7 and Wound30; mouse arms use postoperative days (POD) 0, 2, 7 and 30. Scaling follows the listed training exclusions. Source-only context at test time is permitted for donor-conditioned prediction, while the evaluation target is excluded from fitting in each held-out task. The explicit target column distinguishes prediction for an excluded donor from sensitivity analyses evaluated only in retained donors.

Protocol	Training data	Evaluation target and interpretation
Global Wound7 holdout	Skin, Wound1 and Wound30 from the included donors	Their Wound7 distributions; one globally missing measurement time.
Drop-one-donor sensitivity	Same time exclusions after removing one donor	Wound7 in the retained donors; sensitivity to donor composition, not prediction of the omitted donor.
Donor transfer	All four times in the other two donors	Omitted donor's Wound7 from their Wound1 source; observed interval in the training cohort.
Method benchmark	Same donor-transfer folds	Shared/conditioned fields, three displacement predictors and the training-target marginal; training-donor fits shown separately.
Donor-count curve	One or two of the available training donors	Same held donor and target; common scoring reference across k; seeds and subsets do not add independent donors.
Mouse POD7 audit	POD0, POD2 and POD30 in one source arm	POD7 from POD2; one animal per time point confounds time and individual.

Evaluation first resolves cell and sample identities and selects the lineage. Held-out cells are then defined before standardization is fitted to training cells alone. Endpoints sampled from those training populations supply the optimization objective. The fitted field transports the source cells with a fixed integrator, after which target, unchanged-source and split-half distances are computed. Aggregation preserves the stated donor, seed and training-subset units. Target-distance path scans are diagnostics after prediction and are not used to select the reported target coordinate.

For completeness, the fixed-step RK4 update for step h is

$$\begin{aligned} k_1 &= v(z_n, s_n), & k_2 &= v(z_n + hk_1/2, s_n + h/2), \\ k_3 &= v(z_n + hk_2/2, s_n + h/2), & k_4 &= v(z_n + hk_3, s_n + h), \\ z_{n+1} &= z_n + \frac{h}{6}(k_1 + 2k_2 + 2k_3 + k_4), & h &= (s_b - s_a)/50. \end{aligned} \tag{S1}$$

Equation (S1) gives the numerical update for the direct target predictions. The conditioned field holds the source-derived context fixed throughout this integration. Under standard smoothness assumptions RK4 has fourth-order global convergence, but this property does not establish the actual error of the fitted field or certify the chosen step count in these experiments.

The finite empirical energy distance is nonzero when two independently sampled subsets represent the same population. The split-half reference therefore indicates one realized scale of target-sampling variation. It changes with cell number, latent scaling and random split, and does not include model-fit uncertainty. Comparing a prediction gain to it is a practical diagnostic with no established frequentist size in this analysis.

Likewise, the endpoint-distance ordering condition cannot show that a point lies on a line or a geodesic. In a metric space, even being closer to both endpoints than the endpoints are to each other is only a necessary-looking geometric screen, not an interpolation identity; here the unsquared-root energy quantity is used by convention and no triangle-inequality argument is invoked. PDB is consequently labelled as failing the ordering screen, not as a proven off-path biological state.

The fitted transport moves probability mass between empirical distributions; it does not model absolute cell expansion, death, migration into a biopsy or clinical closure. These unobserved processes and the non-uniqueness of couplings prevent a mechanistic interpretation of an individual streamline.

Tables S15–S23 provide target errors, time-axis comparisons, the six-entry benchmark including stay-still, donor-specific method errors, the seed-averaged donor curve, mouse-arm contrasts and descriptive geometry intervals. The five historical pre-

dictors share test donors and are not independent replications. The added target marginal is reported in Table S35, with matched prediction counts and identical target samples; it supplies no new test donor and does not establish a population ranking.

For the revised count curve, a model first returns its prediction from its training-standardized space to the frozen encoder coordinates. A separate reference estimated from both non-held donors maps every prediction, source and target in that fold to the same scoring coordinates. The reference is used only for retrospective evaluation, not to fit the single-donor models. All 27 shared-field fits retain 8,000 optimization steps, 256-cell batches and 50 RK4 steps, with endpoint and network random generators reset for each configuration. Full target-marginal prediction uses all permitted Wound7 cells with equal donor mass; its matched version averages 50 samples at the held-source particle count.

Table S36 compares the nine saved two-training-donor fields after changing only the prediction source. For each held donor, the true Wound1 source, 50 equal-donor draws from training Wound1 and 50 existing training Wound7 marginal draws share the same target sample and reference coordinates. Source substitution integrates all permitted training Wound1 cells once per field and selects matched-size draws from those saved predictions. Draws and seeds quantify computational and cell-sampling sensitivity in three people, not independent validation. The new fields belong to the common-reference donor-curve experiment; they are not the historical single-seed method-benchmark fits in Table S35.

In the donor curve, two alternative single-donor fits contribute to each $k = 1$ donor/seed value. The three-seed mean improvement is then aggregated over three held donors. Because training sets overlap between folds, the block interval does not estimate calibrated population or predictive uncertainty. Extending the curve requires new independent donors before any plateau rule can be evaluated. There is no statistically identified minimum cohort size from the two observed k values. Table S24 reports all six training-donor comparisons. Tables S25 and S26 separate mean-displacement controls, seed-level means and configuration-specific training variability. Table S21 lists the cell counts behind each mouse computation; each cell group still represents only one animal. The retained-donor sensitivity rows in Table S15 evaluate the two donors left in the fit, not the excluded donor.

Table S15. Human Wound7 evaluation in GSE241132. ED denotes the empirical energy-distance V-statistic in training-standardized latent space; lower values indicate closer predicted and observed distributions. The pooled row excludes Wound7 globally. Drop-one rows exclude the named donor and evaluate pooled Wound7 only in the two retained donors. Transfer rows instead evaluate the excluded donor after fitting all four time points of the other donors. Improvement is 100 times (stay-still ED minus predicted ED) divided by stay-still ED. Split-half ED uses one seeded target split and is not a confidence limit. Every row is a point estimate; these dependent fits do not create additional donors.

Evaluation	Stay-still ED	Predicted ED	Split-half ED	Improvement
Pooled time holdout	1.4069	0.7054	0.0043	49.86%
Drop PWH26; retained pair	1.3357	0.7434	0.0043	44.34%
Drop PWH27; retained pair	1.2780	0.6258	0.0128	51.04%
Drop PWH28; retained pair	1.5460	0.7638	0.0080	50.59%
PWH26 transfer	1.8221	0.7110	0.0171	60.98%
PWH27 transfer	1.5355	0.2786	0.0096	81.85%
PWH28 transfer	1.0704	0.1634	0.0418	84.73%

Table S16. Historical single-seed four-clock diagnostic under global Wound7 holdout; not the revised paired exact-time comparison. The reference distance is 1.3879 and the split-half distance is 0.0061. ED denotes empirical energy distance in training-standardized latent space. Wound7 time is the global transformed coordinate; target distance uses the nearest point of the 60-step Wound1-to-Wound30 scan. Improvement is 100 times gain divided by stay-still distance. The three donors are reused, endpoint draws were not paired across clocks, and the path minimum is not substituted for target error. No interval is shown.

Time coordinate	Wound7 time	Target ED	Improvement	Diverged
rank	0.6667	0.6994	49.61%	False
days	0.2333	1.6411	-18.24%	False
sqrt days	0.4830	0.7695	44.56%	False
log days	0.6055	0.6344	54.29%	False

Table S17. Paired exact-time clock audit: four transforms, three seeds, 12 fields, three healthy donors. Each fit uses 8,000 steps, batch 256, a training-only scaler excluding all day-7 cells, and common source/target identities capped at 1,200 per donor. ED is energy distance in the common standardized encoder coordinates. Means equally weight donors and seeds. The seed range covers three equal-donor means, not a biological confidence interval. The centroid comparator uses the same donor's day-1/day-30 means with the relevant time fraction. The final column is the maximum absolute donor/seed ED change from 50 to 200 RK4 steps, not proof of solver convergence. CPU checkpoint reload reproduces the predictions; no clock is selected using the held target.

Clock	Flow ED	Flow seed range	Centroid ED	Unchanged ED	Max refinement change
rank	0.8104	0.8037 to 0.8229	0.6667	1.4746	4.98e-08
days	2.1542	1.9740 to 2.3100	0.9775	1.4746	2.13e-06
sqrt days	0.8746	0.8588 to 0.8924	0.7476	1.4746	2.78e-08
log days	0.7725	0.7400 to 0.8120	0.6654	1.4746	7.57e-08

Table S18. Common Wound1-to-Wound7 benchmark across the three held-out human donors. ED is empirical energy distance in each training-standardized fold; reported means equally weight donors. Gain is stay-still ED minus method ED. Pass count records gains above the single split-half target reference, a descriptive criterion rather than a significance test. CFM denotes conditional flow matching and OT the entropic optimal-transport displacement extension. All methods share test donors; point estimates without confidence intervals do not establish a population ranking.

Method	Mean ED	Mean gain	Pass count
Stay-still	1.4760	0.0000	0/3
Shared CFM	0.3947	1.0813	3/3
Mean displacement	0.4186	1.0574	3/3
Nearest donor	0.5147	0.9613	3/3
Conditioned CFM	0.6176	0.8584	3/3
Entropic OT extension	0.7284	0.7476	3/3

Table S19. Empirical energy distances for each method and held-out human donor in the common benchmark. Each column uses the other two donors for fitting and scaling, the named donor's Wound1 cells as source and its Wound7 cells only for evaluation. Lower is better; stay-still retains the source distribution. CFM means conditional flow matching; OT is the entropic transport displacement extension. Entries are point estimates with a 1,200-cell distance cap. Training draws and this cap differ from separately run transfer experiments, so those point estimates need not coincide.

Method	PWH26	PWH27	PWH28
Stay-still	1.8221	1.5355	1.0704
Shared CFM	0.7110	0.2881	0.1850
Mean displacement	0.5674	0.3496	0.3388
Nearest donor	0.6792	0.4318	0.4331
Conditioned CFM	1.4730	0.1990	0.1809
Entropic OT extension	0.7201	0.6228	0.8423

Table S20. Seed-averaged donor-count comparison after returning each prediction to the frozen encoder space and scoring in a common reference within held donor. The reference uses both non-held donors and never enters a single-donor fit. ED is empirical energy distance; k is the number of training donors. Single-donor choices are averaged within held donor and seed, followed by seeds 0, 1 and 2. Reduction is k=1 minus k=2. The donor-balanced reduction is 0.1483. The interval [0.0732, 0.2727] is the minimum and maximum of the three donor reductions; with three blocks a percentile bootstrap returns those endpoints, so it is not a calibrated 95% population interval. Overlapping training sets add a further reason not to read it as population coverage.

Held donor	k=1 ED	k=2 ED	Reduction
PWH26	1.0313	0.7586	0.2727
PWH27	0.3359	0.2368	0.0991
PWH28	0.2726	0.1994	0.0732

Table S21. Fibroblast cell counts and target-sampling reference for each mouse arm. POD means postoperative day. POD0, POD2 and POD30 enter training; POD7 is the held-out target. Every arm/timepoint cell group comes from one animal, so these counts quantify distribution sampling and not biological replication. The final column is one seeded split-half POD7 energy distance in the training-standardized space. NDB, PDB and GDB are retained source-study arm labels. All arms share 4,182/7,002 mapped panel genes.

Arm	POD0	POD2	POD7 held out	POD30	Split-half ED
NDB	2818	8177	12914	550	0.00544
PDB	1197	471	2500	813	0.01091
GDB	223	2400	11590	1402	0.00617

Table S22. Mouse POD7 audit for the source-labelled NDB, PDB and GDB arms. Each field uses rank time and POD0, POD2 and POD30 for fitting/scaling, then predicts from POD2 to excluded POD7. ED is empirical energy distance; improvement is 100 times (stay-still ED minus predicted ED) divided by stay-still ED. Each arm covers 4,182/7,002 human-panel genes and has one animal per time point. The ordering check asks whether endpoint distance is at least both endpoint-to-target distances; it is not proof of a geodesic. Values have no confidence intervals, and cell counts do not provide animal replication.

Arm	Stay-still ED	Predicted ED	Improvement	Ordering check
NDB	1.2341	1.2525	-1.49%	Yes
PDB	2.4345	1.6526	32.12%	No
GDB	1.3532	0.6682	50.62%	Yes

Table S23. Descriptive geometry for three healthy acute-wound donors. The first estimate divides the median of 18 within-donor between-time energy distances by the median of 12 between-donor same-time distances. The second is the mean of three pairwise cosines between donor Wound1-to-Wound7 mean-displacement vectors. Endpoints are percentile 95% limits from 10,000 resamples of pairwise distances within class or of cosine pairs. Pairs share donors, so these intervals do not provide independent donor replication or a general population ordering.

Summary	Estimate	Lower endpoint	Upper endpoint
time to donor median ratio	8.4570	6.0173	13.1114
mean displacement cosine	0.9477	0.9432	0.9536

Table S24. All six training-donor comparisons for shared and donor-conditioned flow matching. The first column identifies the donor excluded from that fit; the second is a retained training donor whose Wound1-to-Wound7 prediction is evaluated. ED is empirical energy distance in the fold-standardized coordinates. Both source and target of the evaluated donor were available during fitting, so these are fitting-capacity comparisons rather than donor-transfer tests. Conditioned errors are lower in all six rows. The repeated use of three donors prevents interpreting the rows as six independent biological replicates.

Excluded donor	Evaluated training donor	Stay-still ED	Shared ED	Conditioned ED
PWH26	PWH27	1.6257	0.1277	0.0369
PWH26	PWH28	1.0837	0.0930	0.0305
PWH27	PWH26	1.7375	0.2483	0.0511
PWH27	PWH28	1.0227	0.1248	0.0286
PWH28	PWH26	1.7873	0.3089	0.0584
PWH28	PWH27	1.6191	0.1190	0.0306

Table S25. Donor-count means in common reference coordinates. CFM denotes conditional flow matching; ED is empirical energy distance. The three CFM seeds reset both endpoint and network random generators. Deterministic controls are listed once. Full target-marginal prediction assigns equal mass to each selected training donor; matched prediction averages 50 draws with the same particle count as the held-source distribution. k=1 averages the two available single-training-donor choices before equally weighting held donors. Evaluation targets and scoring coordinates remain fixed across k and seed. Positive reductions favour k=2; these computational configurations retain three biological donors.

Predictor	Seed	k=1 mean ED	k=2 mean ED	Reduction
Shared CFM	0	0.5315	0.3854	0.1462
Mean displacement	Fixed	0.5095	0.4186	0.0909
Target marginal, full	Fixed	0.2222	0.1676	0.0546
Target marginal, matched	Fixed	0.2332	0.1818	0.0514
Unchanged source	Fixed	1.4760	1.4760	0.0000
Shared CFM	1	0.5834	0.4276	0.1558
Shared CFM	2	0.5250	0.3819	0.1430

Table S26. Training stochasticity in common reference coordinates for every shared-field donor/training-subset configuration. Mean ED, sample SD and range summarize seeds 0, 1 and 2; range is maximum minus minimum. The observed source and target indices, target split and reference scaler are fixed within held donor, so variation within each row reflects fitting randomness. Seeds and overlapping training subsets are not independent donors. The single split-half target reference measures finite-cell sampling and omits fitting variation.

Held donor	Training donors	k	Mean ED	SD	Range
PWH26	PWH27	1	1.2253	0.1920	0.3377
PWH26	PWH28	1	0.8373	0.0056	0.0097
PWH26	PWH27|PWH28	2	0.7586	0.1238	0.2334
PWH27	PWH26	1	0.1952	0.0287	0.0573
PWH27	PWH28	1	0.4766	0.1098	0.1962
PWH27	PWH26|PWH28	2	0.2368	0.0516	0.1021
PWH28	PWH26	1	0.2468	0.0351	0.0655
PWH28	PWH27	1	0.2985	0.0816	0.1633
PWH28	PWH26|PWH27	2	0.1994	0.0029	0.0057

Table S27. Fixed-field Runge–Kutta resolution comparison for pooled day-7 holdout. A single seed-0 field was refitted with the specified 8,000 optimization steps and 256-cell batches on the saved expression coordinates, excluding day 7 from fitting and scaling. All rows use the same 1,012 day-1 cells and the same 1,500 sampled target cells. Energy distance is the empirical V-statistic evaluated in double precision; improvement is relative to the same unchanged-source baseline (1.406924). RMS is the root-mean-square coordinate difference from the 200-step solution; it is a numerical discrepancy, not a confidence interval. The comparison is conditional on one field and finite-precision arithmetic and does not establish formal convergence to an exact ODE solution.

RK4 steps	Energy distance	Improvement	RMS vs 200 steps
25	0.705395409	49.862585%	5.46e-07
50	0.705395449	49.862582%	4.76e-07
100	0.705395463	49.862581%	5.36e-07
200	0.705395452	49.862582%	0

Table S28. Evaluation-sampling sensitivity for the same fixed field with 50 RK4 steps. For each cell count, 50 seeded draws sample source and target cells without replacement; prediction and unchanged-source comparisons use identical source indices and identical target indices within a draw. An additional pair of disjoint target subsets at that count defines a split-half reference. Percentages summarize relative improvement over unchanged source; the central 95% draw range measures Monte Carlo sensitivity conditional on this cohort and field, not patient-level confidence. The final column counts draws whose gain exceeds the split-half reference. Cell count and resampling do not increase the three-donor biological sample size.

Cells per distribution	Draws	Median improvement	Central 95% draw range	Gain exceeds reference
250	50	48.46%	44.24 to 51.57%	50/50
500	50	48.39%	46.22 to 51.69%	50/50
1000	50	49.26%	47.86 to 50.60%	50/50

Table S29. Evaluation of the saved pooled 50-step field in training-standardized coordinates, using all 1,012 Wound1 and 3,821 Wound7 cells. ED is the double-precision weighted empirical energy distance, without subsampling. Cell-pooled weights are uniform; donor-balanced weights allocate one third to each donor and equal mass within donor. Predictions retain source weights. The final three rows evaluate training donors separately, not donor-transfer folds. Improvement uses the corresponding unchanged-source ED. These three-donor point estimates have no confidence intervals and differ from capped-distance estimates.

Evaluation	Source / target cells	Stay-still ED	Predicted ED	Improvement
Cell-pooled	1012 / 3821	1.36146	0.68890	49.40%
Equal donor mass	1012 / 3821	1.31723	0.66265	49.69%
PWH26	255 / 920	1.75803	0.92909	47.15%
PWH27	497 / 1992	1.58457	0.84112	46.92%
PWH28	260 / 909	1.07483	0.64420	40.07%

Table S30. Observed author-annotated fibroblasts retained after the sample–barcode join in GSE241132. Donors A, B and C correspond to source donor labels PWH26, PWH27 and PWH28. Each cell-count entry comes from one biopsy; the three donors supply the biological replication. The 12 entries sum to 12,259 cells and provide the material for expression and geometry displays. Day 0 is intact skin, followed by acute wounds at days 1, 7 and 30.

Donor	Day 0 cells	Day 1 cells	Day 7 cells	Day 30 cells
Donor A	724	255	920	843
Donor B	521	497	1992	2096
Donor C	907	260	909	2335

Table S31. Retrospective global missing-time comparison in three donors. Each error is the equally weighted mean of three within-donor, full-cell empirical energy distances in all 15 training-standardized coordinates. The table gives its mean and range over seeds 0, 1 and 2; deterministic baselines have zero seed range. Improvement is relative to the corresponding unchanged-source error. Withheld day 7 uses day 1 as source and day 30 as anchor; withheld day 1 uses intact skin as source and day 7 as anchor. Both have rank-time interpolation fraction one half. The held-time cells are excluded from scaling, fitting and anchor construction. Seed ranges are not donor confidence intervals, and absolute errors from different tasks use different scalers.

Held day	Predictor	Mean ED	Minimum ED	Maximum ED	Mean improvement
7	Unchanged source	1.4725	1.4725	1.4725	0.00%
7	Centroid translation	0.6674	0.6674	0.6674	54.67%
7	Independent bridge	0.9284	0.9224	0.9386	36.95%
7	Shared flow	0.8110	0.8044	0.8237	44.93%
1	Unchanged source	1.3460	1.3460	1.3460	0.00%
1	Centroid translation	0.9293	0.9293	0.9293	30.96%
1	Independent bridge	1.0959	1.0920	1.1007	18.58%
1	Shared flow	1.0571	0.9484	1.1231	21.46%

Table S32. Every donor-specific error in the two global missing-time tasks, summarized over the three computational seeds. ED is the full 15-dimensional empirical energy distance using all available source-derived predictions and target cells in that donor. The mean, minimum and maximum describe seeds, not independent specimens. The centroid baseline has lower seed-averaged error than the field in all three day-7 donors and two of three day-1 donors. Each donor supplies both training endpoints in this retrospective design; these rows are not held-out-donor transfer.

Held day	Donor	Predictor	Mean ED	Minimum ED	Maximum ED
7	Donor A	Unchanged source	1.7580	1.7580	1.7580
7	Donor A	Centroid translation	0.7058	0.7058	0.7058
7	Donor A	Independent bridge	0.8633	0.8508	0.8712
7	Donor A	Shared flow	0.9069	0.8878	0.9291
7	Donor B	Unchanged source	1.5846	1.5846	1.5846
7	Donor B	Centroid translation	0.6898	0.6898	0.6898
7	Donor B	Independent bridge	0.9808	0.9591	0.9980
7	Donor B	Shared flow	0.8645	0.8411	0.8804
7	Donor C	Unchanged source	1.0748	1.0748	1.0748
7	Donor C	Centroid translation	0.6066	0.6066	0.6066
7	Donor C	Independent bridge	0.9411	0.8981	0.9889

Table S32 (continued)

Held day	Donor	Predictor	Mean ED	Minimum ED	Maximum ED
7	Donor C	Shared flow	0.6614	0.6442	0.6868
1	Donor A	Unchanged source	1.1341	1.1341	1.1341
1	Donor A	Centroid translation	0.8704	0.8704	0.8704
1	Donor A	Independent bridge	1.0311	1.0192	1.0471
1	Donor A	Shared flow	0.8410	0.7370	0.8960
1	Donor B	Unchanged source	1.5517	1.5517	1.5517
1	Donor B	Centroid translation	1.0900	1.0900	1.0900
1	Donor B	Independent bridge	1.3196	1.3132	1.3232
1	Donor B	Shared flow	1.3462	1.2486	1.4146
1	Donor C	Unchanged source	1.3522	1.3522	1.3522
1	Donor C	Centroid translation	0.8275	0.8275	0.8275
1	Donor C	Independent bridge	0.9369	0.9258	0.9525
1	Donor C	Shared flow	0.9841	0.8597	1.0585

Table S33. Geometry controls for globally withheld days 7 and 1. Entries average seeds within each donor and then weight the three donors equally. ED is full-cell empirical energy distance in the 15 training-standardized coordinates. Centered ED subtracts each distribution's own mean only for evaluation; it is a separate shape diagnostic and is not an additive error decomposition. Centroid error is Euclidean distance between means. Variance ratio is predicted total coordinate variance divided by observed target variance. Location–scale interpolates only the permitted endpoint means and standard deviations at rank fraction one half; no target mean or dispersion enters prediction. All entries are point estimates, not population uncertainty intervals.

Held day	Predictor	ED	Centered ED	Centroid error	Variance ratio
7	Unchanged source	1.4725	0.1495	2.5971	0.761
7	Centroid translation	0.6674	0.1495	1.5375	0.761
7	Independent bridge	0.9284	0.2691	1.5426	0.319
7	Shared flow	0.8110	0.1441	1.6835	0.543
7	Location-scale	0.6982	0.1466	1.5375	0.622
1	Unchanged source	1.3460	0.0647	2.5213	0.990
1	Centroid translation	0.9293	0.0647	2.0270	0.990
1	Independent bridge	1.0959	0.1445	2.0215	0.579
1	Shared flow	1.0571	0.1220	2.0158	0.641
1	Location-scale	0.9034	0.0667	2.0270	1.107

Table S34. Matched evaluation with 200 source and 200 target cells per donor, sampled without replacement in 50 draws with seed 29. Each draw shares indices across all predictors and fit seeds, then averages seeds within donor and donors equally. Entries are shared-flow error minus comparator error; positive differences favor the comparator. Full ranges describe observed-cell resampling sensitivity, not donor confidence intervals or hypothesis tests. The 50 draws still involve only the same three biological donors.

Held day	Comparator	Mean difference	Full draw range	Comparator lower
7	Centroid translation	0.1400	0.1132 to 0.1716	50/50
7	Location-scale	0.1102	0.0847 to 0.1405	50/50
7	Independent bridge	-0.1151	-0.1576 to -0.0741	0/50
1	Centroid translation	0.1243	0.0977 to 0.1545	50/50
1	Location-scale	0.1497	0.1194 to 0.1763	50/50
1	Independent bridge	-0.0407	-0.0833 to -0.0070	0/50

Table S35. Training-target marginal predictor compared with the historical shared-field benchmark. Each target-marginal draw samples the permitted training donors' Wound7 cells without replacement, matches the original held-source particle count (255, 497 or 260), and assigns each training donor exactly half of total probability mass. It uses no held-source expression. Scaling and target indices are identical to the original benchmark. Means and full ranges describe 50 draws; they are not donor confidence intervals. The shared-field column is its original saved error, because historical per-cell predictions were unavailable for a new paired evaluation. Across donors the matched marginal mean is 0.1818 versus 0.3947 for the field; lower is better.

Held donor	Shared-field ED	Target-marginal ED	Full draw range	Marginal lower
PWH26	0.7110	0.1569	0.1147 to 0.2106	50/50
PWH27	0.2881	0.1416	0.1055 to 0.1769	50/50
PWH28	0.1850	0.2469	0.1860 to 0.3340	0/50

Table S36. Retrospective source-replacement control for the nine saved two-training-donor fields (three held donors by three seeds). All scores are empirical energy distance in the same reference coordinates and on the same held-donor Wound7 target cells; lower is better. Real is the actual held-donor Wound1 distribution. Substituted sources are 50 without-replacement, source-count-matched draws from permitted training donors' Wound1 cells, integrated by the identical saved field, with one-half mass assigned to each training donor. The target marginal instead draws from their Wound7 cells without reading the held source; its draw IDs are not paired with substituted-source draws. Draws are averaged within field, then seeds within held donor; the final row equally weights three donors. The difference is substituted minus real, so a negative value favors the substituted input. These are retrospective, dependent computations on three healthy donors, not confidence intervals, new biological replicates or a model-selection test.

Held donor	Real-source ED	Substituted ED	Target marginal ED	Substituted minus real
PWH26	0.7586	0.2802	0.1569	-0.4785
PWH27	0.2368	0.2732	0.1416	0.0363
PWH28	0.1994	0.1816	0.2469	-0.0179
Three-donor mean	0.3983	0.2450	0.1818	-0.1533

Table S37. Exact target-marginal mixture decomposition in fixed held-fold coordinates. Separate ED is the mean of the two single-training-donor target errors; mixed ED scores their equal-mass empirical mixture. Reduction is separate minus mixed and equals one quarter of their mutual ED, independently of the held target. ED denotes the empirical energy discrepancy without its square root. Complete permitted training-Wound7 distributions and the original held-target indices reproduce the earlier full-marginal curve. Maximum identity residual is 0.0000000000000024. The final row equally weights three donors. This identity does not decompose a jointly fitted neural field, and lower mixture error does not itself identify target-relevant information gain.

Held donor	Separate ED	Mixed ED	Reduction	Mutual ED / 4
PWH26	0.202462	0.137777	0.064685	0.064685
PWH27	0.199092	0.135287	0.063805	0.063805
PWH28	0.265045	0.229730	0.035315	0.035315
Three-donor mean	0.222199	0.167598	0.054601	0.054601

Table S38. Matched-particle source substitutions in nine existing two-training-donor fields (three held donors, seeds 0–2). Each score uses 200 predicted particles: real held-source cells, one individual permitted training-donor source, or a mixture of 100 cells from each permitted donor. Fifty without-replacement draws share fixed held-target indices and reuse source indices across model seeds. Wrong A/B list actual source donors; their identities vary across held folds, so the final columns average fold-specific substitutions, not a single common donor. Scores average draws within field, then seeds within held donor, then donors equally. The individual-substitution error average is 0.3075. These are point energy discrepancies in common reference coordinates, not confidence intervals or new biological replication. Mixture identities are separately checked on the actual 100-cell components, not by equating them to the 200-cell individual distributions. No model is refitted.

Held donor	Real-source ED	Wrong A ED	Wrong B ED	Mixed ED
PWH26	0.7578	PWH27: 0.3003	PWH28: 0.3745	0.2799
PWH27	0.2488	PWH26: 0.3504	PWH28: 0.3057	0.2877
PWH28	0.2043	PWH26: 0.2503	PWH27: 0.2636	0.1904
Three-donor mean	0.4036	0.3003	0.3146	0.2527

Table S39. Observable-space errors against actual held-donor expression in the same three healthy volunteers. All 5,383 assayed frozen-panel genes are retained without a performance filter; 1,619 absent genes are not observed zeros. Each observed cell or decoded particle is normalized on the common panel before equal-cell averaging. Saved target uses the original held-target identities; all day 7 is a predeclared sensitivity. k counts permitted training donors, not test donors. Twenty-seven saved fits use three computational seeds; no model is trained. Averages weight seeds, then training subsets, then held donors equally. TV is total variation between mean profiles; JS is Jensen–Shannon divergence in natural-log units, not its square root; RMSE is the root mean square error over gene proportions. Lower is better. Observed and decoded baselines are distinct; training marginals use only permitted donors. These are mean-profile errors, not single-cell transitions, full distributional fidelity, absolute-count prediction or biological confidence intervals.

Target set	k	Predictor	TV	JS (nats)	Gene RMSE
Saved target	1	Decoded shared flow	0.2934	0.0746	0.000779
Saved target	1	Decoded persistence	0.3529	0.0990	0.001128
Saved target	1	Decoded mean displacement	0.2882	0.0717	0.000759
Saved target	1	Decoded training marginal	0.2590	0.0636	0.000600
Saved target	1	Observed persistence	0.3421	0.0928	0.001268
Saved target	1	Observed training marginal	0.1279	0.0155	0.000370
Saved target	2	Decoded shared flow	0.2823	0.0709	0.000694
Saved target	2	Decoded persistence	0.3529	0.0990	0.001128
Saved target	2	Decoded mean displacement	0.2825	0.0703	0.000714
Saved target	2	Decoded training marginal	0.2546	0.0627	0.000571
Saved target	2	Observed persistence	0.3421	0.0928	0.001268
Saved target	2	Observed training marginal	0.1115	0.0120	0.000322
All day 7	1	Decoded shared flow	0.2929	0.0744	0.000780
All day 7	1	Decoded persistence	0.3522	0.0988	0.001126
All day 7	1	Decoded mean displacement	0.2878	0.0716	0.000760
All day 7	1	Decoded training marginal	0.2584	0.0634	0.000600
All day 7	1	Observed persistence	0.3412	0.0923	0.001265
All day 7	1	Observed training marginal	0.1270	0.0153	0.000369

Table S39 (continued)

Target set	k	Predictor	TV	JS (nats)	Gene RMSE
All day 7	2	Decoded shared flow	0.2818	0.0707	0.000695
All day 7	2	Decoded persistence	0.3522	0.0988	0.001126
All day 7	2	Decoded mean displacement	0.2820	0.0702	0.000714
All day 7	2	Decoded training marginal	0.2541	0.0624	0.000571
All day 7	2	Observed persistence	0.3412	0.0923	0.001265
All day 7	2	Observed training marginal	0.1103	0.0118	0.000320

Table S40. Decoder reconstruction reference for the observed held-donor day-7 cells. The frozen encoder means of the target cells themselves are decoded, restricted to the same 5,383 observed panel genes and normalized per cell before averaging. Errors compare this reconstruction with actual mean observed expression, not with another decoded target. TV is total variation; JS is Jensen–Shannon divergence in nats. The missing-mass column reports mean decoder probability on the 1,619 unmeasured panel genes before renormalization. This reference uses held-target information: it is not a forecast, true expression, a competing predictor or a mathematical lower error bound. The two target sets overlap and do not add independent observations.

Held donor	Target set	Cells	TV	JS (nats)	Unmeasured mass
PWH26	Saved target	920	0.2314	0.0553	1.59%
PWH26	All day 7	920	0.2314	0.0553	1.59%
PWH27	Saved target	1200	0.2549	0.0655	1.62%
PWH27	All day 7	1992	0.2537	0.0651	1.62%
PWH28	Saved target	909	0.2426	0.0584	1.79%
PWH28	All day 7	909	0.2426	0.0584	1.79%

Table S41. Saved finite-budget diagnostic: flow minus the fixed seed-0 training-target draw in each donor fold. Positive differences indicate higher error for the flow. Each cell equally averages three observed-source adjacent intervals and two computational seeds within that donor and budget. Training-only HVG200/PCA15 coordinates, 64-unit two-layer fields, time embedding 16 and batch size 64 differ from the primary frozen neural-coordinate experiment. Equal steps do not imply equal wall time. Fold-specific PCA scales are not pooled into a cross-donor effect, and the baseline sampling is not symmetric with two flow seeds. OT is minibatch optimal-transport coupling; SB is the entropic bridge variant. None of the four flows has lower within-donor mean energy at either budget, but individual intervals may favor flow. No p values, population ranking or new donor replication are claimed.

Held donor	Steps	Independent	Minibatch OT	SB	Conditioned
PWH26	120	0.7096	0.6888	0.7001	0.1986
PWH26	600	0.1630	0.1619	0.1454	0.1843
PWH27	120	1.0705	1.0422	1.0655	0.3248
PWH27	600	0.2432	0.2359	0.2263	0.2379
PWH28	120	0.7610	0.7482	0.7560	0.3122
PWH28	600	0.4134	0.3517	0.3464	0.3553

Table S42. Complete saved within-donor method/budget means behind the diagnostic. Energy is the capped empirical V-distance in that fold's training-only PCA coordinates (200 cells per distribution); lower values are better within a fold. Flow rows average three time pairs and two seeds (six records); baseline rows average three pairs at seed 0 (three records). These are 48 flow fits and 12 baseline jobs with 180 dependent predictions from three donors, not 180 biological replicates. All pairs start from their observed source rather than an autonomous day-0 rollout. SB retained all 4,320 training batches, including 20 failing the frozen two-sided marginal L1 tolerance of 0.001; maximum row and column residuals were 0.002061 and 0.00001014. Prediction-plus-target adaptive MMD is diagnostic only and is not used to rank methods. The historical diagnostic saved predictions, not reloadable fields.

Held donor	Method	Steps	Records	Mean Energy
PWH26	Independent	120	6	0.9513
PWH26	Independent	600	6	0.4047
PWH26	Minibatch OT	120	6	0.9305
PWH26	Minibatch OT	600	6	0.4035
PWH26	SB	120	6	0.9418
PWH26	SB	600	6	0.3871
PWH26	Conditioned	120	6	0.4403
PWH26	Conditioned	600	6	0.4260
PWH26	Persistence	None	3	1.5159
PWH26	Mean shift	None	3	0.3842
PWH26	Location scale	None	3	0.3767
PWH26	Target draw	None	3	0.2417
PWH27	Independent	120	6	1.3054
PWH27	Independent	600	6	0.4781
PWH27	Minibatch OT	120	6	1.2771
PWH27	Minibatch OT	600	6	0.4708
PWH27	SB	120	6	1.3004
PWH27	SB	600	6	0.4612
PWH27	Conditioned	120	6	0.5598

Table S42 (continued)

Held donor	Method	Steps	Records	Mean Energy
PWH27	Conditioned	600	6	0.4728
PWH27	Persistence	None	3	1.8764
PWH27	Mean shift	None	3	0.4724
PWH27	Location scale	None	3	0.4587
PWH27	Target draw	None	3	0.2349
PWH28	Independent	120	6	1.0853
PWH28	Independent	600	6	0.7377
PWH28	Minibatch OT	120	6	1.0725
PWH28	Minibatch OT	600	6	0.6760
PWH28	SB	120	6	1.0803
PWH28	SB	600	6	0.6706
PWH28	Conditioned	120	6	0.6365
PWH28	Conditioned	600	6	0.6795
PWH28	Persistence	None	3	1.6663
PWH28	Mean shift	None	3	0.6217
PWH28	Location scale	None	3	0.6146
PWH28	Target draw	None	3	0.3243

Additional supporting displays




Figure S1. Donor count, source substitution and observed-expression restrictions. (a) Three-seed shared-flow curves alongside saved mean-displacement and exact target-marginal curves, in the same reference geometry and hierarchical donor weighting. (b) Macro seed reductions (navy diamonds; navy line, their mean), all nine donor/seed reductions and the saved descriptive three-block 95% range; shading is not a seed-level interval. (c) Donor means with finer seed-resolved curves. (d) Real source, each alternative donor and their mixture under the saved 200-particle, 50-draw protocol; large points are donor means and smaller points are the three seed means. Other A/B are sorted alternatives within each fold. (e) Total variation against observed mean expression on 5,383 common genes. (f) Jensen–Shannon divergence for the same predictions. e/f distinguish decoded and directly observed predictors; large symbols are equal-donor macros and faint symbols are three donor means after seed/subset averaging, for each training size. Source mixing is not a donor-count experiment, and mean-expression errors are not full cellular-distribution validation. All panels reuse three donors; seeds, draws and overlapping folds add no biological replication.

References

[1] Zhuang Liu, Xiaowei Bian, Lihua Luo, Åsa K. Björklund, Li Li, Letian Zhang, Yongjian Chen, Lei Guo, Juan Gao, Chunyan Cao, Jiating Wang, Wenjun He, Yunting Xiao, Liping Zhu, Karl Annusver, Nusayhah Hudaa Gopee, Daniela Basurto-Lozada, David Horsfall, Clare L.

Bennett, Maria Kasper, Muzlifah Haniffa, Pehr Sommar, Dongqing Li, and Ning Xu Landén. Spatiotemporal single-cell roadmap of human skin wound healing. Cell Stem Cell, 32(3):479–498.e8, 2025. doi: 10.1016/j.stem.2024.11.013.

[2] David R. Roberts, Volker Bahn, Simone Ciuti, Mark S. Boyce, Jane Elith, Gurutzeta Guillera-Arroita, Severin Hauenstein, José J. Lahoz-Monfort, Boris Schröder, Wilfried Thuiller, David I. Warton, Brendan A. Wintle, Florian Hartig, and Carsten F. Dormann. Cross-validation strategies for data with temporal, spatial, hierarchical, or phylogenetic structure. Ecography, 40(8):913–929, 2017. doi: 10.1111/ecog.02881.

[3] Geoffrey Schiebinger, Jian Shu, Marcin Tabaka, Brian Cleary, Vidya Subramanian, Aryeh Solomon, Joshua Gould, Siyan Liu, Stacie Lin, Peter Berube, Lia Lee, Jenny Chen, Justin Brumbaugh, Philippe Rigollet, Konrad Hochedlinger, Rudolf Jaenisch, Aviv Regev, and Eric S. Lander. Optimal-Transport Analysis of Single-Cell Gene Expression Identifies Developmental Trajectories in Reprogramming. Cell, 176 (4):928–943.e22, 2019. doi: 10.1016/j.cell.2019.01.006.

[4] Alexander Tong, Jessie Huang, Guy Wolf, David van Dijk, and Smita Krishnaswamy. TrajectoryNet: A Dynamic Optimal Transport Network for Modeling Cellular Dynamics. In Proceedings of the 37th International Conference on Machine Learning, volume 119 of Proceedings of Machine Learning Research, pages 9526–9536, 2020. URL https://proceedings.mlr.press/v119/tong20a.html.

[5] Gavin C. Cawley and Nicola L. C. Talbot. On Over-fitting in Model Selection and Subsequent Selection Bias in Performance Evaluation. Journal of Machine Learning Research, 11:2079–2107, 2010. URL https://www.jmlr.org/papers/v11/cawley10a.html.

[6] Haiyang Huang, Yingfan Wang, Cynthia Rudin, and Edward P. Browne. Towards a comprehensive evaluation of dimension reduction methods for transcriptomic data visualization. Communications Biology, 5:719, 2022. doi: 10.1038/s42003-022-03628-x.

[7] Caleb Weinreb, Samuel Wolock, Betsabeh K. Tusi, Merav Socolovsky, and Allon M. Klein. Fundamental limits on dynamic inference from single-cell snapshots. Proceedings of the National Academy of Sciences, 115(10):E2467–E2476, 2018. doi: 10.1073/pnas.1714723115.

[8] Hugo Lavenant, Stephen Zhang, Young-Heon Kim, and Geoffrey Schiebinger. Toward a mathematical theory of trajectory inference. The Annals of Applied Probability, 34(1A), 2024. doi: 10.1214/23-AAP1969.

[9] Caleb Weinreb, Alejo Rodriguez-Fraticelli, Fernando D. Camargo, and Allon M. Klein. Lineage tracing on transcriptional landscapes links state to fate during differentiation. Science, 367(6479):eaaw3381, 2020. doi: 10.1126/science.aaw3381.

[10] Gioele La Manno, Ruslan Soldatov, Amit Zeisel, Emelie Braun, Hannah Hochgerner, Viktor Petukhov, Katja Lidschreiber, Maria E. Kastriti, Peter Lönnerberg, Alessandro Furlan, Jean Fan, Lars E. Borm, Zehua Liu, David van Bruggen, Jimin Guo, Xiaoling He, Roger Barker, Erik Sundström, Gonçalo Castelo-Branco, Patrick Cramer, Igor Adameyko, Sten Linnarsson, and Peter V. Kharchenko. RNA velocity of single cells. Nature, 560(7719):494–498, 2018. doi: 10.1038/s41586-018-0414-6.

[11] Volker Bergen, Marius Lange, Stefan Peidli, F. Alexander Wolf, and Fabian J. Theis. Generalizing RNA velocity to transient cell states through dynamical modeling. Nature Biotechnology, 38(12):1408–1414, 2020. doi: 10.1038/s41587-020-0591-3.

[12] Marius Lange, Volker Bergen, Michal Klein, Manu Setty, Bernhard Reuter, Mostafa Bakhti, Heiko Lickert, Meshal Ansari, Janine Schniering, Herbert B. Schiller, Dana Pe'er, and Fabian J. Theis. CellRank for directed single-cell fate mapping. Nature Methods, 19(2):159–170, 2022. doi: 10.1038/s41592-021-01346-6.

[13] Michael E. Vinyard, Anders W. Rasmussen, Ruitong Li, Allon M. Klein, Gad Getz, and Luca Pinello. Learning cell dynamics with neural differential equations. Nature Machine Intelligence, 7(12):1969–1984, 2025. doi: 10.1038/s42256-025-01150-3.

[14] Emanuel Rognoni, Angela Oliveira Pisco, Toru Hiratsuka, Kalle H Sipilä, Julio M Belmonte, Seyedeh Atefeh Mobasseri, Christina Philippeos, Rui Dilão, and Fiona M Watt. Fibroblast state switching orchestrates dermal maturation and wound healing. Molecular Systems Biology, 14(8):e8174, 2018. doi: 10.15252/msb.20178174.

[15] Brett A. Shook, Renee R. Wasko, Guillermo C. Rivera-Gonzalez, Emilio Salazar-Gatzimas, Francesc López-Giráldez, Biraja C. Dash, Andrés R. Muñoz-Rojas, Krystal D. Aultman, Rachel K. Zwick, Vivian Lei, Jack L. Arbiser, Kathryn Miller-Jensen, Damon A. Clark, Henry C. Hsia, and Valerie Horsley. Myofibroblast proliferation and heterogeneity are supported by macrophages during skin repair. Science, 362(6417):eaar2971, 2018. doi: 10.1126/science.aar2971.

[16] Sudhir Varma and Richard Simon. Bias in error estimation when using cross-validation for model selection. BMC Bioinformatics, 7(1): 91, 2006. doi: 10.1186/1471-2105-7-91.

[17] Charlotte Bunne, Stefan G. Stark, Gabriele Gut, Jacobo Sarabia del Castillo, Mitch Levesque, Kjong-Van Lehmann, Lucas Pelkmans, Andreas Krause, and Gunnar Rätsch. Learning single-cell perturbation responses using neural optimal transport. Nature Methods, 20(11): 1759–1768, 2023. doi: 10.1038/s41592-023-01969-x.

[18] Gábor J. Székely and Maria L. Rizzo. Energy statistics: A class of statistics based on distances. Journal of Statistical Planning and Inference, 143(8):1249–1272, 2013. doi: 10.1016/j.jspi.2013.03.018.

[19] Dino Sejdinovic, Bharath Sriperumbudur, Arthur Gretton, and Kenji Fukumizu. Equivalence of distance-based and RKHS-based statistics in hypothesis testing. The Annals of Statistics, 41(5), 2013. doi: 10.1214/13-aos1140.

[20] Malte D. Luecken, M. Büttner, K. Chaichoompu, A. Danese, M. Interlandi, M. F. Mueller, D. C. Strobl, L. Zappia, M. Dugas, M. Colomé-Tatché, and Fabian J. Theis. Benchmarking atlas-level data integration in single-cell genomics. Nature Methods, 19(1):41–50, 2022. doi: 10.1038/s41592-021-01336-8.

[21] Kip D. Zimmerman, Mark A. Espeland, and Carl D. Langefeld. A practical solution to pseudoreplication bias in single-cell studies. Nature Communications, 12(1):738, 2021. doi: 10.1038/s41467-021-21038-1.

[22] Ernst Hairer, Syvert P. Nørsett, and Gerhard Wanner. Solving Ordinary Differential Equations I: Nonstiff Problems. Springer, 2 edition, 1993. doi: 10.1007/978-3-540-78862-1.

[23] Bradley Efron and Robert J. Tibshirani. An Introduction to the Bootstrap. Chapman and Hall, New York, 1993. ISBN 9780412042317. doi: 10.1007/978-1-4899-4541-9.

[24] Yoshua Bengio and Yves Grandvalet. No Unbiased Estimator of the Variance of K-Fold Cross-Validation. Journal of Machine Learning Research, 5:1089–1105, 2004. URL https://www.jmlr.org/papers/v5/grandvalet04a.html.

[25] Andrius Vabalas, Emma Gowen, Ellen Poliakoff, and Alexander J. Casson. Machine learning algorithm validation with a limited sample size. PLOS ONE, 14(11):e0224365, 2019. doi: 10.1371/journal.pone.0224365.

[26] David G. Armstrong, Andrew J.M. Boulton, and Sicco A. Bus. Diabetic Foot Ulcers and Their Recurrence. New England Journal of Medicine, 376(24):2367–2375, 2017. doi: 10.1056/nejmra1615439.