

# Fenchel-type Representations and Large-scale Problems with Convex Structure on Difficult Geometry Domains

Arkadi Nemirovski

H. Milton Stewart School of Industrial and Systems Engineering  
Georgia Institute of Technology

*Joint research with Bruce Cox<sup>†</sup> & Anatoli Juditsky<sup>‡</sup>*

<sup>†</sup>US AirForce <sup>‡</sup>University J. Fourier, Grenoble

Optimization and Big Data 2015  
Edinburgh May 2015

**Problem of Interest:** large-scale problems with convex structure:

- **C**onvex **M**inimization:  $\min_{x \in X} f(x)$ 
  - $X$ : convex compact set in Euclidean space
  - $f : X \rightarrow \mathbb{R}$ : convex & Lipschitz
- Convex-Concave **S**addle **P**oint:  $\min_{u \in U} \max_{v \in V} f(u, v)$
- **V**ariational **I**nequality with monotone operator:  
Find  $x \in X : \langle F(y), y - x \rangle \geq 0 \forall y \in X$

**Note:** **CM** and **SP** reduce to **VI**

- **CM**  $\Rightarrow$  **VI**:  $F$ : bounded section of  $x \mapsto \partial_x f$
- **SP**  $\Rightarrow$  **VI**:  $X := U \times V$ ,  $F$ : bounded section of  
 $[u; v] \mapsto [\partial_u f(u, v); \partial_v [-f(u, v)]]$

$\Rightarrow$  Every problem with convex structure induces a monotone vector field on problem's domain.

**Problem of Interest:** large-scale problems with convex structure:

- **Convex Minimization:**  $\min_{x \in X} f(x)$
- **Convex-Concave Saddle Point:**  $\min_{u \in U} \max_{v \in V} f(u, v)$ 
  - $U, V$ : convex compact sets in Euclidean spaces
  - $f(\cdot, \cdot) : X \rightarrow \mathbb{R}$ : convex-concave & Lipschitz
- **Variational Inequality with monotone operator:**  
Find  $x \in X : \langle F(y), y - x \rangle \geq 0 \forall y \in X$

**Note:** **CM** and **SP** reduce to **VI**

- **CM**  $\Rightarrow$  **VI**:  $F$ : bounded section of  $x \mapsto \partial_x f$
- **SP**  $\Rightarrow$  **VI**:  $X := U \times V$ ,  $F$ : bounded section of  
 $[u; v] \mapsto [\partial_u f(u, v); \partial_v [-f(u, v)]]$

$\Rightarrow$  Every problem with convex structure induces a monotone vector field on problem's domain.

**Problem of Interest:** large-scale problems with convex structure:

- **C**onvex **M**inimization:  $\min_{x \in X} f(x)$
- Convex-Concave **S**addle **P**oint:  $\min_{u \in U} \max_{v \in V} f(u, v)$
- **V**ariational **I**nequality with monotone operator:  
 Find  $x \in X : \langle F(y), y - x \rangle \geq 0 \forall y \in X$ 
  - $X$ : convex compact set in Euclidean space  $E$
  - $F : X \rightarrow E$ : monotone & bounded

**Note:** **CM** and **SP** reduce to **VI**

- **CM**  $\Rightarrow$  **VI**:  $F$ : bounded section of  $x \mapsto \partial_x f$
- **SP**  $\Rightarrow$  **VI**:  $X := U \times V$ ,  $F$ : bounded section of  
 $[u; v] \mapsto [\partial_u f(u, v); \partial_v [-f(u, v)]]$

$\Rightarrow$  Every problem with convex structure induces a monotone vector field on problem's domain.

**Problem of Interest:** large-scale problems with convex structure:

- **Convex Minimization:**  $\min_{x \in X} f(x)$
- **Convex-Concave Saddle Point:**  $\min_{u \in U} \max_{v \in V} f(u, v)$
- **Variational Inequality** with monotone operator:  
Find  $x \in X : \langle F(y), y - x \rangle \geq 0 \forall y \in X$

**Note:** **CM** and **SP** reduce to **VI**

- **CM**  $\Rightarrow$  **VI**:  $F$ : bounded section of  $x \mapsto \partial_x f$
- **SP**  $\Rightarrow$  **VI**:  $X := U \times V$ ,  $F$ : bounded section of  
 $[u; v] \mapsto [\partial_u f(u, v); \partial_v [-f(u, v)]]$

$\Rightarrow$  Every problem with convex structure induces a monotone vector field on problem's domain.

**Problem of Interest:** large-scale problems with convex structure:

- **Convex Minimization:**  $\min_{x \in X} f(x)$
- **Convex-Concave Saddle Point:**  $\min_{u \in U} \max_{v \in V} f(u, v)$
- **Variational Inequality** with monotone operator:  
Find  $x \in X : \langle F(y), y - x \rangle \geq 0 \forall y \in X$

**Note:** **CM** and **SP** reduce to **VI**

- **CM**  $\Rightarrow$  **VI**:  $F$ : bounded section of  $x \mapsto \partial_x f$
- **SP**  $\Rightarrow$  **VI**:  $X := U \times V$ ,  $F$ : bounded section of  
 $[u; v] \mapsto [\partial_u f(u, v); \partial_v [-f(u, v)]]$

$\Rightarrow$  Every problem with convex structure induces a monotone vector field on problem's domain.

**Problem of Interest:** large-scale problems with convex structure:

- **Convex Minimization:**  $\min_{x \in X} f(x)$
- **Convex-Concave Saddle Point:**  $\min_{u \in U} \max_{v \in V} f(u, v)$
- **Variational Inequality** with monotone operator:  
Find  $x \in X : \langle F(y), y - x \rangle \geq 0 \forall y \in X$

**Note:** **CM** and **SP** reduce to **VI**

- **CM**  $\Rightarrow$  **VI**:  $F$ : bounded section of  $x \mapsto \partial_x f$
- **SP**  $\Rightarrow$  **VI**:  $X := U \times V$ ,  $F$ : bounded section of  
 $[u; v] \mapsto [\partial_u f(u, v); \partial_v [-f(u, v)]]$

$\Rightarrow$  *Every problem with convex structure induces a monotone vector field on problem's domain.*

**Fact:** At present, main tools of large-scale convex optimization are proximal First Order Methods where a linear perturbation of fixed *strongly convex* function is minimized over problem's domain  $X$  at every step.

⇒ Proximal FOMs require  $X$  to be *proximal friendly* – to allow for cheap minimization of all linear perturbations of a strongly convex function. In particular,  $X$  should admit a cheap Linear Minimization Oracle – a routine capable to minimize over  $X$  linear forms.

**Fact:** Some important high-dimensional domains admit cheap LMO and are *not* proximal friendly:

- nuclear/trace norm balls [low rank matrix recovery & SDP]
- total variation ball [imaging]
- some combinatorial polytopes

⇒ **Challenge:** *Design of LMO based First Order Methods*



**Fact:** At present, main tools of large-scale convex optimization are proximal First Order Methods where a linear perturbation of fixed *strongly convex* function is minimized over problem's domain  $X$  at every step.

⇒ Proximal FOMs require  $X$  to be *proximal friendly* – to allow for cheap minimization of all linear perturbations of a strongly convex function. In particular,  $X$  should admit a cheap Linear Minimization Oracle – a routine capable to minimize over  $X$  linear forms.

**Fact:** Some important high-dimensional domains admit cheap LMO and are *not* proximal friendly:

- nuclear/trace norm balls [low rank matrix recovery & SDP]
- total variation ball [imaging]
- some combinatorial polytopes

⇒ **Challenge:** *Design of LMO based First Order Methods*

**Fact:** At present, main tools of large-scale convex optimization are proximal First Order Methods where a linear perturbation of fixed *strongly convex* function is minimized over problem's domain  $X$  at every step.

⇒ Proximal FOMs require  $X$  to be *proximal friendly* – to allow for cheap minimization of all linear perturbations of a strongly convex function. In particular,  $X$  should admit a cheap Linear Minimization Oracle – a routine capable to minimize over  $X$  linear forms.

**Fact:** Some important high-dimensional domains admit cheap LMO and are *not* proximal friendly:

- nuclear/trace norm balls [low rank matrix recovery & SDP]
- total variation ball [imaging]
- some combinatorial polytopes

⇒ **Challenge:** Design of LMO based First Order Methods

**Fact:** At present, main tools of large-scale convex optimization are proximal First Order Methods where a linear perturbation of fixed *strongly convex* function is minimized over problem's domain  $X$  at every step.

⇒ Proximal FOMs require  $X$  to be *proximal friendly* – to allow for cheap minimization of all linear perturbations of a strongly convex function. In particular,  $X$  should admit a cheap Linear Minimization Oracle – a routine capable to minimize over  $X$  linear forms.

**Fact:** Some important high-dimensional domains admit cheap LMO and are *not* proximal friendly:

- nuclear/trace norm balls [low rank matrix recovery & SDP]
- total variation ball [imaging]
- some combinatorial polytopes

⇒ **Challenge:** *Design of LMO based First Order Methods*

**Fact:** At present, main tools of large-scale convex optimization are proximal First Order Methods where a linear perturbation of fixed *strongly convex* function is minimized over problem's domain  $X$  at every step.

⇒ Proximal FOMs require  $X$  to be *proximal friendly* – to allow for cheap minimization of all linear perturbations of a strongly convex function. In particular,  $X$  should admit a cheap Linear Minimization Oracle – a routine capable to minimize over  $X$  linear forms.

**Fact:** Some important high-dimensional domains admit cheap LMO and are *not* proximal friendly:

- nuclear/trace norm balls [low rank matrix recovery & SDP]
- total variation ball [imaging]
- some combinatorial polytopes

⇒ **Challenge:** *Design of LMO based First Order Methods*

**Fact:** The only traditional optimization technique capable to solve problems with convex structure on **LMO** represented domains is the classical **Conditional Gradient** algorithm (Frank & Wolfe, 1958).

**But:** The *only* problem with convex structure in the scope of **CG** is *smooth CM*. There are *no* traditional LMO-based **FOMs** for (smooth and nonsmooth alike) **SP** and **VI** and *nonsmooth CM* problems.

**Goal of the talk:** **LMO** based **FOMs** for **SP/VI/nonsmooth CM**.

**Fact:** The only traditional optimization technique capable to solve problems with convex structure on **LMO** represented domains is the classical **Conditional Gradient** algorithm (Frank & Wolfe, 1958).

**But:** The *only* problem with convex structure in the scope of **CG** is *smooth CM*. There are *no* traditional LMO-based **FOMs** for (smooth and nonsmooth alike) **SP** and **VI** and *nonsmooth CM* problems.

**Goal of the talk:** **LMO** based **FOMs** for **SP/VI/nonsmooth CM**.

**Fact:** The only traditional optimization technique capable to solve problems with convex structure on **LMO** represented domains is the classical **Conditional Gradient** algorithm (Frank & Wolfe, 1958).

**But:** The *only* problem with convex structure in the scope of **CG** is *smooth CM*. There are *no* traditional LMO-based **FOMs** for (smooth and nonsmooth alike) **SP** and **VI** and *nonsmooth CM* problems.

**Goal of the talk:** **LMO** based **FOMs** for **SP/VI/nonsmooth CM**.

## Strategy:

- 1 Use *Fenchel-type representation* of the monotone vector field associated with the problem of interest to build the *dual* problem, which again will be a problem with convex structure.
- 2 Solve the dual problem to a desired accuracy  $\epsilon$  by a **FOM** (e.g., a proximal one, provided the domain of the dual is proximal friendly).
- 3 Utilize *accuracy certificates* to build an  $\epsilon$ -solution to the problem of interest from information acquired when processing the dual problem.



## Strategy:

- 1 Use *Fenchel-type representation* of the monotone vector field associated with the problem of interest to build the *dual* problem, which again will be a problem with convex structure.
- 2 Solve the dual problem to a desired accuracy  $\varepsilon$  by a **FOM** (e.g., a proximal one, provided the domain of the dual is proximal friendly).
- 3 Utilize *accuracy certificates* to build an  $\varepsilon$ -solution to the problem of interest from information acquired when processing the dual problem.

## Strategy:

- 1 Use *Fenchel-type representation* of the monotone vector field associated with the problem of interest to build the *dual* problem, which again will be a problem with convex structure.
- 2 Solve the dual problem to a desired accuracy  $\varepsilon$  by a **FOM** (e.g., a proximal one, provided the domain of the dual is proximal friendly).
- 3 Utilize *accuracy certificates* to build an  $\varepsilon$ -solution to the problem of interest from information acquired when processing the dual problem.

**Fenchel representation** of a function  $f : \mathbb{R}^n \rightarrow \mathbb{R} \cup \{+\infty\}$  is

$$f(x) = \sup_y \{ \langle x, y \rangle - f^*(y) \}$$

- $f^*$ : convex, proper, l.sc.

**F.r.** of a convex proper l.sc.  $f$  exists and is unique.

**However:** Fenchel representation “exists in the nature,” but, aside of a fistful of simple cases, is **not** available for numerical use.

**Fenchel-type representation** of a function  $f : \mathbb{R}^n \rightarrow \mathbb{R} \cup \{+\infty\}$  is

$$f(x) = \sup_{y \in Y} \{ \langle x, Ay + a \rangle - \phi(y) \}$$

- $Y$ : convex •  $\phi : Y \rightarrow \mathbb{R}$ : convex.
- *bilinear F.-t.r.*:  $Y$  is closed &  $\phi$  is affine.

**Fact:** F.-t.r.’s (even bilinear ones) admit *fully algorithmic* calculus: as applied to F.-t. represented convex operands, basic convexity preserving operations straightforwardly yield an F.-t.r. of the result.  
 $\Rightarrow$  *Explicit F.-t.r.’s of convex functions are “common commodity.”*

**Fenchel representation** of a function  $f : \mathbb{R}^n \rightarrow \mathbb{R} \cup \{+\infty\}$  is

$$f(x) = \sup_y \{ \langle x, y \rangle - f^*(y) \}$$

- $f^*$ : convex, proper, l.sc.

**F.r.** of a convex proper l.sc.  $f$  exists and is unique.

**However:** Fenchel representation “exists in the nature,” but, aside of a fistful of simple cases, is **not** available for numerical use.

**Fenchel-type representation** of a function  $f : \mathbb{R}^n \rightarrow \mathbb{R} \cup \{+\infty\}$  is

$$f(x) = \sup_{y \in Y} \{ \langle x, Ay + a \rangle - \phi(y) \}$$

- $Y$ : convex
- $\phi : Y \rightarrow \mathbb{R}$ : convex.
- *bilinear F.-t.r.*:  $Y$  is closed &  $\phi$  is affine.

**Fact:** F.-t.r.’s (even bilinear ones) admit *fully algorithmic* calculus: as applied to F.-t. represented convex operands, basic convexity preserving operations straightforwardly yield an F.-t.r. of the result.  
 $\Rightarrow$  *Explicit F.-t.r.’s of convex functions are “common commodity.”*

**Fenchel representation** of a function  $f : \mathbb{R}^n \rightarrow \mathbb{R} \cup \{+\infty\}$  is

$$f(x) = \sup_y \{ \langle x, y \rangle - f^*(y) \}$$

- $f^*$ : convex, proper, l.sc.

**F.r.** of a convex proper l.sc.  $f$  exists and is unique.

**However:** Fenchel representation “exists in the nature,” but, aside of a fistful of simple cases, is **not** available for numerical use.

**Fenchel-type representation** of a function  $f : \mathbb{R}^n \rightarrow \mathbb{R} \cup \{+\infty\}$  is

$$f(x) = \sup_{y \in Y} \{ \langle x, Ay + a \rangle - \phi(y) \}$$

- $Y$ : convex •  $\phi : Y \rightarrow \mathbb{R}$ : convex.
- **bilinear F.-t.r.:**  $Y$  is closed &  $\phi$  is affine.

**Fact:** **F.-t.r.**’s (even bilinear ones) admit *fully algorithmic* calculus: as applied to **F.-t.** represented convex operands, basic convexity preserving operations straightforwardly yield an **F.-t.r.** of the result.  
 $\Rightarrow$  *Explicit F.-t.r.’s of convex functions are “common commodity.”*

**Fenchel representation** of a function  $f : \mathbb{R}^n \rightarrow \mathbb{R} \cup \{+\infty\}$  is

$$f(x) = \sup_y \{ \langle x, y \rangle - f^*(y) \}$$

- $f^*$ : convex, proper, l.sc.

**F.r.** of a convex proper l.sc.  $f$  exists and is unique.

**However:** Fenchel representation “exists in the nature,” but, aside of a fistful of simple cases, is **not** available for numerical use.

**Fenchel-type representation** of a function  $f : \mathbb{R}^n \rightarrow \mathbb{R} \cup \{+\infty\}$  is

$$f(x) = \sup_{y \in Y} \{ \langle x, Ay + a \rangle - \phi(y) \}$$

- $Y$ : convex •  $\phi : Y \rightarrow \mathbb{R}$ : convex.
- *bilinear F.-t.r.*:  $Y$  is closed &  $\phi$  is affine.

**Fact:** **F.-t.r.**’s (even bilinear ones) admit *fully algorithmic* calculus: as applied to **F.-t.** represented convex operands, basic convexity preserving operations straightforwardly yield an **F.-t.r.** of the result.

*⇒ Explicit F.-t.r.’s of convex functions are “common commodity.”*

**Fenchel representation** of a function  $f : \mathbb{R}^n \rightarrow \mathbb{R} \cup \{+\infty\}$  is

$$f(x) = \sup_y \{ \langle x, y \rangle - f^*(y) \}$$

- $f^*$ : convex, proper, l.sc.

**F.r.** of a convex proper l.sc.  $f$  exists and is unique.

**However:** Fenchel representation “exists in the nature,” but, aside of a fistful of simple cases, is **not** available for numerical use.

**Fenchel-type representation** of a function  $f : \mathbb{R}^n \rightarrow \mathbb{R} \cup \{+\infty\}$  is

$$f(x) = \sup_{y \in Y} \{ \langle x, Ay + a \rangle - \phi(y) \}$$

- $Y$ : convex
- $\phi : Y \rightarrow \mathbb{R}$ : convex.
- *bilinear F.-t.r.*:  $Y$  is closed &  $\phi$  is affine.

**Fact:** **F.-t.r.**’s (even bilinear ones) admit *fully algorithmic* calculus: as applied to **F.-t.** represented convex operands, basic convexity preserving operations straightforwardly yield an **F.-t.r.** of the result.

$\Rightarrow$  *Explicit F.-t.r.’s of convex functions are “common commodity.”*

**Example: F.r.** of the sum  $f_1(x) + f_2(x)$  of two functions given by explicit **F.r.'s** is given by computationally demanding inf-convolution:

$$(f_1 + f_2)^*(y) = \inf_{y_1 + y_2 = y} [f_1^*(y_1) + f_2^*(y_2)]$$

**F.-t.r.** of the sum  $\sum_i f_i$  of functions given by **F.-t.r.'s** is immediate:

$$\begin{aligned} \alpha_i \geq 0, f_i(x) &= \sup_{y_i \in Y_i} [\langle x, A_i y_i + a_i \rangle - \phi_i(y_i)] \Rightarrow \\ \sum_i \alpha_i f_i(x) &= \sup_{y := [y_1; \dots; y_k] \in Y_1 \times \dots \times Y_k} \left[ \langle x, \underbrace{\sum_i \alpha_i [A_i y_i + a_i]}_{Ay + a} \rangle - \underbrace{\sum_i \alpha_i \phi_i(y_i)}_{\phi(y)} \right] \end{aligned}$$

**Example:**  $f(x) := \|\mathcal{A}x - b\| = \max_{y \in Y} \{\langle x, \mathcal{A}^* y \rangle - \langle b, y \rangle\}$

- $Y$ : unit ball of the norm  $\|\cdot\|_*$  conjugate to  $\|\cdot\|$ .

**Fact:** *Strictly feasible conic representation of a function  $f$ :*

$$t \geq f(x) \Leftrightarrow \exists u : \mathcal{A}(x, t, u) \in \mathbf{K}$$

- $\mathcal{A}(x, t, u)$ : affine
  - $\mathbf{K}$ : product of second order and semidefinite cones
- can be straightforwardly converted into an explicit F.-t.r. of  $f$ .*



**Example: F.r.** of the sum  $f_1(x) + f_2(x)$  of two functions given by explicit **F.r.'s** is given by computationally demanding inf-convolution:

$$(f_1 + f_2)^*(y) = \inf_{y_1+y_2=y} [f_1^*(y_1) + f_2^*(y_2)]$$

**F.-t.r.** of the sum  $\sum_i f_i$  of functions given by **F.-t.r.'s** is immediate:

$$\alpha_i \geq 0, f_i(x) = \sup_{y_i \in Y_i} [\langle x, A_i y_i + a_i \rangle - \phi_i(y_i)] \Rightarrow$$

$$\sum_i \alpha_i f_i(x) = \sup_{y:=[y_1;\dots;y_k] \in Y_1 \times \dots \times Y_k} \left[ \underbrace{\langle x, \sum_i \alpha_i [A_i y_i + a_i] \rangle}_{Ay+a} - \underbrace{\sum_i \alpha_i \phi_i(y_i)}_{\phi(y)} \right]$$

**Example:**  $f(x) := \|\mathcal{A}x - b\| = \max_{y \in Y} \{ \langle x, \mathcal{A}^* y \rangle - \langle b, y \rangle \}$

•  $Y$ : unit ball of the norm  $\|\cdot\|_*$  conjugate to  $\|\cdot\|$ .

**Fact:** *Strictly feasible conic representation of a function  $f$ :*

$$t \geq f(x) \Leftrightarrow \exists u : \mathcal{A}(x, t, u) \in \mathbf{K}$$

[•  $\mathcal{A}(x, t, u)$ : affine      •  $\mathbf{K}$ : product of second order and semidefinite cones]  
*can be straightforwardly converted into an explicit F.-t.r. of  $f$ .*

**Example: F.r.** of the sum  $f_1(x) + f_2(x)$  of two functions given by explicit **F.r.'s** is given by computationally demanding inf-convolution:

$$(f_1 + f_2)^*(y) = \inf_{y_1+y_2=y} [f_1^*(y_1) + f_2^*(y_2)]$$

**F.-t.r.** of the sum  $\sum_i f_i$  of functions given by **F.-t.r.'s** is immediate:

$$\alpha_i \geq 0, f_i(x) = \sup_{y_i \in Y_i} [\langle x, A_i y_i + a_i \rangle - \phi_i(y_i)] \Rightarrow$$

$$\sum_i \alpha_i f_i(x) = \sup_{y:=[y_1;\dots;y_k] \in Y_1 \times \dots \times Y_k} \left[ \underbrace{\langle x, \sum_i \alpha_i [A_i y_i + a_i] \rangle}_{Ay+a} - \underbrace{\sum_i \alpha_i \phi_i(y_i)}_{\phi(y)} \right]$$

**Example:**  $f(x) := \|\mathcal{A}x - b\| = \max_{y \in Y} \{ \langle x, \mathcal{A}^* y \rangle - \langle b, y \rangle \}$

•  $Y$ : unit ball of the norm  $\|\cdot\|_*$  conjugate to  $\|\cdot\|$ .

**Fact:** *Strictly feasible conic representation of a function  $f$ :*

$$t \geq f(x) \Leftrightarrow \exists u : \mathcal{A}(x, t, u) \in \mathbf{K}$$

[•  $\mathcal{A}(x, t, u)$ : affine      •  $\mathbf{K}$ : product of second order and semidefinite cones]

*can be straightforwardly converted into an explicit F.-t.r. of  $f$ .*

**Example: F.r.** of the sum  $f_1(x) + f_2(x)$  of two functions given by explicit **F.r.'s** is given by computationally demanding inf-convolution:

$$(f_1 + f_2)^*(y) = \inf_{y_1 + y_2 = y} [f_1^*(y_1) + f_2^*(y_2)]$$

**F.-t.r.** of the sum  $\sum_i f_i$  of functions given by **F.-t.r.'s** is immediate:

$$\alpha_i \geq 0, f_i(x) = \sup_{y_i \in Y_i} [\langle x, A_i y_i + a_i \rangle - \phi_i(y_i)] \Rightarrow$$

$$\sum_i \alpha_i f_i(x) = \sup_{y := [y_1; \dots; y_k] \in Y_1 \times \dots \times Y_k} \left[ \underbrace{\langle x, \sum_i \alpha_i [A_i y_i + a_i] \rangle}_{Ay + a} - \underbrace{\sum_i \alpha_i \phi_i(y_i)}_{\phi(y)} \right]$$

**Example:**  $f(x) := \|\mathcal{A}x - b\| = \max_{y \in Y} \{ \langle x, \mathcal{A}^* y \rangle - \langle b, y \rangle \}$

•  $Y$ : unit ball of the norm  $\|\cdot\|_*$  conjugate to  $\|\cdot\|$ .

**Fact:** *Strictly feasible conic representation of a function  $f$ :*

$$t \geq f(x) \Leftrightarrow \exists u : \mathcal{A}(x, t, u) \in \mathbf{K}$$

[•  $\mathcal{A}(x, t, u)$ : affine      •  $\mathbf{K}$ : product of second order and semidefinite cones]  
*can be straightforwardly converted into an explicit F.-t.r. of  $f$ .*

$$\text{Opt}(P) = \min_{x \in X} f(x) \quad (P)$$

- $X$ : convex compact
- $f$ : convex & Lipschitz

Given an **F.t.r.** of  $f$ :

$$f(x) = \max_{y \in Y} [\langle x, Ay + a \rangle - \phi(y)]$$

with compact  $Y$ , we define the problem dual to (P) as

$$\begin{aligned} \text{Opt}(D) &= \min_{y \in Y} [f^*(y) := \phi(y) - \min_{x \in X} \langle x, Ay + a \rangle] \\ [\text{Opt}(P) &= -\text{Opt}(D)] \end{aligned} \quad (D)$$

**Note:** **LMO** for  $X$  and **First Order Oracle** for  $\phi$  induce an **FOO** for  $f^*$ :

$$y \in Y, \phi'(y) \in \partial_Y \phi(y), x(y) \in \text{Argmin}_X \langle x, Ay + a \rangle \Rightarrow \phi'(y) - A^*x(y) \in \partial_Y f^*(y)$$

**Fact:** Assuming **FOO** for  $\phi$  available and  $Y$  proximal friendly, we can solve (D) to a desired accuracy  $\varepsilon$  by proximal **FOM** producing “accuracy certificates” and use these certificates to get an  $\varepsilon$ -solution to (P).

$$\text{Opt}(P) = \min_{x \in X} f(x) \quad (P)$$

- $X$ : convex compact
- $f$ : convex & Lipschitz

Given an **F.t.r.** of  $f$ :

$$f(x) = \max_{y \in Y} [\langle x, Ay + a \rangle - \phi(y)]$$

with compact  $Y$ , we define the problem dual to (P) as

$$\begin{aligned} \text{Opt}(D) &= \min_{y \in Y} [f^*(y) := \phi(y) - \min_{x \in X} \langle x, Ay + a \rangle] \\ [\text{Opt}(P) &= -\text{Opt}(D)] \end{aligned} \quad (D)$$

**Note:** **LMO** for  $X$  and **First Order Oracle** for  $\phi$  induce an **FOO** for  $f^*$ :

$$y \in Y, \phi'(y) \in \partial_Y \phi(y), x(y) \in \text{Argmin}_X \langle x, Ay + a \rangle \Rightarrow \phi'(y) - A^*x(y) \in \partial_Y f^*(y)$$

**Fact:** Assuming **FOO** for  $\phi$  available and  $Y$  proximal friendly, we can solve (D) to a desired accuracy  $\varepsilon$  by proximal **FOM** producing “accuracy certificates” and use these certificates to get an  $\varepsilon$ -solution to (P).

$$\text{Opt}(P) = \min_{x \in X} f(x) \quad (P)$$

- $X$ : convex compact
- $f$ : convex & Lipschitz

Given an **F.t.r.** of  $f$ :

$$f(x) = \max_{y \in Y} [\langle x, Ay + a \rangle - \phi(y)]$$

with compact  $Y$ , we define the problem dual to (P) as

$$\begin{aligned} \text{Opt}(D) &= \min_{y \in Y} [f^*(y) := \phi(y) - \min_{x \in X} \langle x, Ay + a \rangle] \\ [\text{Opt}(P) &= -\text{Opt}(D)] \end{aligned} \quad (D)$$

**Note:** **LMO** for  $X$  and **First Order Oracle** for  $\phi$  induce an **FOO** for  $f^*$ :

$$y \in Y, \phi'(y) \in \partial_Y \phi(y), x(y) \in \text{Argmin}_X \langle x, Ay + a \rangle \Rightarrow \phi'(y) - A^*x(y) \in \partial_Y f^*(y)$$

**Fact:** Assuming **FOO** for  $\phi$  available and  $Y$  proximal friendly, we can solve (D) to a desired accuracy  $\varepsilon$  by proximal **FOM** producing “accuracy certificates” and use these certificates to get an  $\varepsilon$ -solution to (P).

**Situation:**

- We are processing vector field  $F(y) : Y \rightarrow F$  on convex compact set  $Y$  in Euclidean space  $H$ .
- after  $t$  of the process, we have built a sequence of *search points*  $y_i \in Y$  and have computed  $F(y_i)$ ,  $1 \leq i \leq t$ .

**Definition:** An *accuracy certificate* for  $t$ -step execution protocol

$\mathcal{I}_t = \{y_i, F(y_i) : 1 \leq i \leq t\}$  is a vector  $\lambda \in \mathbb{R}_+^t$  with  $\sum_{i=1}^t \lambda_i = 1$ .

The *residual* and the *approximate solution* associated with  $(\mathcal{I}_t, \lambda)$  are

$$\begin{aligned} \text{Res}(\mathcal{I}_t, \lambda | Y) &= \max_{y \in Y} \sum_{i=1}^t \lambda_i \langle F(y_i), y_i - y \rangle && \text{[residual]} \\ y^t &:= y^t(\mathcal{I}_t, \lambda) = \sum_{i=1}^t \lambda_i y_i && \text{[approximate solution]} \end{aligned}$$

**Note:**  $y^t \in Y$ .

**Theorem A.** Let  $F : Y \rightarrow H$  be vector field induced by **CM/SP/VI** problem with convex structure and problem's domain  $Y \subset H$ . Inaccuracy of the approximate solution  $y^t$  associated with  $t$ -step execution protocol  $\mathcal{I}_t = \{y_i, F(y_i) : 1 \leq i \leq t\}$  and accuracy certificate  $\lambda$  is upper-bounded by the residual.

**Situation:**

- We are processing vector field  $F(y) : Y \rightarrow F$  on convex compact set  $Y$  in Euclidean space  $H$ .
- after  $t$  of the process, we have built a sequence of *search points*  $y_i \in Y$  and have computed  $F(y_i)$ ,  $1 \leq i \leq t$ .

**Definition:** An *accuracy certificate* for *t-step execution protocol*

$\mathcal{I}_t = \{y_i, F(y_i) : 1 \leq i \leq t\}$  is a vector  $\lambda \in \mathbb{R}_+^t$  with  $\sum_{i=1}^t \lambda_i = 1$ .

The *residual* and the *approximate solution* associated with  $(\mathcal{I}_t, \lambda)$  are

$$\begin{aligned} \text{Res}(\mathcal{I}_t, \lambda | Y) &= \max_{y \in Y} \sum_{i=1}^t \lambda_i \langle F(y_i), y_i - y \rangle && \text{[residual]} \\ y^t &:= y^t(\mathcal{I}_t, \lambda) = \sum_{i=1}^t \lambda_i y_i && \text{[approximate solution]} \end{aligned}$$

**Note:**  $y^t \in Y$ .

**Theorem A.** Let  $F : Y \rightarrow H$  be vector field induced by **CM/SP/VI** problem with convex structure and problem's domain  $Y \subset H$ . Inaccuracy of the approximate solution  $y^t$  associated with *t-step execution protocol*  $\mathcal{I}_t = \{y_i, F(y_i) : 1 \leq i \leq t\}$  and accuracy certificate  $\lambda$  is upper-bounded by the residual.



**Situation:**

- We are processing vector field  $F(y) : Y \rightarrow F$  on convex compact set  $Y$  in Euclidean space  $H$ .
- after  $t$  of the process, we have built a sequence of *search points*  $y_i \in Y$  and have computed  $F(y_i)$ ,  $1 \leq i \leq t$ .

**Definition:** An *accuracy certificate* for *t-step execution protocol*

$\mathcal{I}_t = \{y_i, F(y_i) : 1 \leq i \leq t\}$  is a vector  $\lambda \in \mathbb{R}_+^t$  with  $\sum_{i=1}^t \lambda_i = 1$ .

The *residual* and the *approximate solution* associated with  $(\mathcal{I}_t, \lambda)$  are

$$\begin{aligned} \text{Res}(\mathcal{I}_t, \lambda | Y) &= \max_{y \in Y} \sum_{i=1}^t \lambda_i \langle F(y_i), y_i - y \rangle && \text{[residual]} \\ y^t &:= y^t(\mathcal{I}_t, \lambda) = \sum_{i=1}^t \lambda_i y_i && \text{[approximate solution]} \end{aligned}$$

**Note:**  $y^t \in Y$ .

**Theorem A.** Let  $F : Y \rightarrow H$  be vector field induced by **CM/SP/VI** problem with convex structure and problem's domain  $Y \subset H$ . Inaccuracy of the approximate solution  $y^t$  associated with *t-step execution protocol*  $\mathcal{I}_t = \{y_i, F(y_i) : 1 \leq i \leq t\}$  and accuracy certificate  $\lambda$  is upper-bounded by the residual.

**Situation:**

- We are processing vector field  $F(y) : Y \rightarrow F$  on convex compact set  $Y$  in Euclidean space  $H$ .
- after  $t$  of the process, we have built a sequence of *search points*  $y_i \in Y$  and have computed  $F(y_i)$ ,  $1 \leq i \leq t$ .

**Definition:** An *accuracy certificate* for *t-step execution protocol*

$\mathcal{I}_t = \{y_i, F(y_i) : 1 \leq i \leq t\}$  is a vector  $\lambda \in \mathbb{R}_+^t$  with  $\sum_{i=1}^t \lambda_i = 1$ .

The *residual* and the *approximate solution* associated with  $(\mathcal{I}_t, \lambda)$  are

$$\begin{aligned} \text{Res}(\mathcal{I}_t, \lambda | Y) &= \max_{y \in Y} \sum_{i=1}^t \lambda_i \langle F(y_i), y_i - y \rangle && \text{[residual]} \\ y^t &:= y^t(\mathcal{I}_t, \lambda) = \sum_{i=1}^t \lambda_i y_i && \text{[approximate solution]} \end{aligned}$$

**Note:**  $y^t \in Y$ .

**Theorem A.** Let  $F : Y \rightarrow H$  be vector field induced by **CM/SP/VI** problem with convex structure and problem's domain  $Y \subset H$ . Inaccuracy of the approximate solution  $y^t$  associated with *t-step execution protocol*  $\mathcal{I}_t = \{y_i, F(y_i) : 1 \leq i \leq t\}$  and accuracy certificate  $\lambda$  is upper-bounded by the residual.

**Situation:**

- We are processing vector field  $F(y) : Y \rightarrow F$  on convex compact set  $Y$  in Euclidean space  $H$ .
- after  $t$  of the process, we have built a sequence of *search points*  $y_i \in Y$  and have computed  $F(y_i)$ ,  $1 \leq i \leq t$ .

**Definition:** An *accuracy certificate* for *t-step execution protocol*

$\mathcal{I}_t = \{y_i, F(y_i) : 1 \leq i \leq t\}$  is a vector  $\lambda \in \mathbb{R}_+^t$  with  $\sum_{i=1}^t \lambda_i = 1$ .

The *residual* and the *approximate solution* associated with  $(\mathcal{I}_t, \lambda)$  are

$$\begin{aligned} \text{Res}(\mathcal{I}_t, \lambda | Y) &= \max_{y \in Y} \sum_{i=1}^t \lambda_i \langle F(y_i), y_i - y \rangle && \text{[residual]} \\ y^t &:= y^t(\mathcal{I}_t, \lambda) = \sum_{i=1}^t \lambda_i y_i && \text{[approximate solution]} \end{aligned}$$

**Note:**  $y^t \in Y$ .

**Theorem A.** Let  $F : Y \rightarrow H$  be vector field induced by **CM/SP/VI** problem with convex structure and problem's domain  $Y \subset H$ . Inaccuracy of the approximate solution  $y^t$  associated with *t-step execution protocol*  $\mathcal{I}_t = \{y_i, F(y_i) : 1 \leq i \leq t\}$  and accuracy certificate  $\lambda$  is upper-bounded by the residual.

**Theorem A** [Nem., Onn, Rothblum, '10] *Let  $F : Y \rightarrow H$  be vector field induced by **CM/SP/VI** problem with domain  $Y \subset H$ . Inaccuracy of the approximate solution associated with protocol  $\mathcal{I}_t = \{y_i, F(y_i) : 1 \leq i \leq t\}$  and certificate  $\lambda$  is upper-bounded by the residual:*

- *for **CM** problem  $\min_Y g(y)$ :*

$$\epsilon_{\text{CM}}(y^t | g, Y) := g(y^t) - \min_Y g \leq \text{Res}(\mathcal{I}_t, \lambda | Y)$$

- $\epsilon_{\text{CM}}$  is the standard *optimality gap*

**Theorem A** [Nem., Onn., Rothblum, '10] *Let  $F : Y \rightarrow H$  be vector field induced by **CM/SP/VI** problem with domain  $Y \subset H$ . Inaccuracy of the approximate solution associated with protocol  $\mathcal{I}_t = \{y_i, F(y_i) : 1 \leq i \leq t\}$  and certificate  $\lambda$  is upper-bounded by the residual:*



• *for **SP** problem  $\min_{u \in U} \max_{v \in V} g(u, v)$ :*

$$\epsilon_{\text{SP}}(y^t = [u^t; v^t] | g, U, V) := \max_{v \in V} g(u^t, v) - \min_{u \in U} g(u, v^t) \leq \text{Res}(\mathcal{I}_t, \lambda | Y)$$

•  $\epsilon_{\text{SP}}([\bar{u}; \bar{v}] | g, U, V)$  is the *duality gap* – the sum of non-optimality of  $\bar{u}$  and  $\bar{v}$  as approximate solutions to the associated with **SP** primal and dual optimization problems

$$\begin{aligned} \text{Opt(Pr)} &= \min_{u \in U} [\bar{g}(u) := \max_{v \in V} g(u, v)] \\ \text{Opt(Dl)} &= \max_{v \in V} [\underline{g}(v) := \min_{u \in U} g(u, v)] \end{aligned}$$



**Theorem A** [Nem.,Onn,Rothblum,'10] *Let  $F : Y \rightarrow H$  be vector field induced by **CM/SP/VI** problem with domain  $Y \subset H$ . Inaccuracy of the approximate solution associated with protocol  $\mathcal{I}_t = \{y_i, F(y_i) : 1 \leq i \leq t\}$  and certificate  $\lambda$  is upper-bounded by the residual:*

- 
- 
- *for **VI** “find  $y \in Y : \langle F(z), z - y \rangle \geq 0 \forall z \in Y$ .”*  

$$\epsilon_{\text{VI}}(y^t | F, Y) := \sup_{z \in Y} \langle F(z), y^t - z \rangle \leq \text{Res}(\mathcal{I}_t, \lambda | Y)$$
  - $\epsilon_{\text{VI}}(\cdot | \cdot, \cdot)$  is the *dual gap function* associated with **VI**.

$\text{Opt}(P)$	$=$	$\min_{x \in X} \{f(x) := \max_{y \in Y} [\langle x, Ay + a \rangle - \phi(y)]\}$	(P)
$-\text{Opt}(P) = \text{Opt}(D)$	$=$	$\min_{y \in Y} \{f^*(y) := \phi(y) - \min_{x \in X} \langle x, Ay + a \rangle\}$	(D)

**Theorem B** [Cox,Juditsky,Nem.,'13] *Let*

- $F(y) = \phi'(y) - A^*x(y)$ ,  $x(y) \in \text{Argmin}_X \langle x, Ay + a \rangle$  be the monotone operator associated with the dual problem (D),
- $\lambda$  be an accuracy certificate for  $t$ -step execution protocol  $\{y_i \in Y, F(y_i) = \phi'(y_i) - A^*x(y_i)\}$  produced by an **FOM** as applied to the dual problem.

Setting  $x^t = \sum_{i=1}^t \lambda_i x(y_i)$ , we get a feasible solution to the primal problem (P) such that

$$\begin{aligned}
 \epsilon_{\text{CM}}(x^t | f, X) &\leq \text{Gap}(\mathcal{I}_t, \lambda | Y) \\
 &:= \left[ \min_{i \leq t} f^*(y_i) - \sum_{i=1}^t \lambda_i f^*(y_i) \right] + \text{Res}(\mathcal{I}_t, \lambda | Y) \\
 &\leq \text{Res}(\mathcal{I}_t, \lambda | Y)
 \end{aligned}$$

$\text{Opt}(P)$	$=$	$\min_{x \in X} \{f(x) := \max_{y \in Y} [\langle x, Ay + a \rangle - \phi(y)]\}$	(P)
$-\text{Opt}(P) = \text{Opt}(D)$	$=$	$\min_{y \in Y} \{f^*(y) := \phi(y) - \min_{x \in X} \langle x, Ay + a \rangle\}$	(D)

**Theorem B** [Cox,Juditsky,Nem.,'13] *Let*

- $F(y) = \phi'(y) - A^*x(y)$ ,  $x(y) \in \text{Argmin}_X \langle x, Ay + a \rangle$  be the monotone operator associated with the dual problem (D),
- $\lambda$  be an accuracy certificate for  $t$ -step execution protocol  $\{y_i \in Y, F(y_i) = \phi'(y_i) - A^*x(y_i)\}$  produced by an **FOM** as applied to the dual problem.

Setting  $x^t = \sum_{i=1}^t \lambda_i x(y_i)$ , we get a feasible solution to the primal problem (P) such that

$$\begin{aligned}
 \epsilon_{\text{CM}}(x^t | f, X) &\leq \text{Gap}(\mathcal{I}_t, \lambda | Y) \\
 &:= \left[ \min_{i \leq t} f^*(y_i) - \sum_{i=1}^t \lambda_i f^*(y_i) \right] + \text{Res}(\mathcal{I}_t, \lambda | Y) \\
 &\leq \text{Res}(\mathcal{I}_t, \lambda | Y)
 \end{aligned}$$



## Fact: Standard FOMs:

- Subgradient/Mirror Descent and its Bundle versions,
- Cutting plane algorithms, including Ellipsoid/Inscribed Ellipsoid methods,
- Nesterov's Fast Gradients,...

produce certificates with gap/residual obeying the standard efficiency estimates of the algorithms.

**Example:** Let  $F$  be a vector field on convex domain  $Y$  of diameter  $D$  w.r.t. a norm  $\|\cdot\|$ .

- $t$ -step *Mirror Descent* as applied to  $(F, Y)$  ensures  $\text{Res} \leq C_{\|\cdot\|} \frac{LD}{\sqrt{t}}$ , provided  $\|F(y)\|_* \leq L \forall y \in Y$ .  
For good norms ( $\ell_p$  with  $1 \leq p \leq 2$ , nuclear,...),  $C_{\|\cdot\|}$  is (nearly) dimension independent.
- $t$ -step *Mirror Prox* as applied to  $(F, Y)$  ensures  $\text{Res} \leq C_{\|\cdot\|}^2 \frac{LD^2}{t}$ , provided  $\|F(y) - F(z)\|_* \leq L\|y - z\| \forall y, z \in Y$

$\Rightarrow$  **F-t.r.'s** with proximal friendly and bounded dual domains induce  $O(1/\sqrt{t})$ -converging **LMO-based algorithms for Nonsmooth CM**

## Fact: Standard FOMs:

- Subgradient/Mirror Descent and its Bundle versions,
- Cutting plane algorithms, including Ellipsoid/Inscribed Ellipsoid methods,
- Nesterov's Fast Gradients,...

produce certificates with gap/residual obeying the standard efficiency estimates of the algorithms.

**Example:** Let  $F$  be a vector field on convex domain  $Y$  of diameter  $D$  w.r.t. a norm  $\|\cdot\|$ .

- $t$ -step *Mirror Descent* as applied to  $(F, Y)$  ensures  $\text{Res} \leq C_{\|\cdot\|} \frac{LD}{\sqrt{t}}$ , provided  $\|F(y)\|_* \leq L \forall y \in Y$ .

*For good norms ( $\ell_p$  with  $1 \leq p \leq 2$ , nuclear,...),  $C_{\|\cdot\|}$  is (nearly) dimension independent.*

- $t$ -step *Mirror Prox* as applied to  $(F, Y)$  ensures  $\text{Res} \leq C_{\|\cdot\|}^2 \frac{LD^2}{t}$ , provided  $\|F(y) - F(z)\|_* \leq L\|y - z\| \forall y, z \in Y$

$\Rightarrow$  **F-t.r.'s** with proximal friendly and bounded dual domains induce  $O(1/\sqrt{t})$ -converging **LMO-based algorithms for Nonsmooth CM**

## Fact: Standard FOMs:

- Subgradient/Mirror Descent and its Bundle versions,
- Cutting plane algorithms, including Ellipsoid/Inscribed Ellipsoid methods,
- Nesterov's Fast Gradients,...

produce certificates with gap/residual obeying the standard efficiency estimates of the algorithms.

**Example:** Let  $F$  be a vector field on convex domain  $Y$  of diameter  $D$  w.r.t. a norm  $\|\cdot\|$ .

- $t$ -step *Mirror Descent* as applied to  $(F, Y)$  ensures  $\text{Res} \leq C_{\|\cdot\|} \frac{LD}{\sqrt{t}}$ , provided  $\|F(y)\|_* \leq L \forall y \in Y$ .  
For good norms ( $\ell_p$  with  $1 \leq p \leq 2$ , nuclear,...),  $C_{\|\cdot\|}$  is (nearly) dimension independent.
- $t$ -step *Mirror Prox* as applied to  $(F, Y)$  ensures  $\text{Res} \leq C_{\|\cdot\|}^2 \frac{LD^2}{t}$ , provided  $\|F(y) - F(z)\|_* \leq L\|y - z\| \forall y, z \in Y$

$\Rightarrow$  **F-t.r.'s** with proximal friendly and bounded dual domains induce  $O(1/\sqrt{t})$ -converging **LMO-based algorithms for Nonsmooth CM**

## Fact: Standard FOMs:

- Subgradient/Mirror Descent and its Bundle versions,
- Cutting plane algorithms, including Ellipsoid/Inscribed Ellipsoid methods,
- Nesterov's Fast Gradients,...

produce certificates with gap/residual obeying the standard efficiency estimates of the algorithms.

**Example:** Let  $F$  be a vector field on convex domain  $Y$  of diameter  $D$  w.r.t. a norm  $\|\cdot\|$ .

- $t$ -step *Mirror Descent* as applied to  $(F, Y)$  ensures  $\text{Res} \leq C_{\|\cdot\|} \frac{LD}{\sqrt{t}}$ , provided  $\|F(y)\|_* \leq L \forall y \in Y$ .  
For good norms ( $\ell_p$  with  $1 \leq p \leq 2$ , nuclear,...),  $C_{\|\cdot\|}$  is (nearly) dimension independent.
- $t$ -step *Mirror Prox* as applied to  $(F, Y)$  ensures  $\text{Res} \leq C_{\|\cdot\|}^2 \frac{LD^2}{t}$ , provided  $\|F(y) - F(z)\|_* \leq L\|y - z\| \forall y, z \in Y$

$\Rightarrow$  F-t.r.'s with proximal friendly and bounded dual domains induce  $O(1/\sqrt{t})$ -converging LMO-based algorithms for Nonsmooth CM

## Fact: Standard FOMs:

- Subgradient/Mirror Descent and its Bundle versions,
- Cutting plane algorithms, including Ellipsoid/Inscribed Ellipsoid methods,
- Nesterov's Fast Gradients,...

produce certificates with gap/residual obeying the standard efficiency estimates of the algorithms.

**Example:** Let  $F$  be a vector field on convex domain  $Y$  of diameter  $D$  w.r.t. a norm  $\|\cdot\|$ .

- $t$ -step *Mirror Descent* as applied to  $(F, Y)$  ensures  $\text{Res} \leq C_{\|\cdot\|} \frac{LD}{\sqrt{t}}$ , provided  $\|F(y)\|_* \leq L \forall y \in Y$ .  
For good norms ( $\ell_p$  with  $1 \leq p \leq 2$ , nuclear,...),  $C_{\|\cdot\|}$  is (nearly) dimension independent.
- $t$ -step *Mirror Prox* as applied to  $(F, Y)$  ensures  $\text{Res} \leq C_{\|\cdot\|}^2 \frac{LD^2}{t}$ , provided  $\|F(y) - F(z)\|_* \leq L\|y - z\| \forall y, z \in Y$

$\Rightarrow$  **F-t.r.'s** with proximal friendly and bounded dual domains induce  $O(1/\sqrt{t})$ -converging **LMO-based algorithms for Nonsmooth CM.**

**Observation: F.-t.r.**  $f(x) = \max_{y \in Y} \{\langle x, Ay + a \rangle - \phi(x)\}$  of a convex function  $f$  with proximal friendly  $Y$  and affine  $\phi$  induces easy to compute smooth approximation of  $f$

$$f_\epsilon(x) = \max_{y \in Y} [\langle x, Ay + a \rangle - \phi(y) + \epsilon d(y)]$$

$d(\cdot)$ : strongly convex function with easy-to-minimize linear perturbations

and one can minimize this approximation over an **LMO**-represented domain  $X$  by Conditional Gradients, resulting in the same  $O(1/\sqrt{t})$  convergence as **F.-t.r./Accuracy Certificates** machinery.

**Note:** This is nothing but *Nesterov's smoothing* with Conditional Gradients in the role of Fast Gradients.

**Observation:** **F.-t.r.**  $f(x) = \max_{y \in Y} \{\langle x, Ay + a \rangle - \phi(x)\}$  of a convex function  $f$  with proximal friendly  $Y$  and affine  $\phi$  induces easy to compute smooth approximation of  $f$

$$f_\epsilon(x) = \max_{y \in Y} [\langle x, Ay + a \rangle - \phi(y) + \epsilon d(y)]$$

$d(\cdot)$ : strongly convex function with easy-to-minimize linear perturbations

and one can minimize this approximation over an **LMO**-represented domain  $X$  by Conditional Gradients, resulting in the same  $O(1/\sqrt{t})$  convergence as **F.-t.r./Accuracy Certificates** machinery.

**Note:** This is nothing but *Nesterov's smoothing* with Conditional Gradients in the role of Fast Gradients.

## Definition [Juditsky,Nem.,'14] Let

- $\Phi(\cdot) : X \rightarrow E$  be a monotone operator on convex domain  $X$  in Euclidean space  $E$
- $H$  be Euclidean space and  $y \mapsto Ay + a : H \rightarrow E$  be an affine mapping
- $G(\cdot) : Y \rightarrow H$  be monotone operator on convex domain  $Y \subset H$ , and for all  $x \in X$  the VI “find  $y \in Y : \langle G(z) - A^*x, z - y \rangle \geq 0 \forall z \in Y$ ” has a *strong* solution  $y(x) \in Y : \langle G(y(x)) - A^*x, z - y(x) \rangle \geq 0 \forall z \in Y$ .
- It holds  $\Phi(x) = Ay(x) + a \forall x \in X$

We say that

- $H, A, a, Y, G(\cdot), y(\cdot)$  form a Fenchel-type representation of  $\Phi$  on  $X$ .

Informally: **F.-t.r.** of  $\Phi$  is

$$\Phi(x) = AG^{-1}(A^*x) + a$$

- the VI “find  $y \in Y : \langle \Psi(v), v - y \rangle \geq 0 \forall v \in Y$ ” with

$$\Psi(v) = G(v) - A^*x(v), \quad x(v) \in \text{Argmin}_{x \in X} \langle x, Av + a \rangle$$

is the *dual* of the *primal* VI “find  $x \in X : \langle \Phi(u), u - x \rangle \geq 0 \forall u \in X$ ”

**Note:**  $\Psi(\cdot)$  is monotone on  $Y$ , and computing  $\Psi(\cdot)$  reduces to computing  $G(\cdot)$  and a call to **LMO** for  $X$ .



**Definition** [Juditsky,Nem.,'14] Let

- $\Phi(\cdot) : X \rightarrow E$  be a monotone operator on convex domain  $X$  in Euclidean space  $E$
- $H$  be Euclidean space and  $y \mapsto Ay + a : H \rightarrow E$  be an affine mapping
- $G(\cdot) : Y \rightarrow H$  be monotone operator on convex domain  $Y \subset H$ , and for all  $x \in X$  the VI “find  $y \in Y : \langle G(z) - A^*x, z - y \rangle \geq 0 \forall z \in Y$ ” has a *strong* solution  $y(x) \in Y : \langle G(y(x)) - A^*x, z - y(x) \rangle \geq 0 \forall z \in Y$ .
- It holds  $\Phi(x) = Ay(x) + a \forall x \in X$

We say that

- $H, A, a, Y, G(\cdot), y(\cdot)$  form a Fenchel-type representation of  $\Phi$  on  $X$ .

**Informally:** F-t.r. of  $\Phi$  is

$$\Phi(x) = AG^{-1}(A^*x) + a$$

- the VI “find  $y \in Y : \langle \Psi(v), v - y \rangle \geq 0 \forall v \in Y$ ” with  
 $\Psi(v) = G(v) - A^*x(v), x(v) \in \text{Argmin}_{x \in X} \langle x, Av + a \rangle$

is the *dual* of the *primal* VI “find  $x \in X : \langle \Phi(u), u - x \rangle \geq 0 \forall u \in X$ ”

**Note:**  $\Psi(\cdot)$  is monotone on  $Y$ , and computing  $\Psi(\cdot)$  reduces to computing  $G(\cdot)$  and a call to **LMO** for  $X$ .

**Definition** [Juditsky,Nem.,'14] Let

- $\Phi(\cdot) : X \rightarrow E$  be a monotone operator on convex domain  $X$  in Euclidean space  $E$
- $H$  be Euclidean space and  $y \mapsto Ay + a : H \rightarrow E$  be an affine mapping
- $G(\cdot) : Y \rightarrow H$  be monotone operator on convex domain  $Y \subset H$ , and for all  $x \in X$  the VI “find  $y \in Y : \langle G(z) - A^*x, z - y \rangle \geq 0 \forall z \in Y$ ” has a *strong* solution  $y(x) \in Y : \langle G(y(x)) - A^*x, z - y(x) \rangle \geq 0 \forall z \in Y$ .
- It holds  $\Phi(x) = Ay(x) + a \forall x \in X$

We say that

- $H, A, a, Y, G(\cdot), y(\cdot)$  form a Fenchel-type representation of  $\Phi$  on  $X$ .

**Informally:** F-t.r. of  $\Phi$  is

$$\Phi(x) = AG^{-1}(A^*x) + a$$

- the VI “find  $y \in Y : \langle \Psi(v), v - y \rangle \geq 0 \forall v \in Y$ ” with  
 $\Psi(v) = G(v) - A^*x(v), x(v) \in \text{Argmin}_{x \in X} \langle x, Av + a \rangle$

is the *dual* of the *primal* VI “find  $x \in X : \langle \Phi(u), u - x \rangle \geq 0 \forall u \in X$ ”

**Note:**  $\Psi(\cdot)$  is monotone on  $Y$ , and computing  $\Psi(\cdot)$  reduces to computing  $G(\cdot)$  and a call to LMO for  $X$ .

**Definition** [Juditsky,Nem.,'14] Let

- $\Phi(\cdot) : X \rightarrow E$  be a monotone operator on convex domain  $X$  in Euclidean space  $E$
- $H$  be Euclidean space and  $y \mapsto Ay + a : H \rightarrow E$  be an affine mapping
- $G(\cdot) : Y \rightarrow H$  be monotone operator on convex domain  $Y \subset H$ , and for all  $x \in X$  the VI “find  $y \in Y : \langle G(z) - A^*x, z - y \rangle \geq 0 \forall z \in Y$ ” has a *strong* solution  $y(x) \in Y : \langle G(y(x)) - A^*x, z - y(x) \rangle \geq 0 \forall z \in Y$ .
- It holds  $\Phi(x) = Ay(x) + a \forall x \in X$

We say that

- $H, A, a, Y, G(\cdot), y(\cdot)$  form a Fenchel-type representation of  $\Phi$  on  $X$ .

**Informally:** F-t.r. of  $\Phi$  is

$$\Phi(x) = AG^{-1}(A^*x) + a$$

- the VI “find  $y \in Y : \langle \Psi(v), v - y \rangle \geq 0 \forall v \in Y$ ” with  
 $\Psi(v) = G(v) - A^*x(v), x(v) \in \text{Argmin}_{x \in X} \langle x, Av + a \rangle$

is the *dual* of the *primal* VI “find  $x \in X : \langle \Phi(u), u - x \rangle \geq 0 \forall u \in X$ ”

**Note:**  $\Psi(\cdot)$  is monotone on  $Y$ , and computing  $\Psi(\cdot)$  reduces to computing  $G(\cdot)$  and a call to **LMO** for  $X$ .

**Example:** Affine monotone operator  $\Phi(x) = Sx + s : E \rightarrow E$  on a convex compact domain  $X \subset E$  admits **F.-t.r.** with the data

$$H = E, Ay + a \equiv Sy + s, Y \supset X, G(y) = S^*y, y(x) = x$$

**Fact:**

- **F.-t.r.**'s of monotone operators admit *fully algorithmic* calculus: as applied to **F.-t.** represented monotone operands, basic monotonicity-preserving operations with vector fields:

- summation with nonnegative weights,
- direct summation,
- affine substitution of argument,

straightforwardly yield an **F.-t.r.** of the result.

$\Rightarrow$  **F.-t.r.**'s of monotone operators form a “common commodity.”

**Example:** Affine monotone operator  $\Phi(x) = Sx + s : E \rightarrow E$  on a convex compact domain  $X \subset E$  admits **F.-t.r.** with the data

$$H = E, Ay + a \equiv Sy + s, Y \supset X, G(y) = S^*y, y(x) = x$$

### Fact:

- **F.-t.r.**'s of monotone operators admit *fully algorithmic* calculus: as applied to **F.-t.** represented monotone operands, basic monotonicity-preserving operations with vector fields:

- summation with nonnegative weights,
- direct summation,
- affine substitution of argument,

straightforwardly yield an **F.-t.r.** of the result.

⇒ **F.-t.r.**'s of monotone operators form a “common commodity.”

**Example:** Affine monotone operator  $\Phi(x) = Sx + s : E \rightarrow E$  on a convex compact domain  $X \subset E$  admits **F.-t.r.** with the data

$$H = E, Ay + a \equiv Sy + s, Y \supset X, G(y) = S^*y, y(x) = x$$

### Fact:

- **F.-t.r.**'s of monotone operators admit *fully algorithmic* calculus: as applied to **F.-t.** represented monotone operands, basic monotonicity-preserving operations with vector fields:

- summation with nonnegative weights,
- direct summation,
- affine substitution of argument,

straightforwardly yield an **F.-t.r.** of the result.

$\Rightarrow$  **F.-t.r.**'s of monotone operators form a “common commodity.”

$$\text{find } x \in X : \langle \Phi(u), u - x \rangle \geq 0 \quad \forall u \in X \quad (\text{P})$$

$$\Phi(x) = Ay(x) + a \text{ \& } \langle G(y(x)) - A^*x, v - y(x) \rangle \geq 0 \quad \forall v \in Y \quad (1)$$

$$\Psi(y) := G(y) - A^*x(y) \text{ \& } x(y) \in \text{Argmin}_{x \in X} \langle x, Ay + a \rangle \quad (2)$$

$$\Rightarrow \text{find } y \in Y : \langle \Psi(v), v - y \rangle \geq 0 \quad \forall v \in Y \quad (\text{D})$$

## Theorem C [Juditsky,Nem.,'14] *Let*

- $\Phi(\cdot) : X \rightarrow E$  be a monotone vector field with **F-t.r.** (1) on convex compact subset  $X$  of Euclidean space  $E$ , and

- $\lambda$  be an accuracy certificate for  $t$ -step execution protocol

$$\mathcal{I}_t = \{y_i, \Psi(y_i) : 1 \leq i \leq t\}$$

produced by an **FOM** as applied to the **dual VI (D)**.

The point  $x^t = \sum_{i=1}^t \lambda_i x(y_i)$  is a feasible solution to the primal **VI (P)**, and the inaccuracy of  $x^t$  w.r.t. (P) can be upper-bounded as

$$\epsilon_{\text{VI}}(x^t | F, X) \leq \text{Res}(\mathcal{I}_t, \lambda | \Psi, Y).$$

When  $\Phi$  is the affine monotone vector field associated with **SP** problem

$$\min_{u \in U} \max_{v \in V} g(u, v)$$

with bilinear  $g(u, v)$  and convex compact  $U, V$ , we have also

$$\epsilon_{\text{SP}}(x^t | g, U, V) \leq \text{Res}(\mathcal{I}_t, \lambda | \Psi, Y).$$

find $x \in X : \langle \Phi(u), u - x \rangle \geq 0 \quad \forall u \in X$	(P)
$\Phi(x) = Ay(x) + a \ \& \ \langle G(y(x)) - A^*x, v - y(x) \rangle \geq 0 \quad \forall v \in Y$	(1)
$\Psi(y) := G(y) - A^*x(y) \ \& \ x(y) \in \text{Argmin}_{x \in X} \langle x, Ay + a \rangle$	(2)
$\Rightarrow \text{find } y \in Y : \langle \Psi(v), v - y \rangle \geq 0 \quad \forall v \in Y$	(D)

**Theorem C** [Juditsky,Nem.,'14] *Let*

- $\Phi(\cdot) : X \rightarrow E$  be a monotone vector field with **F-t.r.** (1) on convex compact subset  $X$  of Euclidean space  $E$ , and
- $\lambda$  be an accuracy certificate for  $t$ -step execution protocol

$$\mathcal{I}_t = \{y_i, \Psi(y_i) : 1 \leq i \leq t\}$$

produced by an **FOM** as applied to the **dual VI (D)**.

The point  $x^t = \sum_{i=1}^t \lambda_i x(y_i)$  is a feasible solution to the primal **VI (P)**, and the inaccuracy of  $x^t$  w.r.t. (P) can be upper-bounded as

$$\epsilon_{\text{VI}}(x^t | F, X) \leq \text{Res}(\mathcal{I}_t, \lambda | \Psi, Y).$$

When  $\Phi$  is the affine monotone vector field associated with **SP** problem

$$\min_{u \in U} \max_{v \in V} g(u, v)$$

with bilinear  $g(u, v)$  and convex compact  $U, V$ , we have also

$$\epsilon_{\text{SP}}(x^t | g, U, V) \leq \text{Res}(\mathcal{I}_t, \lambda | \Psi, Y).$$



$$\text{find } x \in X : \langle \Phi(u), u - x \rangle \geq 0 \quad \forall u \in X \quad (\text{P})$$

$$\Phi(x) = Ay(x) + a \text{ \& } \langle G(y(x)) - A^*x, v - y(x) \rangle \geq 0 \quad \forall v \in Y \quad (1)$$

$$\Psi(y) := G(y) - A^*x(y) \text{ \& } x(y) \in \text{Argmin}_{x \in X} \langle x, Ay + a \rangle \quad (2)$$

$$\Rightarrow \text{find } y \in Y : \langle \Psi(v), v - y \rangle \geq 0 \quad \forall v \in Y \quad (\text{D})$$

**Theorem C** [Juditsky,Nem.,'14] *Let*

- $\Phi(\cdot) : X \rightarrow E$  be a monotone vector field with **F-t.r.** (1) on convex compact subset  $X$  of Euclidean space  $E$ , and

- $\lambda$  be an accuracy certificate for  $t$ -step execution protocol

$$\mathcal{I}_t = \{y_i, \Psi(y_i) : 1 \leq i \leq t\}$$

produced by an **FOM** as applied to the **dual VI (D)**.

The point  $x^t = \sum_{i=1}^t \lambda_i x(y_i)$  is a feasible solution to the primal **VI (P)**, and the inaccuracy of  $x^t$  w.r.t. (P) can be upper-bounded as

$$\epsilon_{\text{VI}}(x^t | F, X) \leq \text{Res}(\mathcal{I}_t, \lambda | \Psi, Y).$$

When  $\Phi$  is the affine monotone vector field associated with **SP** problem

$$\min_{u \in U} \max_{v \in V} g(u, v)$$

with bilinear  $g(u, v)$  and convex compact  $U, V$ , we have also

$$\epsilon_{\text{SP}}(x^t | g, U, V) \leq \text{Res}(\mathcal{I}_t, \lambda | \Psi, Y).$$

$$\begin{array}{ll}
 \text{find } x \in X : \langle \Phi(u), u - x \rangle \geq 0 \quad \forall u \in X & \text{(P)} \\
 \Phi(x) = Ay(x) + a \text{ \& } \langle G(y(x)) - A^*x, v - y(x) \rangle \geq 0 \quad \forall v \in Y & (1) \\
 \Psi(y) := G(y) - A^*x(y) \text{ \& } x(y) \in \text{Argmin}_{x \in X} \langle x, Ay + a \rangle & (2) \\
 \Downarrow & \\
 \text{find } y \in Y : \langle \Psi(v), v - y \rangle \geq 0 \quad \forall v \in Y & \text{(D)}
 \end{array}$$

**Corollary:** When  $Y$  is proximal friendly and  $G(\cdot)$  is bounded on  $Y$ , **F.t.r.** (1) induces  $O(1/\sqrt{t})$ -converging **LMO**-based algorithm for (P).

**Corollary:** **F.t.r.** (1) with bounded  $Y \subset \mathbb{R}^n$  induces **LMO**-based linearly converging algorithm for (P) with convergence ratio  $\vartheta$  depending solely on  $n$  (e.g.,  $\vartheta = 1 - O(1/n)$  or  $\vartheta = 1 - O(1/n^2)$ )

**Note:** In some applications,  $\dim X$  is huge and  $\dim Y$  is small (tens/hundreds)...

$$\begin{array}{ll}
 \text{find } x \in X : \langle \Phi(u), u - x \rangle \geq 0 \quad \forall u \in X & \text{(P)} \\
 \Phi(x) = Ay(x) + a \text{ \& } \langle G(y(x)) - A^*x, v - y(x) \rangle \geq 0 \quad \forall v \in Y & (1) \\
 \Psi(y) := G(y) - A^*x(y) \text{ \& } x(y) \in \text{Argmin}_{x \in X} \langle x, Ay + a \rangle & (2) \\
 \Downarrow & \\
 \text{find } y \in Y : \langle \Psi(v), v - y \rangle \geq 0 \quad \forall v \in Y & \text{(D)}
 \end{array}$$

**Corollary:** When  $Y$  is proximal friendly and  $G(\cdot)$  is bounded on  $Y$ , **F.t.r.** (1) induces  $O(1/\sqrt{t})$ -converging **LMO**-based algorithm for (P).

**Corollary:** **F.t.r.** (1) with bounded  $Y \subset \mathbb{R}^n$  induces **LMO**-based linearly converging algorithm for (P) with convergence ratio  $\vartheta$  depending solely on  $n$  (e.g.,  $\vartheta = 1 - O(1/n)$  or  $\vartheta = 1 - O(1/n^2)$ )

**Note:** In some applications,  $\dim X$  is huge and  $\dim Y$  is small (tens/hundreds)...

$$\begin{array}{ll}
 \text{find } x \in X : \langle \Phi(u), u - x \rangle \geq 0 \quad \forall u \in X & \text{(P)} \\
 \Phi(x) = Ay(x) + a \text{ \& } \langle G(y(x)) - A^*x, v - y(x) \rangle \geq 0 \quad \forall v \in Y & \text{(1)} \\
 \Psi(y) := G(y) - A^*x(y) \text{ \& } x(y) \in \text{Argmin}_{x \in X} \langle x, Ay + a \rangle & \text{(2)} \\
 \Downarrow & \\
 \text{find } y \in Y : \langle \Psi(v), v - y \rangle \geq 0 \quad \forall v \in Y & \text{(D)}
 \end{array}$$

**Corollary:** When  $Y$  is proximal friendly and  $G(\cdot)$  is bounded on  $Y$ , **F.t.r.** (1) induces  $O(1/\sqrt{t})$ -converging **LMO**-based algorithm for (P).

**Corollary:** **F.t.r.** (1) with bounded  $Y \subset \mathbb{R}^n$  induces **LMO**-based linearly converging algorithm for (P) with convergence ratio  $\vartheta$  depending solely on  $n$  (e.g.,  $\vartheta = 1 - O(1/n)$  or  $\vartheta = 1 - O(1/n^2)$ )

**Note:** In some applications,  $\dim X$  is huge and  $\dim Y$  is small (tens/hundreds)...

**Fact:** *Nonsmooth CM* problems *with smooth F.-t.r.'s* and *smooth SP/VI's* on *proximal friendly* domains admit  $O(1/t)$ -converging *proximal FOMs*.

**In contrast,** *the best known convergence rate of LMO-based FOMs for nonsmooth CM and smooth/nonsmooth SP/VI problems is  $O(1/\sqrt{t})$ .*

**Reason:** For a typical **LMO**-based **FOM**, *approximate solution generated in  $t$  calls to LMO belongs to the linear span of the minimizers of the  $t$  linear forms processed by LMO in these calls. Under this restriction, already pretty simple nonsmooth CM's and smooth/nonsmooth SP's/VI's on domains like  $\ell_1$ /nuclear norm balls provably cannot be solved faster than at  $O(1/\sqrt{t})$  rate.*

**Fact:** *Nonsmooth CM* problems *with smooth F.-t.r.'s* and *smooth SP/VI's* on *proximal friendly* domains admit  $O(1/t)$ -converging *proximal FOMs*.

**In contrast,** *the best known convergence rate of LMO-based FOMs for nonsmooth CM and smooth/nonsmooth SP/VI problems is  $O(1/\sqrt{t})$ .*

**Reason:** For a typical LMO-based FOM, *approximate solution generated in  $t$  calls to LMO belongs to the linear span of the minimizers of the  $t$  linear forms processed by LMO in these calls. Under this restriction, already pretty simple nonsmooth CM's and smooth/nonsmooth SP's/VI's on domains like  $\ell_1$ /nuclear norm balls provably cannot be solved faster than at  $O(1/\sqrt{t})$  rate.*

**Fact:** *Nonsmooth CM* problems *with smooth F.-t.r.'s* and *smooth SP/VI's* on *proximal friendly* domains admit  $O(1/t)$ -converging *proximal FOMs*.

**In contrast,** *the best known convergence rate of LMO-based FOMs for nonsmooth CM and smooth/nonsmooth SP/VI problems is  $O(1/\sqrt{t})$ .*

**Reason:** For a typical **LMO**-based **FOM**, *approximate solution generated in  $t$  calls to LMO belongs to the linear span of the minimizers of the  $t$  linear forms processed by LMO in these calls.*

*Under this restriction, already pretty simple nonsmooth CM's and smooth/nonsmooth SP's/VI's on domains like  $\ell_1$ /nuclear norm balls provably cannot be solved faster than at  $O(1/\sqrt{t})$  rate.*

**Fact:** *Nonsmooth CM* problems *with smooth F.-t.r.'s* and *smooth SP/VI's* on *proximal friendly* domains admit  $O(1/t)$ -converging *proximal FOMs*.

**In contrast,** *the best known convergence rate of LMO-based FOMs for nonsmooth CM and smooth/nonsmooth SP/VI problems is  $O(1/\sqrt{t})$ .*

**Reason:** For a typical **LMO**-based **FOM**, *approximate solution generated in  $t$  calls to LMO belongs to the linear span of the minimizers of the  $t$  linear forms processed by LMO in these calls. Under this restriction, already pretty simple nonsmooth CM's and smooth/nonsmooth SP's/VI's on domains like  $\ell_1$ /nuclear norm balls provably cannot be solved faster than at  $O(1/\sqrt{t})$  rate.*



**Problem:** Given noisy observations  $a_{ij}$ ,  $(i, j) \in \Omega$ , of  $N$  entries in a  $p \times p$  matrix, find matrix of nuclear norm  $\leq 1$  which is the best fit to the observations in the uniform norm.

**Note:**

- **LMO** for the domain  $X$  of the problem of interest (unit Nuclear norm ball in  $\mathbb{R}^{p \times p}$ ) is readily given by Power method.
- The objective admits bilinear **F-t.r.**, the domain  $Y$  of the dual problem is the proximal-friendly unit  $\ell_1$ -ball in  $\mathbb{R}^N$ .

**Worst-case efficiency:**  $\epsilon_{\text{CM}}(x^t) \leq O(1) \frac{\ln(N)[1 + \max_{i,j} |a_{ij}|]}{\sqrt{t}}$

**Results, I:** Restricted Memory Bundle-Level algorithm on the dual of a low size ( $p = 512$ ,  $N = 512$ ) Matrix Completion:

Memory depth	1	33	65	129
$\text{Gap}_1 / \text{Gap}_{1024}$	114	164	350	3253

**Results, II:** Subgradient Descent on the dual of Matrix Completion:

$p$	$N$	$\text{Gap}_1$	$\text{Gap}_1 / \text{Gap}_{32}$	$\text{Gap}_1 / \text{Gap}_{128}$	$\text{Gap}_1 / \text{Gap}_{1024}$	CPU, sec
2048	8192	1.81e-1	171.2	213.8	451.4	521.3
4096	16384	3.74e-1	335.4	1060.8	1287.3	1524.8
8192	16384	2.54e-1	37.8	875.8	1183.6	3644.0

**Problem:** Given noisy observations  $a_{ij}$ ,  $(i, j) \in \Omega$ , of  $N$  entries in a  $p \times p$  matrix, find matrix of nuclear norm  $\leq 1$  which is the best fit to the observations in the uniform norm.

**Note:**

- **LMO** for the domain  $X$  of the problem of interest (unit Nuclear norm ball in  $\mathbb{R}^{p \times p}$ ) is readily given by Power method.
- The objective admits bilinear **F-t.r.**, the domain  $Y$  of the dual problem is the proximal-friendly unit  $\ell_1$ -ball in  $\mathbb{R}^N$ .

**Worst-case efficiency:**  $\epsilon_{\text{CM}}(x^t) \leq O(1) \frac{\ln(N)[1 + \max_{i,j} |a_{ij}|]}{\sqrt{t}}$

**Results, I:** Restricted Memory Bundle-Level algorithm on the dual of a low size ( $p = 512$ ,  $N = 512$ ) Matrix Completion:

Memory depth	1	33	65	129
Gap <sub>1</sub> /Gap <sub>1024</sub>	114	164	350	3253

**Results, II:** Subgradient Descent on the dual of Matrix Completion:

$p$	$N$	Gap <sub>1</sub>	Gap <sub>1</sub> /Gap <sub>32</sub>	Gap <sub>1</sub> /Gap <sub>128</sub>	Gap <sub>1</sub> /Gap <sub>1024</sub>	CPU, sec
2048	8192	1.81e-1	171.2	213.8	451.4	521.3
4096	16384	3.74e-1	335.4	1060.8	1287.3	1524.8
8192	16384	2.54e-1	37.8	875.8	1183.6	3644.0

**Problem:** Given noisy observations  $a_{ij}, (i, j) \in \Omega$ , of  $N$  entries in a  $p \times p$  matrix, find matrix of nuclear norm  $\leq 1$  which is the best fit to the observations in the uniform norm.

**Note:**

- **LMO** for the domain  $X$  of the problem of interest (unit Nuclear norm ball in  $\mathbb{R}^{p \times p}$ ) is readily given by Power method.
- The objective admits bilinear **F-t.r.**, the domain  $Y$  of the dual problem is the proximal-friendly unit  $\ell_1$ -ball in  $\mathbb{R}^N$ .

**Worst-case efficiency:**  $\epsilon_{\text{CM}}(x^t) \leq O(1) \frac{\ln(N)[1 + \max_{i,j} |a_{ij}|]}{\sqrt{t}}$

**Results, I:** Restricted Memory Bundle-Level algorithm on the dual of a low size ( $p = 512, N = 512$ ) Matrix Completion:

Memory depth	1	33	65	129
$\text{Gap}_1 / \text{Gap}_{1024}$	114	164	350	3253

**Results, II:** Subgradient Descent on the dual of Matrix Completion:

$p$	$N$	$\text{Gap}_1$	$\text{Gap}_1 / \text{Gap}_{32}$	$\text{Gap}_1 / \text{Gap}_{128}$	$\text{Gap}_1 / \text{Gap}_{1024}$	CPU, sec
2048	8192	1.81e-1	171.2	213.8	451.4	521.3
4096	16384	3.74e-1	335.4	1060.8	1287.3	1524.8
8192	16384	2.54e-1	37.8	875.8	1183.6	3644.0

**Problem:** Given noisy observations  $a_{ij}, (i, j) \in \Omega$ , of  $N$  entries in a  $p \times p$  matrix, find matrix of nuclear norm  $\leq 1$  which is the best fit to the observations in the uniform norm.

**Note:**

- **LMO** for the domain  $X$  of the problem of interest (unit Nuclear norm ball in  $\mathbb{R}^{p \times p}$ ) is readily given by Power method.
- The objective admits bilinear **F-t.r.**, the domain  $Y$  of the dual problem is the proximal-friendly unit  $\ell_1$ -ball in  $\mathbb{R}^N$ .

**Worst-case efficiency:**  $\epsilon_{\text{CM}}(x^t) \leq O(1) \frac{\ln(N)[1 + \max_{i,j} |a_{ij}|]}{\sqrt{t}}$

**Results, I:** Restricted Memory Bundle-Level algorithm on the dual of a low size ( $p = 512, N = 512$ ) Matrix Completion:

Memory depth	1	33	65	129
$\text{Gap}_1 / \text{Gap}_{1024}$	114	164	350	3253

**Results, II:** Subgradient Descent on the dual of Matrix Completion:

$p$	$N$	$\text{Gap}_1$	$\text{Gap}_1 / \text{Gap}_{32}$	$\text{Gap}_1 / \text{Gap}_{128}$	$\text{Gap}_1 / \text{Gap}_{1024}$	CPU, sec
2048	8192	1.81e-1	171.2	213.8	451.4	521.3
4096	16384	3.74e-1	335.4	1060.8	1287.3	1524.8
8192	16384	2.54e-1	37.8	875.8	1183.6	3644.0

**Problem:** Given a noisy observation  $b \in \mathbb{R}^{q \times q}$  of the linear image  $u \mapsto \mathcal{A}u : \mathbb{R}^{p \times p} \rightarrow \mathbb{R}^{q \times q}$ , find  $p \times p$  matrix of nuclear norm  $\leq 1$  which is the best fit to the observation in the spectral norm.

**Note:**

- The problem reduces to VI with affine monotone operator  $\Phi$  on LMO-represented domain (product of two nuclear norm balls)
- $\Phi$  admits explicit F.-t.r. with affine  $G(\cdot)$  and proximal-friendly dual domain (product of two Euclidean balls)

**Worst-case efficiency:**  $\epsilon_{\text{CM}}(v^t|f, U_p) \leq \epsilon_{\text{SP}}([v^t; w^t]|g, U_p, U_q) \leq O(1)L[A]/\sqrt{t}$

- $L[A]$ : Lipschitz constant of  $u \mapsto \mathcal{A}u$  w.r.t. Frobenius norms.

**Accuracy of primal solutions** yielded by Subgradient Descent applied to the dual VI:

		Iteration count $t$						
		1	65	129	257	385	449	512
$p = 8192$ $q = 4048$	$\epsilon_{\text{SP}}^t$	0.1193	0.0232	0.0134	0.0054	0.0035	0.0034	0.0034
	$\epsilon_{\text{SP}}^t / \epsilon_{\text{SP}}^1$	1.00	5.14	8.90	22.00	33.93	34.85	35.14
	cpu, sec	6.5	289.9	683.8	1816.0	3648.3	4572.2	5490.8
$p = 16384$ $q = 8192$	$\epsilon_{\text{SP}}^t$	0.1196	0.0214	0.0146	0.0085			
	$\epsilon_{\text{SP}}^t / \epsilon_{\text{SP}}^1$	1.00	5.60	8.19	14.01			
	cpu, sec	21.7	920.4	2050.2	4902.2			

Platform: 4 x 3.40 GHz desktop with 16 GB RAM, 64 bit Windows 7 OS.

**Note:** The design dimension of the largest instance is  $2^{28} = 268\,435\,456$ .

**Problem:** Given a noisy observation  $b \in \mathbb{R}^{q \times q}$  of the linear image  $u \mapsto \mathcal{A}u : \mathbb{R}^{p \times p} \rightarrow \mathbb{R}^{q \times q}$ , find  $p \times p$  matrix of nuclear norm  $\leq 1$  which is the best fit to the observation in the spectral norm.

**Note:**

- The problem reduces to **VI** with affine monotone operator  $\Phi$  on **LMO**-represented domain (product of two nuclear norm balls)
- $\Phi$  admits explicit **F-t.r.** with affine  $G(\cdot)$  and proximal-friendly dual domain (product of two Euclidean balls)

**Worst-case efficiency:**  $\epsilon_{\text{CM}}(v^t | f, U_p) \leq \epsilon_{\text{SP}}([v^t; w^t] | g, U_p, U_q) \leq O(1)L[A]/\sqrt{t}$

- $L[A]$ : Lipschitz constant of  $u \mapsto \mathcal{A}u$  w.r.t. Frobenius norms.

**Accuracy of primal solutions** yielded by Subgradient Descent applied to the dual **VI**:

		Iteration count $t$						
		1	65	129	257	385	449	512
$p = 8192$ $q = 4048$	$\epsilon_{\text{SP}}^t$	0.1193	0.0232	0.0134	0.0054	0.0035	0.0034	0.0034
	$\epsilon_{\text{SP}}^t / \epsilon_{\text{SP}}^1$	1.00	5.14	8.90	22.00	33.93	34.85	35.14
	cpu, sec	6.5	289.9	683.8	1816.0	3648.3	4572.2	5490.8
$p = 16384$ $q = 8192$	$\epsilon_{\text{SP}}^t$	0.1196	0.0214	0.0146	0.0085			
	$\epsilon_{\text{SP}}^t / \epsilon_{\text{SP}}^1$	1.00	5.60	8.19	14.01			
	cpu, sec	21.7	920.4	2050.2	4902.2			

Platform: 4 x 3.40 GHz desktop with 16 GB RAM, 64 bit Windows 7 OS.

**Note:** The design dimension of the largest instance is  $2^{28} = 268\,435\,456$ .

**Problem:** Given a noisy observation  $b \in \mathbb{R}^{q \times q}$  of the linear image  $u \mapsto \mathcal{A}u : \mathbb{R}^{p \times p} \rightarrow \mathbb{R}^{q \times q}$ , find  $p \times p$  matrix of nuclear norm  $\leq 1$  which is the best fit to the observation in the spectral norm.

**Note:**

- The problem reduces to **VI** with affine monotone operator  $\Phi$  on **LMO**-represented domain (product of two nuclear norm balls)
- $\Phi$  admits explicit **F-t.r.** with affine  $G(\cdot)$  and proximal-friendly dual domain (product of two Euclidean balls)

**Worst-case efficiency:**  $\epsilon_{\text{CM}}(v^t | f, U_p) \leq \epsilon_{\text{SP}}([v^t; w^t] | g, U_p, U_q) \leq O(1)L[A]/\sqrt{t}$

- $L[A]$ : Lipschitz constant of  $u \mapsto \mathcal{A}u$  w.r.t. Frobenius norms.

**Accuracy of primal solutions** yielded by Subgradient Descent applied to the dual **VI**:

		Iteration count $t$						
		1	65	129	257	385	449	512
$p = 8192$ $q = 4048$	$\epsilon_{\text{SP}}^t$	0.1193	0.0232	0.0134	0.0054	0.0035	0.0034	0.0034
	$\epsilon_{\text{SP}}^t / \epsilon_{\text{SP}}^1$	1.00	5.14	8.90	22.00	33.93	34.85	35.14
	cpu, sec	6.5	289.9	683.8	1816.0	3648.3	4572.2	5490.8
$p = 16384$ $q = 8192$	$\epsilon_{\text{SP}}^t$	0.1196	0.0214	0.0146	0.0085			
	$\epsilon_{\text{SP}}^t / \epsilon_{\text{SP}}^1$	1.00	5.60	8.19	14.01			
	cpu, sec	21.7	920.4	2050.2	4902.2			

Platform: 4 x 3.40 GHz desktop with 16 GB RAM, 64 bit Windows 7 OS.

**Note:** The design dimension of the largest instance is  $2^{28} = 268\,435\,456$ .

**Problem:** Given a noisy observation  $b \in \mathbb{R}^{q \times q}$  of the linear image  $u \mapsto \mathcal{A}u : \mathbb{R}^{p \times p} \rightarrow \mathbb{R}^{q \times q}$ , find  $p \times p$  matrix of nuclear norm  $\leq 1$  which is the best fit to the observation in the spectral norm.

**Note:**

- The problem reduces to **VI** with affine monotone operator  $\Phi$  on **LMO**-represented domain (product of two nuclear norm balls)
- $\Phi$  admits explicit **F-t.r.** with affine  $G(\cdot)$  and proximal-friendly dual domain (product of two Euclidean balls)

**Worst-case efficiency:**  $\epsilon_{\text{CM}}(v^t | f, U_p) \leq \epsilon_{\text{SP}}([v^t; w^t] | g, U_p, U_q) \leq O(1)L[\mathcal{A}]/\sqrt{t}$

- $L[\mathcal{A}]$ : Lipschitz constant of  $u \mapsto \mathcal{A}u$  w.r.t. Frobenius norms.

**Accuracy of primal solutions** yielded by Subgradient Descent applied to the dual **VI**:

		Iteration count $t$						
		1	65	129	257	385	449	512
$p = 8192$ $q = 4048$	$\epsilon_{\text{SP}}^t$	0.1193	0.0232	0.0134	0.0054	0.0035	0.0034	0.0034
	$\epsilon_{\text{SP}}^t / \epsilon_{\text{SP}}^1$	1.00	5.14	8.90	22.00	33.93	34.85	35.14
	cpu, sec	6.5	289.9	683.8	1816.0	3648.3	4572.2	5490.8
$p = 16384$ $q = 8192$	$\epsilon_{\text{SP}}^t$	0.1196	0.0214	0.0146	0.0085			
	$\epsilon_{\text{SP}}^t / \epsilon_{\text{SP}}^1$	1.00	5.60	8.19	14.01			
	cpu, sec	21.7	920.4	2050.2	4902.2			

Platform: 4 x 3.40 GHz desktop with 16 GB RAM, 64 bit Windows 7 OS.

**Note:** The design dimension of the largest instance is  $2^{28} = 268\,435\,456$



**Problem:** Given a noisy observation  $b \in \mathbb{R}^{q \times q}$  of the linear image  $u \mapsto \mathcal{A}u : \mathbb{R}^{p \times p} \rightarrow \mathbb{R}^{q \times q}$ , find  $p \times p$  matrix of nuclear norm  $\leq 1$  which is the best fit to the observation in the spectral norm.

**Note:**

- The problem reduces to **VI** with affine monotone operator  $\Phi$  on **LMO**-represented domain (product of two nuclear norm balls)
- $\Phi$  admits explicit **F-t.r.** with affine  $G(\cdot)$  and proximal-friendly dual domain (product of two Euclidean balls)

**Worst-case efficiency:**  $\epsilon_{\text{CM}}(v^t | f, U_p) \leq \epsilon_{\text{SP}}([v^t; w^t] | g, U_p, U_q) \leq O(1)L[\mathcal{A}]/\sqrt{t}$

- $L[\mathcal{A}]$ : Lipschitz constant of  $u \mapsto \mathcal{A}u$  w.r.t. Frobenius norms.

**Accuracy of primal solutions** yielded by Subgradient Descent applied to the dual **VI**:

		Iteration count $t$						
		1	65	129	257	385	449	512
$p = 8192$ $q = 4048$	$\epsilon_{\text{SP}}^t$	0.1193	0.0232	0.0134	0.0054	0.0035	0.0034	0.0034
	$\epsilon_{\text{SP}}^t / \epsilon_{\text{SP}}^1$	1.00	5.14	8.90	22.00	33.93	34.85	35.14
	cpu, sec	6.5	289.9	683.8	1816.0	3648.3	4572.2	5490.8
$p = 16384$ $q = 8192$	$\epsilon_{\text{SP}}^t$	0.1196	0.0214	0.0146	0.0085			
	$\epsilon_{\text{SP}}^t / \epsilon_{\text{SP}}^1$	1.00	5.60	8.19	14.01			
	cpu, sec	21.7	920.4	2050.2	4902.2			

Platform: 4 x 3.40 GHz desktop with 16 GB RAM, 64 bit Windows 7 OS.

**Note:** The design dimension of the largest instance is  $2^{28} = 268\,435\,456$ .

**Observation:** Structured Convex-Concave “master” SP problem

$$\text{SadVal}(M) = \min_{[u;w] \in U \times W} \max_{[v;z] \in V \times Z} \Phi(u, w; v, z) \quad (M)$$

gives rise to *primal-dual pair* of Convex-Concave SP's

$$\text{SadVal}(\text{Pr}) = \min_{u \in U} \max_{v \in V} [\phi(u; v) := \min_{w \in W} \max_{z \in Z} \Phi(u, w; v, z)] \quad (\text{Pr})$$

$$\text{SadVal}(\text{DI}) = \min_{w \in W} \max_{z \in Z} [\psi(w; z) = \min_{u \in U} \max_{v \in V} \Phi(u, w; v, z)] \quad (\text{DI})$$

with  $\text{SadVal}(M) = \text{SadVal}(\text{Pr}) = \text{SadVal}(\text{DI})$ .

**Fact** [Cox, Juditsky, Nem.'15]: Assuming  $U, W, V, Z$  compact and  $\Phi$  Lipschitz and differentiable, a  $t$ -step execution protocol  $\mathcal{I}_t$  and accuracy certificate  $\lambda$  for (Pr) induce feasible approximate solution  $[u^t; w^t; v^t; z^t]$  to (M) such that all three saddle point inaccuracies

$\epsilon_{\text{SP}}([u^t; v^t] | \phi, U, V)$ ,  $\epsilon_{\text{SP}}([w^t; z^t] | \psi, W, Z)$ ,  $\epsilon_{\text{SP}}([u^t; w^t; v^t; z^t] | \Phi, U \times W, V \times Z)$  do not exceed  $\text{Res}(\mathcal{I}_t, \lambda | U \times V)$ .

**Observation:** Structured Convex-Concave “master” SP problem

$$\text{SadVal}(M) = \min_{[u;w] \in U \times W} \max_{[v;z] \in V \times Z} \Phi(u, w; v, z) \quad (M)$$

gives rise to *primal-dual pair* of Convex-Concave SP's

$$\text{SadVal}(\text{Pr}) = \min_{u \in U} \max_{v \in V} [\phi(u; v) := \min_{w \in W} \max_{z \in Z} \Phi(u, w; v, z)] \quad (\text{Pr})$$

$$\text{SadVal}(\text{DI}) = \min_{w \in W} \max_{z \in Z} [\psi(w; z) = \min_{u \in U} \max_{v \in V} \Phi(u, w; v, z)] \quad (\text{DI})$$

with  $\text{SadVal}(M) = \text{SadVal}(\text{Pr}) = \text{SadVal}(\text{DI})$ .

**Fact** [Cox,Juditsky,Nem.'15]: Assuming  $U, W, V, Z$  compact and  $\Phi$  Lipschitz and differentiable, a  $t$ -step execution protocol  $\mathcal{I}_t$  and accuracy certificate  $\lambda$  for (Pr) induce feasible approximate solution  $[u^t; w^t; v^t; z^t]$  to (M) such that all three saddle point inaccuracies

$\epsilon_{\text{SP}}([u^t; v^t] | \phi, U, V), \epsilon_{\text{SP}}([w^t; z^t] | \psi, W, Z), \epsilon_{\text{SP}}([u^t; w^t; v^t; z^t] | \Phi, U \times W, V \times Z)$  do not exceed  $\text{Res}(\mathcal{I}_t, \lambda | U \times V)$ .

**Example 1:** To solve bilinear SP problem

$$\min_{w \in W} \max_{z \in Z} [\psi(w, z) := \langle p, w \rangle + \langle q, z \rangle + \langle z, Sw \rangle] \quad (!)$$

with LMO-represented  $W, Z$ , set the master program as

$$\begin{aligned} \min_{[u; w] \in U \times W} \max_{[v; z] \in V \times Z} [\Phi(u, w; v, z) = \langle w, p + S^T v \rangle + \langle z, q + Su \rangle - \langle v, Su \rangle] \\ [U \supset W, V \supset Z] \end{aligned}$$

$\Rightarrow (!)$  is the dual SP problem induced by (M). The induced primal is

$$\min_{u \in U} \max_{v \in V} \left[ \min_{w \in W} \langle w, p + S^T v \rangle + \max_{z \in Z} \langle z, q + Su \rangle - \langle v, Su \rangle \right] \quad (\text{Pr})$$

**Note:**

- LMO's for  $W, Z$  induce First Order oracle for (Pr)
- We can make  $U, V$  proximal-friendly

$\Rightarrow$  Solving (Pr) by proximal algorithm with certificates, we get  $O(1/\sqrt{t})$  algorithm for bilinear SP on LMO-represented domains.

**Example 1:** To solve bilinear SP problem

$$\min_{w \in W} \max_{z \in Z} [\psi(w, z) := \langle p, w \rangle + \langle q, z \rangle + \langle z, Sw \rangle] \quad (!)$$

with LMO-represented  $W, Z$ , set the master program as

$$\begin{aligned} \min_{[u; w] \in U \times W} \max_{[v; z] \in V \times Z} [\Phi(u, w; v, z) = \langle w, p + S^T v \rangle + \langle z, q + Su \rangle - \langle v, Su \rangle] \\ [U \supset W, V \supset Z] \end{aligned}$$

$\Rightarrow (!)$  is the dual SP problem induced by (M). The induced primal is

$$\min_{u \in U} \max_{v \in V} \left[ \min_{w \in W} \langle w, p + S^T v \rangle + \max_{z \in Z} \langle z, q + Su \rangle - \langle v, Su \rangle \right] \quad (\text{Pr})$$

**Note:**

- LMO's for  $W, Z$  induce First Order oracle for (Pr)
- We can make  $U, V$  proximal-friendly

$\Rightarrow$  Solving (Pr) by proximal algorithm with certificates, we get  $O(1/\sqrt{t})$  algorithm for bilinear SP on LMO-represented domains.

**Example 1:** To solve bilinear SP problem

$$\min_{w \in W} \max_{z \in Z} [\psi(w, z) := \langle p, w \rangle + \langle q, z \rangle + \langle z, Sw \rangle] \quad (!)$$

with LMO-represented  $W, Z$ , set the master program as

$$\begin{aligned} \min_{[u; w] \in U \times W} \max_{[v; z] \in V \times Z} [\Phi(u, w; v, z) = \langle w, p + S^T v \rangle + \langle z, q + Su \rangle - \langle v, Su \rangle] \\ [U \supset W, V \supset Z] \end{aligned}$$

$\Rightarrow (!)$  is the dual SP problem induced by (M). The induced primal is

$$\min_{u \in U} \max_{v \in V} \left[ \min_{w \in W} \langle w, p + S^T v \rangle + \max_{z \in Z} \langle z, q + Su \rangle - \langle v, Su \rangle \right] \quad (\text{Pr})$$

**Note:**

- LMO's for  $W, Z$  induce First Order oracle for (Pr)
- We can make  $U, V$  proximal-friendly

$\Rightarrow$  Solving (Pr) by proximal algorithm with certificates, we get  $O(1/\sqrt{t})$  algorithm for bilinear SP on LMO-represented domains.

**Example 1:** To solve bilinear SP problem

$$\min_{w \in W} \max_{z \in Z} [\psi(w, z) := \langle p, w \rangle + \langle q, z \rangle + \langle z, Sw \rangle] \quad (!)$$

with LMO-represented  $W, Z$ , set the master program as

$$\begin{aligned} \min_{[u, w] \in U \times W} \max_{[v, z] \in V \times Z} [\Phi(u, w; v, z) = \langle w, p + S^T v \rangle + \langle z, q + Su \rangle - \langle v, Su \rangle] \\ [U \supset W, V \supset Z] \end{aligned}$$

$\Rightarrow (!)$  is the dual SP problem induced by (M). The induced primal is

$$\min_{u \in U} \max_{v \in V} \left[ \min_{w \in W} \langle w, p + S^T v \rangle + \max_{z \in Z} \langle z, q + Su \rangle - \langle v, Su \rangle \right] \quad (\text{Pr})$$

**Note:**

- LMO's for  $W, Z$  induce First Order oracle for (Pr)
- We can make  $U, V$  proximal-friendly

$\Rightarrow$  Solving (Pr) by proximal algorithm with certificates, we get  $O(1/\sqrt{t})$  algorithm for bilinear SP on LMO-represented domains.

**Example 2:** To solve matrix game

$$\min_{w \in \Delta_M} \max_{z \in \Delta_N} [\psi(w, z) = \langle Az, Dw \rangle] \quad (!)$$

$$[\Delta_k = \{x \in \mathbb{R}_+^k : \sum_i x_i = 1\}, D : K \times M, A : K \times N]$$

set the master program as

$$\min_{[u; w] \in U \times \Delta_M} \max_{[v; z] \in V \times \Delta_N} [\Phi(u, w; v, z) = \langle u, Az \rangle + \langle v, Dw \rangle - \langle v, u \rangle] \quad (M)$$

$$[U \supset DW, V \supset AZ]$$

$\Rightarrow (!)$  is the dual SP problem induced by (M). The induced primal is

$$\min_{u \in U \subset \mathbb{R}^K} \max_{v \in V \subset \mathbb{R}^K} [\phi(u, v) = \text{Max}(A^T u) - \text{Min}(D^T v) - \langle u, v \rangle] \quad (\text{Pr})$$

**Note:** In some applications,

- the dimension  $K$  of (Pr) is small
- matrices  $A$  and  $D$  are *well organized*: it is easy to find a column making the largest inner product with a given  $x \in \mathbb{R}^K$ .

$\Rightarrow$  Applying to (Pr) the Ellipsoid algorithm, we solve (!) to high accuracy with effort independent of the sizes  $M, N$  of (!)



**Example 2:** To solve matrix game

$$\min_{w \in \Delta_M} \max_{z \in \Delta_N} [\psi(w, z) = \langle Az, Dw \rangle] \quad (!)$$

$$[\Delta_k = \{x \in \mathbb{R}_+^k : \sum_i x_i = 1\}, D : K \times M, A : K \times N]$$

set the master program as

$$\min_{[u; w] \in U \times \Delta_M} \max_{[v; z] \in V \times \Delta_N} [\Phi(u, w; v, z) = \langle u, Az \rangle + \langle v, Dw \rangle - \langle v, u \rangle] \quad (M)$$

$$[U \supset DW, V \supset AZ]$$

$\Rightarrow (!)$  is the dual SP problem induced by (M). The induced primal is

$$\min_{u \in U \subset \mathbb{R}^K} \max_{v \in V \subset \mathbb{R}^K} [\phi(u, v) = \text{Max}(A^T u) - \text{Min}(D^T v) - \langle u, v \rangle] \quad (\text{Pr})$$

**Note:** In some applications,

- the dimension  $K$  of (Pr) is small
- matrices  $A$  and  $D$  are *well organized*: it is easy to find a column making the largest inner product with a given  $x \in \mathbb{R}^K$ .

$\Rightarrow$  Applying to (Pr) the Ellipsoid algorithm, we solve (!) to high accuracy with effort independent of the sizes  $M, N$  of (!)

**Example 2:** To solve matrix game

$$\min_{w \in \Delta_M} \max_{z \in \Delta_N} [\psi(w, z) = \langle Az, Dw \rangle] \quad (!)$$

$$[\Delta_k = \{x \in \mathbb{R}_+^k : \sum_i x_i = 1\}, D : K \times M, A : K \times N]$$

set the master program as

$$\min_{[u; w] \in U \times \Delta_M} \max_{[v; z] \in V \times \Delta_N} [\Phi(u, w; v, z) = \langle u, Az \rangle + \langle v, Dw \rangle - \langle v, u \rangle] \quad (M)$$

$$[U \supset DW, V \supset AZ]$$

$\Rightarrow (!)$  is the dual SP problem induced by (M). The induced primal is

$$\min_{u \in U \subset \mathbb{R}^K} \max_{v \in V \subset \mathbb{R}^K} [\phi(u, v) = \text{Max}(A^T u) - \text{Min}(D^T v) - \langle u, v \rangle] \quad (\text{Pr})$$

**Note:** In some applications,

- the dimension  $K$  of (Pr) is small
- matrices  $A$  and  $D$  are *well organized*: it is easy to find a column making the largest inner product with a given  $x \in \mathbb{R}^K$ .

$\Rightarrow$  Applying to (Pr) the Ellipsoid algorithm, we solve (!) to high accuracy with effort independent of the sizes  $M, N$  of (!)

**Example 2:** To solve matrix game

$$\min_{w \in \Delta_M} \max_{z \in \Delta_N} [\psi(w, z) = \langle Az, Dw \rangle] \quad (!)$$

$$[\Delta_k = \{x \in \mathbb{R}_+^k : \sum_i x_i = 1\}, D : K \times M, A : K \times N]$$

set the master program as

$$\min_{[u; w] \in U \times \Delta_M} \max_{[v; z] \in V \times \Delta_N} [\Phi(u, w; v, z) = \langle u, Az \rangle + \langle v, Dw \rangle - \langle v, u \rangle] \quad (M)$$

$$[U \supset DW, V \supset AZ]$$

$\Rightarrow (!)$  is the dual SP problem induced by (M). The induced primal is

$$\min_{u \in U \subset \mathbb{R}^K} \max_{v \in V \subset \mathbb{R}^K} [\phi(u, v) = \text{Max}(A^T u) - \text{Min}(D^T v) - \langle u, v \rangle] \quad (\text{Pr})$$

**Note:** In some applications,

- the dimension  $K$  of (Pr) is small
- matrices  $A$  and  $D$  are *well organized*: it is easy to find a column making the largest inner product with a given  $x \in \mathbb{R}^K$ .

$\Rightarrow$  Applying to (Pr) the Ellipsoid algorithm, we solve (!) to high accuracy with effort independent of the sizes  $M, N$  of (!)