ML in PL CONFERENCE 2026

8 - 11 OCTOBER / COPERNICUS SCIENCE CENTRE, WARSAW, POLAND


KAUST
King Abdullah University of Science and Technology

How to Think About Optimization

The Broximal Point Method and Its Impact on Optimization Theory and Machine Learning

Peter Richtárik
King Abdullah University of Science and Technology
Kingdom of Saudi Arabia







KAUST
King Abdullah University of Science and Technology

Optimization & Machine Learning Lab



Optimization & Machine Learning Lab

KAUST
King Abdullah University of Science and Technology

KAUST (King Abdullah University of Science and ...
Oct 23, 2025 • Edited

#KAUST PhD student Kaja Gruntkowska has been awarded a Google PhD Fellowship, earning the 2025 fellowship in Algorithms & Optimization.

Guided by her advisor, Prof. Peter Richtarik, Kaja's innovative research bridges mathematics, statistics, and AI, advancing both the theory and practice of optimization for machine learning to make AI training faster, more cost-effective, and more resource-efficient.

She shares: "My goal is to bridge rigorous mathematical theory with practical solutions that scale to today's trillion-parameter AI models. This fellowship will help turn these ideas into tools that keep AI breakthroughs within reach."

At KAUST, we are proud to empower the next generation of AI leaders and drive innovation from Saudi Arabia to the world.



695 17 6



Optimization & Machine Learning Lab

KAUST
King Abdullah University of Science and Technology

KAUST (King Abdullah University of Science and ...
Oct 23, 2025 · Edited

#KAUST PhD student Kaja Gruntkowska has been awarded a Google PhD Fellowship, earning the 2025 fellowship in Algorithms & Optimization.

Guided by her advisor, Prof. Peter Richtarik, Kaja's innovative research bridges mathematics, statistics, and AI, advancing both the theory and practice of optimization for machine learning to make AI training faster, more cost-effective, and more resource-efficient.

She shares: "My goal is to bridge rigorous mathematical theory with practical solutions that scale to today's trillion-parameter AI models. This fellowship will help turn these ideas into tools that keep AI breakthroughs within reach."

At KAUST, we are proud to empower the next generation of AI leaders and drive innovation from Saudi Arabia to the world.



695 17 6



Optimization & Machine Learning Lab

KAUST
King Abdullah University of Science and Technology

KAUST (King Abdullah University of Science and ...
Oct 23, 2025 · Edited

#KAUST PhD student Kaja Gruntkowska has been awarded a Google PhD Fellowship, earning the 2025 fellowship in Algorithms & Optimization.

Guided by her advisor, Prof. Peter Richtarik, Kaja's innovative research bridges mathematics, statistics, and AI, advancing both the theory and practice of optimization for machine learning to make AI training faster, more cost-effective, and more resource-efficient.

She shares: "My goal is to bridge rigorous mathematical theory with practical solutions that scale to today's trillion-parameter AI models. This fellowship will help turn these ideas into tools that keep AI breakthroughs within reach."


At KAUST, we are proud to empower the next generation of AI leaders and drive innovation from Saudi Arabia to the world.



695 17 6




A Few Random Photos of KAUST



Optimization

$$\min_{x \in \mathbb{R}^d} f(x)$$

Optimization




x = weights of a neural network

GPT-3: $d \approx 1.75 \cdot 10^{11}$

$$\min_{x \in \mathbb{R}^d} f(x)$$

number of trainable parameters

Optimization




x = weights of a neural network

GPT-3: $d \approx 1.75 \cdot 10^{11}$


$$\min_{x \in \mathbb{R}^d} f(x)$$

number of trainable parameters

$$f : \mathbb{R}^d \to \mathbb{R} \cup \{+\infty\}$$




Optimization



x = weights of a neural network

GPT-3: $d \approx 1.75 \cdot 10^{11}$

$$\min_{x \in \mathbb{R}^d} f(x)$$




number of trainable parameters

$$f : \mathbb{R}^d \to \mathbb{R} \cup \{+\infty\}$$

f is proper

Optimization




x = weights of a neural network

GPT-3: $d \approx 1.75 \cdot 10^{11}$

$$\min_{x \in \mathbb{R}^d} f(x)$$

number of trainable parameters



$$f : \mathbb{R}^d \to \mathbb{R} \cup \{+\infty\}$$


f is proper

f is closed (= lower semi-continuous)


Outline

$$\min_{x \in \mathbb{R}^d} f(x) \qquad d = 2$$

- Part 1 BPM: Convex Regime
- Part 2 BPM: Convergence Theory
- Part 3 BPM: Nonconvex Regime
- Part 4 BPM: Practical Approximations
- Part 5 BPM: Four Formulations



BPM = Broximal Point Method


arXiv:2003.08078v1 [math.OC] 18 Mar 2020

Acceleration with a Ball Optimization Oracle

Yair Carmon* Arun Jambulapati* Qijia Jiang* Yujia Jin* Yin Tat Lee†
Aaron Sidford* Kevin Tian*

Abstract

Consider an oracle which takes a point x and returns the minimizer of a convex function f in an ℓ_2 ball of radius r around x. It is straightforward to show that roughly $r^{-1}\log\frac{1}{\epsilon}$ calls to the oracle suffice to find an ϵ-approximate minimizer of f in an ℓ_2 unit ball. Perhaps surprisingly, this is not optimal: we design an accelerated algorithm which attains an ϵ-approximate minimizer with roughly $r^{-2/3}\log\frac{1}{\epsilon}$ oracle queries, and give a matching lower bound. Further, we implement ball optimization oracles for functions with locally stable Hessians using a variant of Newton's method. The resulting algorithm applies to a number of problems of practical and theoretical import, improving upon previous results for logistic and ℓ_∞ regression and achieving guarantees comparable to the state-of-the-art for ℓ_p regression.

1 Introduction

We study unconstrained minimization of a smooth convex objective $f : \mathbb{R}^d \to \mathbb{R}$, which we access through a *ball optimization oracle* $\mathcal{O}_{\mathrm{ball}}$, that when queried at any point x, returns the minimizer[1] of f restricted a ball of radius r around x, i.e.,

$$\mathcal{O}_{\mathrm{ball}}(x) = \operatorname*{arg\,min}_{x' \text{ s.t. } \|x'-x\|\le r} f(x').$$

Such oracles underlie trust region methods [12] and, as we demonstrate via applications, encapsulate problems with local stability. Iterating $x_{k+1} \leftarrow \mathcal{O}_{\mathrm{ball}}(x_k)$ minimizes f in $\widetilde{O}(R/r)$ iterations (see Appendix A), where R is the initial distance to the minimizer, $x^\star$, and $\widetilde{O}(\cdot)$ hides polylogarithmic factors in problem parameters, including the desired accuracy.

Given the fundamental geometric nature of the ball optimization abstraction, the central question motivating our work is whether it is possible to improve upon this $\widetilde{O}(R/r)$ query complexity. It is natural to conjecture that the answer is negative: we require R/r oracle calls to observe the entire line from x_0 to the optimum, and therefore finding a solution using less queries would require jumping into completely unobserved regions. Nevertheless, we prove that the optimal query complexity scales as $(R/r)^{2/3}$. This result has positive implications for the complexity for several key regression tasks, for which we can efficiently implement the ball optimization oracles.

1.1 Our contributions

Here we overview the main contributions of our paper: accelerating ball optimization oracles (with a matching lower bound), implementing them under Hessian stability, and applying the resulting techniques to regression problems.

[*]Stanford University. {yairc,jsidford}@stanford.edu.
[†]University of Washington. yintat@uw.edu.
[1]In the introduction we discuss exact oracles for simplicity, but our results account for inexactness.


arXiv:2502.02002v1 [math.OC] 4 Feb 2025

The Ball-Proximal (=“Broximal”) Point Method:
a New Algorithm, Convergence Theory, and Applications

Kaja Gruntkowska[1] Hanmin Li[1] Aadi Rane*[1,2] Peter Richtárik[1]

Abstract

Non-smooth and non-convex global optimization poses significant challenges across various applications, where standard gradient-based methods often struggle. We propose the *Ball-Proximal Point Method*, *Broximal Point Method*, or *Ball Point Method* (BPM) for short – a novel algorithmic framework inspired by the classical Proximal Point Method (PPM) (Rockafellar, 1976), which, as we show, sheds new light on several foundational optimization paradigms and phenomena, including non-convex and non-smooth optimization, acceleration, smoothing, adaptive stepsize selection, and trust-region methods. At the core of BPM lies the *ball-proximal (“broximal”)* operator, which arises from the classical proximal operator by replacing the quadratic distance penalty by a ball constraint. Surprisingly, and in sharp contrast with the sublinear rate of PPM in the nonsmooth convex regime, we prove that BPM converges *linearly* and in a *finite* number of steps in the same regime. Furthermore, by introducing the concept of ball-convexity, we prove that BPM retains the same global convergence guarantees under weaker assumptions, making it a powerful tool for a broader class of potentially non-convex optimization problems. Just like PPM plays the role of a conceptual method inspiring the development of practically efficient algorithms and algorithmic elements, e.g., gradient descent, adaptive step sizes, acceleration (Ahn & Sra, 2020), and “W” in AdamW (Zhuang et al., 2022), we believe that BPM should be understood in the same manner: as a blueprint and inspiration for further development.

[1]King Abdullah University of Science and Technology, Thuwal, Saudi Arabia [2]University of California, Berkeley, USA.
*The work of Aadi Rane was conducted during a VSRP internship at KAUST.

1. Introduction

The minimization of non-convex functions is a fundamental challenge across many fields, including machine learning, optimization, applied mathematics, signal processing and operations research. Solving such problems is integral to most machine learning algorithms arising in both training and inference, where non-convex objectives or constraints are often necessary to capture complex prediction tasks.

1.1. Global nonconvex optimization

In this paper, we propose a new meta-algorithm (see Section 1.3) capable of finding *global* minimizers for a specific (new) class of non-convex functions. In particular, we introduce an algorithmic framework designed to solve the (potentially non-convex) optimization problem

$$\min_{x \in \mathbb{R}^d} f(x), \tag{1}$$

where $f : \mathbb{R}^d \mapsto \mathbb{R} \cup \{+\infty\}$ is assumed to be proper (which means that the set $\mathrm{dom} f := \{x \in \mathbb{R}^d : f(x) < +\infty\}$ is nonempty), closed, and have at least one minimizer. We let $\mathcal{X}_f$ be the set of all minimizers of f, and $f_\star := \min_x f(x)$.

1.2. The ball-proximal operator


A key inspiration for our method stems from the well-known Proximal Point Method (PPM) (Rockafellar, 1976), which iteratively adds a quadratic penalty term to the objective (see Section 3 for more details) and solves a modified subproblem at each step. Building on this idea, we introduce the *ball-proximal (“broximal”) operator*:

Definition 1.1 (Ball-Proximal Operator)**.** *The ball-proximal (“broximal”) operator* with radius $t > 0$ associated with a function $f : \mathbb{R}^d \mapsto \mathbb{R} \cup \{+\infty\}$ is given by

$$\mathrm{brox}_f^t(x) := \operatorname*{arg\,min}_{z \in B_t(x)} f(z), \tag{2}$$

where $B_t(x) := \{z \in \mathbb{R}^d : \|z - x\| \le t\}$ and $\|\cdot\|$ is the standard Euclidean norm.

According to this definition, for a given input point $x \in \mathbb{R}^d$, $\mathrm{brox}_f^t(x)$ returns the minimizer(s) of f within the ball of radius t centered at x.


arXiv:2510.00823v1 [math.OC] 1 Oct 2025

NON-EUCLIDEAN BROXIMAL POINT METHOD:
A BLUEPRINT FOR GEOMETRY-AWARE OPTIMIZATION

Kaja Gruntkowska
KAUST*
Center of Excellence for Generative AI
Thuwal, Saudi Arabia

Peter Richtárik
KAUST*
Center of Excellence for Generative AI
Thuwal, Saudi Arabia

ABSTRACT

The recently proposed Broximal Point Method (BPM) (Gruntkowska et al., 2025) offers an idealized optimization framework based on iteratively minimizing the objective function over norm balls centered at the current iterate. It enjoys striking global convergence guarantees, converging linearly and in a finite number of steps for proper, closed and convex functions. However, its theoretical analysis has so far been confined to the *Euclidean* geometry. At the same time, emerging trends in deep learning optimization, exemplified by algorithms such as Muon [Jordan et al., 2024] and Scion [Pethick et al., 2025], demonstrate the practical advantages of minimizing over balls defined via *non-Euclidean* norms which better align with the underlying geometry of the associated loss landscapes. In this note, we ask whether the convergence theory of BPM can be extended to this more general, non-Euclidean setting. We give a positive answer, showing that most of the elegant guarantees of the original method carry over to arbitrary norm geometries. Along the way, we clarify which properties are preserved and which necessarily break down when leaving the Euclidean realm. Our analysis positions Non-Euclidean BPM as a conceptual blueprint for understanding a broad class of geometry-aware optimization algorithms, shedding light on the principles behind their practical effectiveness.


1 Introduction

Optimization stands as a cornerstone of modern machine learning, powering the success of virtually everything, from deep neural networks to state-of-the-art large language models. As models grow ever larger and more complex, the demand for better optimization algorithms evolves in tandem. What is needed now is a new class of methods–ones that are not only computationally efficient and scalable, but also inherently robust to the nonconvex, high-dimensional landscapes that characterize contemporary deep learning.

The field of deep learning optimization has long relied on Adam and related algorithms [Kingma and Ba, 2014, Loshchilov and Hutter, 2017], whose core innovation lies in adaptive moment estimation. While these methods have achieved significant empirical success, their theoretical behavior, particularly in nonconvex landscapes, remains only partially understood. Recently, however, a new class of optimization methods has begun to challenge this long-standing dominance. Algorithms such as Muon [Jordan et al., 2024], Scion [Pethick et al., 2025] and Gluon [Riabinin et al., 2025] break away from the adaptive moment paradigm. Instead, they adopt a different principle: structured updates derived by minimizing a linear approximation of the loss function over a carefully chosen norm ball. Crucially, these norm balls are *non-Euclidean*, aiming to reflect the intrinsic geometry of the optimization problem (see Section 3.1). Operating at the individual layer level, these algorithms combine simplicity, scalability, and promising empirical performance, with several studies suggesting their potential to outperform Adam(W) in large-scale deep learning tasks [Liu et al., 2025, Pethick et al., 2025, Shah et al., 2025, Thérien et al., 2025, Yvet et al., 2025].

While Muon and Scion—both of which utilize spectral norm balls—are among the most prominent and actively studied examples in this emerging line of work, the underlying principle extends well beyond these specific instances. By changing the geometry through altering the norm, one can recover a range of familiar optimization methods. For

[1]King Abdullah University of Science and Technology


arXiv:2604.13483v1 [math.OC] 15 Apr 2026

Broximal Alignment for Global Non-Convex Optimization

Kaja Gruntkowska[1] Hanmin Li[1] Xun Qian[1] Peter Richtárik[1]

Abstract

Most non-convex optimization theory is built around gradient dynamics, leaving global convergence largely unexplored. The dominant paradigm focuses on stationarity, certifying only that the gradient norm vanishes, which is often a weak proxy for actual optimization success. In practice, gradient norms can stagnate or even increase during training, and stationary points may be far from global solutions. In this work, we propose a new framework for global non-convex optimization that avoids gradient-based reasoning altogether. We revisit the Ball Proximal Point Method (BPM), a trust-region–style algorithm introduced by Gruntkowska et al. (2025), and propose a novel structural condition – *Broximal Alignment* – under which BPM provably converges to a global minimizer. Our condition requires no convexity, smoothness, or Lipschitz assumptions, and it permits multiple and disconnected global minima as well as non-optimal local minima. We show that this class generalizes standard non-convex frameworks such as quasi-convexity, star convexity, quasar convexity, and the aiming condition. Our results provide a new conceptual foundation for global non-convex optimization beyond stationarity.

1. Introduction

Non-convex optimization lies at the heart of modern machine learning (ML). Matrix factorization (Koren et al., 2009), phase retrieval (Patterson, 1934; Candes et al., 2015), representation learning, and most prominently deep neural networks (LeCun et al., 2015; Goodfellow et al., 2016) all lead to objective functions with highly non-convex landscapes. Although such functions provide the expressive capacity needed to model complex learning tasks, they often pose significant challenges for optimizer designers. The underlying landscape can be highly irregular, with many poor local minima and saddle points that can trap optimization algorithms (Pascanu et al., 2014; Vidal et al., 2017; Safran & Shamir, 2018; Jin et al., 2021).

Unlike convex optimization, where global minimizers can be efficiently found (Nemirovski & Yudin, 1983; Nemirovski & Nesterov, 1985; Nesterov, 2003), we lack a comparable toolkit for non-convex problems. Indeed, many non-convex optimization tasks are NP-hard to solve (Murty & Kabadi, 1987), and a number of them are NP-hard even to approximate (Meka et al., 2008; Jain & Kar, 2017). As a result, rather than aiming for (approximate) global solutions, most theoretical results for non-convex optimization, especially those for gradient-based methods, are framed in terms of *stationarity*: convergence to points where the gradient norm is small. Although mathematically convenient, such guarantees provide limited insight into *global* optimization behavior.

1.1. Limitations of Stationarity-Based Non-convex Optimization Theory

To fix notation, we consider solving the general (potentially non-convex) optimization problem

$$\min_{x \in \mathbb{R}^d} f(x), \tag{1}$$

where $f : \mathbb{R}^d \mapsto \mathbb{R} \cup \{+\infty\}$ is proper (that is, the set $\mathrm{dom} f := \{x \in \mathbb{R}^d : f(x) < +\infty\}$ is nonempty), closed, and has at least one minimizer. We denote the set of all minimizers by $\mathcal{X}_f$ and write $f_\star := \min_x f(x)$.

The dominant theoretical paradigm for such tasks is built around *stationarity*. Convergence guarantees are typically stated in terms of the norm of the gradient, asserting that an algorithm reaches a point where $\|\nabla f(x)\|$ is small. This framework underlies much of the modern theory of stochastic gradient methods.

From the modeling perspective, however, stationarity-based guarantees are fundamentally misaligned with the goals of learning. A stationary point can be a global minimizer, a poor local minimizer, or a saddle point; first-order information alone does not distinguish among these possibilities. Therefore, a small gradient provides little guidance on whether an algorithm meaningfully solves the under-

Preprint. April 16, 2026.


arXiv:2609.35147v1 [math.OC] 28 Sep 2026

A Sharper Theory of Ball-Proximal Optimization:
Convergence and Radius Selection

Peter Richtárik Hanmin Li
King Abdullah University of Science and Technology
Thuwal 23955-6900, Saudi Arabia

28 September 2026

Abstract

We study the exact Euclidean ball-proximal point method for proper, closed, convex functions, where each iteration minimizes the objective over a ball centered at the current point. Retaining the objective gap in the decrease of squared distance yields sharper bounds on objective values, stationarity, and the symmetric Bregman distance to a minimizer. For constant radius $t > 0$ and initial distance $D_0 > 0$ to the solution set, the objective gap after K iterations is at most its initial value multiplied by $\exp(-2K^2t^2/D_0^2)$.

We characterize convergence for arbitrary positive radius sequences. If their sum diverges, the method reaches a minimizer in finitely many iterations whenever one exists; otherwise, the objective values converge to the infimum and the iterates escape every bounded set. If the radii are summable, the iterates converge to a possibly nonoptimal point. Self-contraction gives finite trajectory length for bounded iterates and a constant-radius termination bound $O_d(1 + D_0/t)$, whose implicit constant depends only on the dimension. A polyhedral family shows that the dimension-independent quadratic bound remains asymptotically sharp.

We also identify a minimum successful geometric decay factor and analyze adaptive radius rules based on subgradients or objective gaps, epigraph reformulation, and relaxed updates. Together, these results strengthen the foundations and convergence guarantees of the method without assuming smoothness or strong convexity.

Keywords: ball-proximal point method; convex optimization; finite termination; self-contracted sequences; radius selection; oracle complexity.
Mathematics Subject Classification (2020): Primary 90C25; Secondary 90C60, 65K05.

1 Introduction

We study the convex optimization problem

$$\min_{x \in \mathbb{R}^d} f(x), \tag{1}$$

where $f : \mathbb{R}^d \to \mathbb{R} \cup \{+\infty\}$ is proper, closed, and convex. Unless stated otherwise, we assume neither that the infimum is attained nor that f is smooth. We focus on the exact ball-oracle iteration considered by Carmon et al. (2020, Appendix A) and subsequently developed as the ball-proximal (broximal) point framework by Gruntkowska et al. (2025). Starting from $x^0 \in \mathrm{dom}\, f$, the ball-proximal point method (BPM) minimizes f over a Euclidean ball centered at the current point:

$$x^{k+1} \in \mathrm{brox}_f^t(x^k) := \operatorname*{arg\,min}_{u \in B(x^k, t)} f(u), \qquad t_k > 0, \tag{BPM}$$

Part 1
Broximal Point Method: Convex Regime



Kaja Gruntkowska, Hanmin Li, Aadi Rane, P.R.
The ball-proximal (="broximal") point method: a new algorithm, convergence theory, and applications
arXiv:2502.02002, 2025



Kaja Gruntkowska, P.R.
Non-Euclidean broximal point method: a blueprint for geometry-aware optimization
arXiv:2510.00823, 2025

BPM with ℓ_2 Balls


$$x_{k+1} \in \operatorname{Argmin} \{ f(y) \ : \ \|y - x_k\|_2 \leq t_k \}$$

Example: $f(x) = \frac{1}{2}\left(|\langle a, x\rangle| - 2\right)_+^2 + 8\langle b, x\rangle^2, \qquad a = (\cos 20^\circ, \sin 20^\circ), \ \ b = (-\sin 20^\circ, \cos 20^\circ)$

BPM with ℓ_2 Balls

$$x_{k+1} \in \operatorname{Argmin}\{f(y) \ : \ \|y - x_k\|_2 \le t_k\}$$

Example: $f(x) = \frac{1}{2}\left(|\langle a, x\rangle| - 2\right)_+^2 + 8\langle b, x\rangle^2, \qquad a = (\cos 20^\circ, \sin 20^\circ), \ \ b = (-\sin 20^\circ, \cos 20^\circ)$



level sets of f
through the iterates
(darker = lower f)


BPM with ℓ_2 Balls

$$x_{k+1} \in \operatorname{Argmin}\{f(y) \;:\; \|y - x_k\|_2 \le t_k\}$$

Example: $f(x) = \frac{1}{2}\left(|\langle a, x\rangle| - 2\right)_+^2 + 8\langle b, x\rangle^2, \qquad a = (\cos 20^\circ, \sin 20^\circ), \;\; b = (-\sin 20^\circ, \cos 20^\circ)$

$\operatorname{Argmin} f = [-2a,\ 2a]$

$f_\star := \min f = 0$



level sets of f
through the iterates
(darker = lower f)

BPM with ℓ_2 Balls

$$x_{k+1} \in \operatorname{Argmin}\{f(y) \ : \ \|y - x_k\|_2 \le t_k\}$$

Example: $f(x) = \frac{1}{2}\left(|\langle a, x\rangle| - 2\right)_+^2 + 8\langle b, x\rangle^2, \qquad a = (\cos 20^\circ, \sin 20^\circ), \ \ b = (-\sin 20^\circ, \cos 20^\circ)$

BPM run

$d_k = \operatorname{dist}_2(x_k, \operatorname{Argmin} f)$


$f_k = f(x_k) - f_\star$

$g_k = \|\nabla f(x_k)\|_2$

$d_0 = 5.06 \quad f_0 = 35.78 \quad g_0 = 28.40$

$\operatorname{Argmin} f \ = \ [-2a, \ 2a]$

$f_\star := \min f \ = \ 0$



BPM with ℓ_2 Balls

$$x_{k+1} \in \operatorname{Argmin}\{f(y) \;:\; \|y - x_k\|_2 \le t_k\}$$

Example: $f(x) = \frac{1}{2}\left(|\langle a, x\rangle| - 2\right)_+^2 + 8\langle b, x\rangle^2, \qquad a = (\cos 20^\circ, \sin 20^\circ), \;\; b = (-\sin 20^\circ, \cos 20^\circ)$

BPM run

$d_k = \operatorname{dist}_2(x_k, \operatorname{Argmin} f)$


$f_k = f(x_k) - f_\star$

$g_k = \|\nabla f(x_k)\|_2$

$d_0 = 5.06 \quad f_0 = 35.78 \quad g_0 = 28.40$

$\operatorname{Argmin} f = [-2a,\ 2a]$

$f_\star := \min f = 0$



level sets of f
through the iterates
(darker = lower f)

BPM with ℓ_2 Balls

$$x_{k+1} \in \operatorname{Argmin}\{f(y) \;:\; \|y - x_k\|_2 \le t_k\}$$

Example: $f(x) = \frac{1}{2}\left(|\langle a, x\rangle| - 2\right)_+^2 + 8\langle b, x\rangle^2, \qquad a = (\cos 20^\circ, \sin 20^\circ), \;\; b = (-\sin 20^\circ, \cos 20^\circ)$

BPM run

$d_k = \operatorname{dist}_2(x_k, \operatorname{Argmin} f)$


$f_k = f(x_k) - f_\star$

$g_k = \|\nabla f(x_k)\|_2$

	$d_0 = 5.06$	$f_0 = 35.78$	$g_0 = 28.40$
$t_0 = 1.0$	$d_1 = 4.50$	$f_1 = 14.97$	$g_1 = 13.61$

$\operatorname{Argmin} f = [-2a,\ 2a]$

$f_\star := \min f = 0$



level sets of f through the iterates (darker = lower f)

BPM with ℓ_2 Balls

$$x_{k+1} \in \operatorname{Argmin}\{f(y) \;:\; \|y - x_k\|_2 \le t_k\}$$

Example: $f(x) = \frac{1}{2}\left(|\langle a, x\rangle| - 2\right)_+^2 + 8\langle b, x\rangle^2, \qquad a = (\cos 20^\circ, \sin 20^\circ), \;\; b = (-\sin 20^\circ, \cos 20^\circ)$

BPM run

$d_k = \operatorname{dist}_2(x_k, \operatorname{Argmin} f)$


$f_k = f(x_k) - f_\star$

$g_k = \|\nabla f(x_k)\|_2$

	$d_0 = 5.06$	$f_0 = 35.78$	$g_0 = 28.40$
$t_0 = 1.0$	$d_1 = 4.50$	$f_1 = 14.97$	$g_1 = 13.61$

$\operatorname{Argmin} f = [-2a,\ 2a]$

$f_\star := \min f = 0$



level sets of f
through the iterates
(darker = lower f)

BPM with ℓ_2 Balls

$$x_{k+1} \in \operatorname{Argmin}\{f(y) \;:\; \|y - x_k\|_2 \le t_k\}$$

Example: $f(x) = \frac{1}{2}\left(|\langle a, x\rangle| - 2\right)_+^2 + 8\langle b, x\rangle^2, \qquad a = (\cos 20^\circ, \sin 20^\circ), \;\; b = (-\sin 20^\circ, \cos 20^\circ)$

BPM run

$d_k = \operatorname{dist}_2(x_k, \operatorname{Argmin} f)$


$f_k = f(x_k) - f_\star$

$g_k = \|\nabla f(x_k)\|_2$

	$d_0 = 5.06$	$f_0 = 35.78$	$g_0 = 28.40$
$t_0 = 1.0$	$d_1 = 4.50$	$f_1 = 14.97$	$g_1 = 13.61$
$t_1 = 1.3$	$d_2 = 3.32$	$f_2 = 5.62$	$g_2 = 3.89$

$\operatorname{Argmin} f = [-2a,\ 2a]$

$f_\star := \min f = 0$



level sets of f through the iterates (darker = lower f)

BPM with ℓ_2 Balls

$$x_{k+1} \in \operatorname{Argmin}\{f(y) \ : \ \|y - x_k\|_2 \le t_k\}$$

Example: $f(x) = \frac{1}{2}\left(|\langle a, x\rangle| - 2\right)_+^2 + 8\langle b, x\rangle^2, \qquad a = (\cos 20^\circ, \sin 20^\circ), \ \ b = (-\sin 20^\circ, \cos 20^\circ)$

BPM run

$d_k = \operatorname{dist}_2(x_k, \operatorname{Argmin} f)$


$f_k = f(x_k) - f_\star$

$g_k = \|\nabla f(x_k)\|_2$

	$d_0 = 5.06$	$f_0 = 35.78$	$g_0 = 28.40$
$t_0 = 1.0$	$d_1 = 4.50$	$f_1 = 14.97$	$g_1 = 13.61$
$t_1 = 1.3$	$d_2 = 3.32$	$f_2 = 5.62$	$g_2 = 3.89$

$\operatorname{Argmin} f = [-2a,\ 2a]$

$f_\star := \min f = 0$



BPM with ℓ_2 Balls

$$x_{k+1} \in \operatorname{Argmin}\{f(y) \;:\; \|y - x_k\|_2 \le t_k\}$$

Example: $f(x) = \frac{1}{2}\left(|\langle a, x\rangle| - 2\right)_+^2 + 8\langle b, x\rangle^2, \qquad a = (\cos 20^\circ, \sin 20^\circ), \;\; b = (-\sin 20^\circ, \cos 20^\circ)$

BPM run

$d_k = \operatorname{dist}_2(x_k, \operatorname{Argmin} f)$


$f_k = f(x_k) - f_\star$

$g_k = \|\nabla f(x_k)\|_2$

	$d_0 = 5.06$	$f_0 = 35.78$	$g_0 = 28.40$
$t_0 = 1.0$	$d_1 = 4.50$	$f_1 = 14.97$	$g_1 = 13.61$
$t_1 = 1.3$	$d_2 = 3.32$	$f_2 = 5.62$	$g_2 = 3.89$
$t_2 = 1.6$	$d_3 = 1.72$	$f_3 = 1.48$	$g_3 = 1.72$

$\operatorname{Argmin} f = [-2a,\ 2a]$

$f_\star := \min f = 0$



BPM with ℓ_2 Balls

$$x_{k+1} \in \operatorname{Argmin}\{f(y) \;:\; \|y - x_k\|_2 \le t_k\}$$

Example: $f(x) = \frac{1}{2}\left(|\langle a, x\rangle| - 2\right)_+^2 + 8\langle b, x\rangle^2, \qquad a = (\cos 20^\circ, \sin 20^\circ), \;\; b = (-\sin 20^\circ, \cos 20^\circ)$

BPM run

$d_k = \operatorname{dist}_2(x_k, \operatorname{Argmin} f)$


$f_k = f(x_k) - f_\star$

$g_k = \|\nabla f(x_k)\|_2$

	$d_0 = 5.06$	$f_0 = 35.78$	$g_0 = 28.40$
$t_0 = 1.0$	$d_1 = 4.50$	$f_1 = 14.97$	$g_1 = 13.61$
$t_1 = 1.3$	$d_2 = 3.32$	$f_2 = 5.62$	$g_2 = 3.89$
$t_2 = 1.6$	$d_3 = 1.72$	$f_3 = 1.48$	$g_3 = 1.72$

$\operatorname{Argmin} f = [-2a,\ 2a]$

$f_\star := \min f = 0$



level sets of f through the iterates (darker = lower f)

BPM with ℓ_2 Balls

$$x_{k+1} \in \operatorname{Argmin}\{f(y) \;:\; \|y - x_k\|_2 \le t_k\}$$

Example: $f(x) = \frac{1}{2}\left(|\langle a, x\rangle| - 2\right)_+^2 + 8\langle b, x\rangle^2, \qquad a = (\cos 20^\circ, \sin 20^\circ),\ \ b = (-\sin 20^\circ, \cos 20^\circ)$

BPM run

$d_k = \operatorname{dist}_2(x_k, \operatorname{Argmin} f)$


$f_k = f(x_k) - f_\star$

$g_k = \|\nabla f(x_k)\|_2$

	$d_0 = 5.06$	$f_0 = 35.78$	$g_0 = 28.40$
$t_0 = 1.0$	$d_1 = 4.50$	$f_1 = 14.97$	$g_1 = 13.61$
$t_1 = 1.3$	$d_2 = 3.32$	$f_2 = 5.62$	$g_2 = 3.89$
$t_2 = 1.6$	$d_3 = 1.72$	$f_3 = 1.48$	$g_3 = 1.72$
$t_3 = 1.8$	$d_4 = 0.00$	$f_4 = 0.00$	$g_4 = 0.00$

$\operatorname{Argmin} f = [-2a,\ 2a]$

$f_\star := \min f = 0$



level sets of f through the iterates (darker = lower f)

BPM with ℓ_2 Balls

$$x_{k+1} \in \operatorname{Argmin}\{f(y) \ : \ \|y - x_k\|_2 \le t_k\}$$

Example: $f(x) = \frac{1}{2}\left(|\langle a, x\rangle| - 2\right)_+^2 + 8\langle b, x\rangle^2, \qquad a = (\cos 20^\circ, \sin 20^\circ), \ \ b = (-\sin 20^\circ, \cos 20^\circ)$

BPM run

$d_k = \operatorname{dist}_2(x_k, \operatorname{Argmin} f)$

$f_k = f(x_k) - f_\star$


$g_k = \|\nabla f(x_k)\|_2$

	$d_0 = 5.06$	$f_0 = 35.78$	$g_0 = 28.40$
$t_0 = 1.0$	$d_1 = 4.50$	$f_1 = 14.97$	$g_1 = 13.61$
$t_1 = 1.3$	$d_2 = 3.32$	$f_2 = 5.62$	$g_2 = 3.89$
$t_2 = 1.6$	$d_3 = 1.72$	$f_3 = 1.48$	$g_3 = 1.72$
$t_3 = 1.8$	$d_4 = 0.00$	$f_4 = 0.00$	$g_4 = 0.00$

d_k, f_k and g_k
all decrease at every step

$\operatorname{Argmin} f = [-2a, \ 2a]$

$f_\star := \min f = 0$



level sets of f
through the iterates
(darker = lower f)

Proximal Point vs. Broximal Point

PPM: Proximal Point Method (Martinet, 1970)

$$x_{k+1} = \operatorname*{argmin}_y \left\{ f(y) + \frac{1}{2\gamma}\|y - x_k\|_2^2 \right\}$$

$$\|x_{k+1} - x_k\|_2 = \gamma \|\nabla f(x_{k+1})\|_2 \to 0$$

f differentiable: $x_k \notin \operatorname{Argmin} f$ for every k




Proximal Point vs. Broximal Point

PPM: Proximal Point Method (Martinet, 1970)

$$x_{k+1} = \operatorname*{argmin}_y \left\{ f(y) + \frac{1}{2\gamma}\|y - x_k\|_2^2 \right\}$$

$$\|x_{k+1} - x_k\|_2 = \gamma \|\nabla f(x_{k+1})\|_2 \to 0$$

f differentiable: $x_k \notin \operatorname{Argmin} f$ for every k



Proximal Point vs. Broximal Point

PPM: Proximal Point Method (Martinet, 1970)

$$x_{k+1} = \operatorname*{argmin}_y \left\{ f(y) + \frac{1}{2\gamma}\|y - x_k\|_2^2 \right\}$$

$$\|x_{k+1} - x_k\|_2 = \gamma \|\nabla f(x_{k+1})\|_2 \;\to\; 0$$


f differentiable: $x_k \notin \operatorname{Argmin} f$ for every k

BPM: Broximal (= ball-proximal) Point Method

$$x_{k+1} \in \operatorname*{Argmin}_y \left\{ f(y) \;:\; \|y - x_k\|_2 \le t \right\}$$

$\|x_{k+1} - x_k\|_2 = t$ until the ball meets $\operatorname{Argmin} f$

exact after $\lceil d_0^2/t^2 \rceil$ steps; no condition number



- PPM, $\gamma = 0.155$
- BPM, $t = 1.4$

same x_0, same first step

after 4 steps: $d_4 = 0$ vs. $d_4 = 2.67$

$d_k := \operatorname{dist}_2(x_k, \operatorname{Argmin} f)$

Proximal Point vs. Broximal Point

PPM: Proximal Point Method (Martinet, 1970)

$$x_{k+1} = \operatorname*{argmin}_y \left\{ f(y) + \frac{1}{2\gamma}\|y - x_k\|_2^2 \right\}$$

$$\|x_{k+1} - x_k\|_2 = \gamma \|\nabla f(x_{k+1})\|_2 \;\to\; 0$$


f differentiable: $x_k \notin \operatorname{Argmin} f$ for every k

BPM: Broximal (= ball-proximal) Point Method

$$x_{k+1} \in \operatorname*{Argmin}_y \left\{ f(y) \;:\; \|y - x_k\|_2 \le t \right\}$$

$\|x_{k+1} - x_k\|_2 = t$ until the ball meets $\operatorname{Argmin} f$

exact after $\lceil d_0^2/t^2 \rceil$ steps; no condition number



- PPM, $\gamma = 0.155$
- BPM, $t = 1.4$

same x_0, same first step

after 4 steps: $d_4 = 0$ vs. $d_4 = 2.67$

$d_k := \operatorname{dist}_2(x_k, \operatorname{Argmin} f)$

Penalty → constraint: the steps no longer shrink, so BPM reaches Argmin f in finitely many steps.

BPM with ℓ_1 Balls


$$x_{k+1} \in \operatorname{Argmin} \{ f(y) \ : \ \|y - x_k\|_1 \leq t_k \}$$

Example: $f(x) = \frac{1}{2}\left(|\langle a, x\rangle| - 2\right)_+^2 + 8\langle b, x\rangle^2, \qquad a = (\cos 20^\circ, \sin 20^\circ), \ \ b = (-\sin 20^\circ, \cos 20^\circ)$

BPM with ℓ_1 Balls

$$x_{k+1} \in \operatorname{Argmin}\{f(y) \ : \ \|y - x_k\|_1 \leq t_k\}$$

Example: $f(x) = \frac{1}{2}\left(|\langle a, x\rangle| - 2\right)_+^2 + 8\langle b, x\rangle^2, \qquad a = (\cos 20^\circ, \sin 20^\circ), \ \ b = (-\sin 20^\circ, \cos 20^\circ)$



level sets of f
through the iterates
(darker = lower f)


BPM with ℓ_1 Balls

$$x_{k+1} \in \operatorname{Argmin}\{f(y) \ : \ \|y - x_k\|_1 \le t_k\}$$

Example: $f(x) = \frac{1}{2}\left(|\langle a, x\rangle| - 2\right)_+^2 + 8\langle b, x\rangle^2, \qquad a = (\cos 20^\circ, \sin 20^\circ), \ \ b = (-\sin 20^\circ, \cos 20^\circ)$

$\operatorname{Argmin} f \ = \ [-2a, \ 2a]$

$f_\star := \min f \ = \ 0$



level sets of f
through the iterates
(darker = lower f)

BPM with ℓ_1 Balls

$$x_{k+1} \in \operatorname{Argmin} \{ f(y) \ : \ \|y - x_k\|_1 \le t_k \}$$

Example: $f(x) = \frac{1}{2}\left(|\langle a, x\rangle| - 2\right)_+^2 + 8\langle b, x\rangle^2, \qquad a = (\cos 20^\circ, \sin 20^\circ), \ \ b = (-\sin 20^\circ, \cos 20^\circ)$

BPM run

$d_k = \operatorname{dist}_1(x_k, \operatorname{Argmin} f)$


$f_k = f(x_k) - f_\star$

$g_k = \|\nabla f(x_k)\|_\infty$ (dual norm)

$d_0 = 5.08 \qquad f_0 = 35.78 \quad g_0 = 24.69$

$\operatorname{Argmin} f \ = \ [-2a, \ 2a]$

$f_\star := \min f \ = \ 0$



BPM with ℓ_1 Balls

$$x_{k+1} \in \operatorname{Argmin}\{f(y) \;:\; \|y - x_k\|_1 \le t_k\}$$

Example: $f(x) = \frac{1}{2}\left(|\langle a, x\rangle| - 2\right)_+^2 + 8\langle b, x\rangle^2, \qquad a = (\cos 20^\circ, \sin 20^\circ),\ \ b = (-\sin 20^\circ, \cos 20^\circ)$

BPM run

$d_k = \operatorname{dist}_1(x_k, \operatorname{Argmin} f)$


$f_k = f(x_k) - f_\star$

$g_k = \|\nabla f(x_k)\|_\infty$ (dual norm)

$d_0 = 5.08 \qquad f_0 = 35.78 \quad g_0 = 24.69$

$\operatorname{Argmin} f = [-2a,\ 2a]$

$f_\star := \min f = 0$



level sets of f
through the iterates
(darker = lower f)

BPM with ℓ_1 Balls

$$x_{k+1} \in \operatorname{Argmin}\{f(y) \;:\; \|y - x_k\|_1 \le t_k\}$$

Example: $f(x) = \frac{1}{2}\left(|\langle a, x\rangle| - 2\right)_+^2 + 8\langle b, x\rangle^2, \qquad a = (\cos 20^\circ, \sin 20^\circ), \;\; b = (-\sin 20^\circ, \cos 20^\circ)$

BPM run

$d_k = \operatorname{dist}_1(x_k, \operatorname{Argmin} f)$

$f_k = f(x_k) - f_\star$


$g_k = \|\nabla f(x_k)\|_\infty$ (dual norm)

$d_0 = 5.08 \quad f_0 = 35.78 \quad g_0 = 24.69$

$t_0 = 1.0 \quad d_1 = 6.04 \uparrow \quad f_1 = 18.22 \quad g_1 = 10.44$

$\operatorname{Argmin} f = [-2a,\ 2a]$

$f_\star := \min f = 0$



BPM with ℓ_1 Balls

$$x_{k+1} \in \operatorname{Argmin}\{f(y) \;:\; \|y - x_k\|_1 \le t_k\}$$

Example: $f(x) = \frac{1}{2}\left(|\langle a, x\rangle| - 2\right)_+^2 + 8\langle b, x\rangle^2, \qquad a = (\cos 20^\circ, \sin 20^\circ), \;\; b = (-\sin 20^\circ, \cos 20^\circ)$

BPM run

$d_k = \operatorname{dist}_1(x_k, \operatorname{Argmin} f)$

$f_k = f(x_k) - f_\star$


$g_k = \|\nabla f(x_k)\|_\infty$ (dual norm)

$d_0 = 5.08 \qquad f_0 = 35.78 \quad g_0 = 24.69$

$t_0 = 1.0 \quad d_1 = 6.04 \uparrow \quad f_1 = 18.22 \quad g_1 = 10.44$

$\operatorname{Argmin} f = [-2a,\ 2a]$

$f_\star := \min f = 0$



level sets of f
through the iterates
(darker = lower f)

BPM with ℓ_1 Balls

$$x_{k+1} \in \operatorname{Argmin} \{ f(y) \ : \ \|y - x_k\|_1 \le t_k \}$$

Example: $f(x) = \frac{1}{2}\left(|\langle a, x\rangle| - 2\right)_+^2 + 8\langle b, x\rangle^2, \qquad a = (\cos 20^\circ, \sin 20^\circ), \ \ b = (-\sin 20^\circ, \cos 20^\circ)$

BPM run

$d_k = \operatorname{dist}_1(x_k, \operatorname{Argmin} f)$


$f_k = f(x_k) - f_\star$

$g_k = \|\nabla f(x_k)\|_\infty$ (dual norm)

	$d_0 = 5.08$	$f_0 = 35.78$	$g_0 = 24.69$
$t_0 = 1.0$	$d_1 = 6.04 \uparrow$	$f_1 = 18.22$	$g_1 = 10.44$
$t_1 = 1.3$	$d_2 = 4.74$	$f_2 = 8.56$	$g_2 = 5.64$

$\operatorname{Argmin} f = [-2a, \ 2a]$

$f_\star := \min f = 0$



level sets of f through the iterates (darker = lower f)

BPM with ℓ_1 Balls

$$x_{k+1} \in \operatorname{Argmin}\{f(y) \;:\; \|y - x_k\|_1 \le t_k\}$$

Example: $f(x) = \frac{1}{2}\left(|\langle a, x\rangle| - 2\right)_+^2 + 8\langle b, x\rangle^2, \qquad a = (\cos 20^\circ, \sin 20^\circ), \;\; b = (-\sin 20^\circ, \cos 20^\circ)$

BPM run

$d_k = \operatorname{dist}_1(x_k, \operatorname{Argmin} f)$


$f_k = f(x_k) - f_\star$

$g_k = \|\nabla f(x_k)\|_\infty$ (dual norm)

	$d_0 = 5.08$	$f_0 = 35.78$	$g_0 = 24.69$
$t_0 = 1.0$	$d_1 = 6.04 \uparrow$	$f_1 = 18.22$	$g_1 = 10.44$
$t_1 = 1.3$	$d_2 = 4.74$	$f_2 = 8.56$	$g_2 = 5.64$

$\operatorname{Argmin} f = [-2a,\ 2a]$

$f_\star := \min f = 0$



level sets of f through the iterates (darker = lower f)

BPM with ℓ_1 Balls

$$x_{k+1} \in \operatorname{Argmin}\{f(y) \;:\; \|y - x_k\|_1 \le t_k\}$$

Example: $f(x) = \frac{1}{2}\left(|\langle a, x\rangle| - 2\right)_+^2 + 8\langle b, x\rangle^2, \qquad a = (\cos 20^\circ, \sin 20^\circ), \;\; b = (-\sin 20^\circ, \cos 20^\circ)$

BPM run

$d_k = \operatorname{dist}_1(x_k, \operatorname{Argmin} f)$


$f_k = f(x_k) - f_\star$

$g_k = \|\nabla f(x_k)\|_\infty$ (dual norm)

	$d_0 = 5.08$	$f_0 = 35.78$	$g_0 = 24.69$
$t_0 = 1.0$	$d_1 = 6.04 \uparrow$	$f_1 = 18.22$	$g_1 = 10.44$
$t_1 = 1.3$	$d_2 = 4.74$	$f_2 = 8.56$	$g_2 = 5.64$
$t_2 = 1.6$	$d_3 = 3.14$	$f_3 = 2.96$	$g_3 = 1.89$

$\operatorname{Argmin} f = [-2a,\ 2a]$

$f_\star := \min f = 0$



level sets of f through the iterates (darker = lower f)

BPM with ℓ_1 Balls

$$x_{k+1} \in \operatorname{Argmin}\{f(y) \;:\; \|y - x_k\|_1 \le t_k\}$$

Example: $f(x) = \frac{1}{2}\left(|\langle a, x\rangle| - 2\right)_+^2 + 8\langle b, x\rangle^2, \qquad a = (\cos 20^\circ, \sin 20^\circ),\ \ b = (-\sin 20^\circ, \cos 20^\circ)$

BPM run

$d_k = \operatorname{dist}_1(x_k, \operatorname{Argmin} f)$


$f_k = f(x_k) - f_\star$

$g_k = \|\nabla f(x_k)\|_\infty$ (dual norm)

	$d_0 = 5.08$	$f_0 = 35.78$	$g_0 = 24.69$
$t_0 = 1.0$	$d_1 = 6.04$ ↑	$f_1 = 18.22$	$g_1 = 10.44$
$t_1 = 1.3$	$d_2 = 4.74$	$f_2 = 8.56$	$g_2 = 5.64$
$t_2 = 1.6$	$d_3 = 3.14$	$f_3 = 2.96$	$g_3 = 1.89$

$\operatorname{Argmin} f = [-2a,\ 2a]$

$f_\star := \min f = 0$



level sets of f through the iterates (darker = lower f)

BPM with ℓ_1 Balls

$$x_{k+1} \in \operatorname{Argmin} \{ f(y) \ : \ \|y - x_k\|_1 \le t_k \}$$

Example: $f(x) = \frac{1}{2}\left(|\langle a, x\rangle| - 2\right)_+^2 + 8\langle b, x\rangle^2, \qquad a = (\cos 20^\circ, \sin 20^\circ), \ \ b = (-\sin 20^\circ, \cos 20^\circ)$

BPM run

$d_k = \operatorname{dist}_1(x_k, \operatorname{Argmin} f)$


$f_k = f(x_k) - f_\star$

$g_k = \|\nabla f(x_k)\|_\infty$ (dual norm)

	$d_0 = 5.08$	$f_0 = 35.78$	$g_0 = 24.69$
$t_0 = 1.0$	$d_1 = 6.04$ ↑	$f_1 = 18.22$	$g_1 = 10.44$
$t_1 = 1.3$	$d_2 = 4.74$	$f_2 = 8.56$	$g_2 = 5.64$
$t_2 = 1.6$	$d_3 = 3.14$	$f_3 = 2.96$	$g_3 = 1.89$
$t_3 = 1.8$	$d_4 = 1.34$	$f_4 = 0.54$	$g_4 = 0.81$

$\operatorname{Argmin} f = [-2a, \ 2a]$

$f_\star := \min f = 0$



level sets of f
through the iterates
(darker = lower f)

BPM with ℓ_1 Balls

$$x_{k+1} \in \operatorname{Argmin}\{f(y) \;:\; \|y - x_k\|_1 \le t_k\}$$

Example: $f(x) = \frac{1}{2}\left(|\langle a, x\rangle| - 2\right)_+^2 + 8\langle b, x\rangle^2, \qquad a = (\cos 20^\circ, \sin 20^\circ),\ \ b = (-\sin 20^\circ, \cos 20^\circ)$

BPM run

$d_k = \operatorname{dist}_1(x_k, \operatorname{Argmin} f)$


$f_k = f(x_k) - f_\star$

$g_k = \|\nabla f(x_k)\|_\infty$ (dual norm)

	$d_0 = 5.08$	$f_0 = 35.78$	$g_0 = 24.69$
$t_0 = 1.0$	$d_1 = 6.04 \uparrow$	$f_1 = 18.22$	$g_1 = 10.44$
$t_1 = 1.3$	$d_2 = 4.74$	$f_2 = 8.56$	$g_2 = 5.64$
$t_2 = 1.6$	$d_3 = 3.14$	$f_3 = 2.96$	$g_3 = 1.89$
$t_3 = 1.8$	$d_4 = 1.34$	$f_4 = 0.54$	$g_4 = 0.81$

$\operatorname{Argmin} f = [-2a,\ 2a]$

$f_\star := \min f = 0$



level sets of f through the iterates (darker = lower f)

BPM with ℓ_1 Balls

$$x_{k+1} \in \operatorname{Argmin}\{f(y) \;:\; \|y - x_k\|_1 \le t_k\}$$

Example: $f(x) = \frac{1}{2}\left(|\langle a, x\rangle| - 2\right)_+^2 + 8\langle b, x\rangle^2, \qquad a = (\cos 20^\circ, \sin 20^\circ),\ \ b = (-\sin 20^\circ, \cos 20^\circ)$

BPM run

$d_k = \operatorname{dist}_1(x_k, \operatorname{Argmin} f)$


$f_k = f(x_k) - f_\star$

$g_k = \|\nabla f(x_k)\|_\infty$ (dual norm)

	$d_0 = 5.08$	$f_0 = 35.78$	$g_0 = 24.69$
$t_0 = 1.0$	$d_1 = 6.04$ ↑	$f_1 = 18.22$	$g_1 = 10.44$
$t_1 = 1.3$	$d_2 = 4.74$	$f_2 = 8.56$	$g_2 = 5.64$
$t_2 = 1.6$	$d_3 = 3.14$	$f_3 = 2.96$	$g_3 = 1.89$
$t_3 = 1.8$	$d_4 = 1.34$	$f_4 = 0.54$	$g_4 = 0.81$
$t_4 = 2.0$	$d_5 = 0.00$	$f_5 = 0.00$	$g_5 = 0.00$

$\operatorname{Argmin} f = [-2a,\ 2a]$

$f_\star := \min f = 0$



level sets of f through the iterates (darker = lower f)

BPM with ℓ_1 Balls

$$x_{k+1} \in \operatorname{Argmin} \{ f(y) \ : \ \|y - x_k\|_1 \le t_k \}$$

Example: $f(x) = \frac{1}{2}\left(|\langle a, x\rangle| - 2\right)_+^2 + 8\langle b, x\rangle^2, \qquad a = (\cos 20^\circ, \sin 20^\circ), \ \ b = (-\sin 20^\circ, \cos 20^\circ)$

BPM run

$d_k = \operatorname{dist}_1(x_k, \operatorname{Argmin} f)$

$f_k = f(x_k) - f_\star$


$g_k = \|\nabla f(x_k)\|_\infty$ (dual norm)

	$d_0 = 5.08$	$f_0 = 35.78$	$g_0 = 24.69$
$t_0 = 1.0$	$d_1 = 6.04 \uparrow$	$f_1 = 18.22$	$g_1 = 10.44$
$t_1 = 1.3$	$d_2 = 4.74$	$f_2 = 8.56$	$g_2 = 5.64$
$t_2 = 1.6$	$d_3 = 3.14$	$f_3 = 2.96$	$g_3 = 1.89$
$t_3 = 1.8$	$d_4 = 1.34$	$f_4 = 0.54$	$g_4 = 0.81$
$t_4 = 2.0$	$d_5 = 0.00$	$f_5 = 0.00$	$g_5 = 0.00$

f_k and g_k decrease at every step, but $d_1 > d_0$!

$\operatorname{Argmin} f = [-2a,\ 2a]$

$f_\star := \min f = 0$



level sets of f through the iterates (darker = lower f)

BPM with ℓ_∞ Balls


$$x_{k+1} \in \operatorname{Argmin} \{ f(y) \ : \ \|y - x_k\|_\infty \leq t_k \}$$

Example: $f(x) = \frac{1}{2}\left(|\langle a, x\rangle| - 2\right)_+^2 + 8\langle b, x\rangle^2, \qquad a = (\cos 20^\circ, \sin 20^\circ), \ \ b = (-\sin 20^\circ, \cos 20^\circ)$

BPM with ℓ_∞ Balls

$$x_{k+1} \in \operatorname{Argmin}\{f(y) \ : \ \|y - x_k\|_\infty \le t_k\}$$

Example: $f(x) = \frac{1}{2}\left(|\langle a, x\rangle| - 2\right)_+^2 + 8\langle b, x\rangle^2, \qquad a = (\cos 20^\circ, \sin 20^\circ), \ \ b = (-\sin 20^\circ, \cos 20^\circ)$



level sets of f
through the iterates
(darker = lower f)


BPM with ℓ_∞ Balls

$$x_{k+1} \in \operatorname{Argmin}\{f(y) \ : \ \|y - x_k\|_\infty \le t_k\}$$

Example: $f(x) = \frac{1}{2}\left(|\langle a, x\rangle| - 2\right)_+^2 + 8\langle b, x\rangle^2, \qquad a = (\cos 20^\circ, \sin 20^\circ), \ \ b = (-\sin 20^\circ, \cos 20^\circ)$

$\operatorname{Argmin} f \ = \ [-2a, \ 2a]$

$f_\star := \min f \ = \ 0$



level sets of f
through the iterates
(darker = lower f)

BPM with ℓ_∞ Balls

$$x_{k+1} \in \operatorname{Argmin}\{f(y) \;:\; \|y - x_k\|_\infty \le t_k\}$$

Example: $f(x) = \frac{1}{2}\left(|\langle a, x\rangle| - 2\right)_+^2 + 8\langle b, x\rangle^2, \qquad a = (\cos 20^\circ, \sin 20^\circ), \;\; b = (-\sin 20^\circ, \cos 20^\circ)$

BPM run

$d_k = \operatorname{dist}_\infty(x_k, \operatorname{Argmin} f)$


$f_k = f(x_k) - f_\star$

$g_k = \|\nabla f(x_k)\|_1$ (dual norm)

$d_0 = 3.57 \qquad f_0 = 49.53 \quad g_0 = 43.60$

$\operatorname{Argmin} f = [-2a,\ 2a]$

$f_\star := \min f = 0$



BPM with ℓ_∞ Balls

$$x_{k+1} \in \operatorname{Argmin}\{f(y) \;:\; \|y - x_k\|_\infty \le t_k\}$$

Example: $f(x) = \frac{1}{2}\left(|\langle a, x\rangle| - 2\right)_+^2 + 8\langle b, x\rangle^2, \qquad a = (\cos 20^\circ, \sin 20^\circ), \;\; b = (-\sin 20^\circ, \cos 20^\circ)$

BPM run

$d_k = \operatorname{dist}_\infty(x_k, \operatorname{Argmin} f)$


$f_k = f(x_k) - f_\star$

$g_k = \|\nabla f(x_k)\|_1$ (dual norm)

$d_0 = 3.57 \qquad f_0 = 49.53 \quad g_0 = 43.60$

$\operatorname{Argmin} f = [-2a,\ 2a]$

$f_\star := \min f = 0$



BPM with ℓ_∞ Balls

$$x_{k+1} \in \operatorname{Argmin}\{f(y) \;:\; \|y - x_k\|_\infty \le t_k\}$$

Example: $f(x) = \frac{1}{2}\left(|\langle a, x\rangle| - 2\right)_+^2 + 8\langle b, x\rangle^2, \qquad a = (\cos 20^\circ, \sin 20^\circ), \;\; b = (-\sin 20^\circ, \cos 20^\circ)$

BPM run

$d_k = \operatorname{dist}_\infty(x_k, \operatorname{Argmin} f)$


$f_k = f(x_k) - f_\star$

$g_k = \|\nabla f(x_k)\|_1$ (dual norm)

	$d_0 = 3.57$	$f_0 = 49.53$	$g_0 = 43.60$
$t_0 = 1.0$	$d_1 = 4.22 \uparrow$	$f_1 = 19.25$	$g_1 = 16.96$

$\operatorname{Argmin} f = [-2a,\ 2a]$

$f_\star := \min f = 0$



BPM with ℓ_∞ Balls

$$x_{k+1} \in \operatorname{Argmin}\{f(y) \;:\; \|y - x_k\|_\infty \le t_k\}$$

Example: $f(x) = \frac{1}{2}\left(|\langle a, x\rangle| - 2\right)_+^2 + 8\langle b, x\rangle^2, \qquad a = (\cos 20^\circ, \sin 20^\circ), \;\; b = (-\sin 20^\circ, \cos 20^\circ)$

BPM run

$d_k = \operatorname{dist}_\infty(x_k, \operatorname{Argmin} f)$


$f_k = f(x_k) - f_\star$

$g_k = \|\nabla f(x_k)\|_1$ (dual norm)

	$d_0 = 3.57$	$f_0 = 49.53$	$g_0 = 43.60$
$t_0 = 1.0$	$d_1 = 4.22 \uparrow$	$f_1 = 19.25$	$g_1 = 16.96$

$\operatorname{Argmin} f = [-2a,\ 2a]$

$f_\star := \min f = 0$



BPM with ℓ_∞ Balls

$$x_{k+1} \in \operatorname{Argmin}\{f(y) \;:\; \|y - x_k\|_\infty \le t_k\}$$

Example: $f(x) = \frac{1}{2}\left(|\langle a, x\rangle| - 2\right)_+^2 + 8\langle b, x\rangle^2, \qquad a = (\cos 20^\circ, \sin 20^\circ), \;\; b = (-\sin 20^\circ, \cos 20^\circ)$

BPM run

$d_k = \operatorname{dist}_\infty(x_k, \operatorname{Argmin} f)$


$f_k = f(x_k) - f_\star$

$g_k = \|\nabla f(x_k)\|_1$ (dual norm)

	$d_0 = 3.57$	$f_0 = 49.53$	$g_0 = 43.60$
$t_0 = 1.0$	$d_1 = 4.22 \uparrow$	$f_1 = 19.25$	$g_1 = 16.96$
$t_1 = 1.3$	$d_2 = 2.92$	$f_2 = 5.35$	$g_2 = 5.91$

$\operatorname{Argmin} f = [-2a,\ 2a]$

$f_\star := \min f = 0$



BPM with ℓ_∞ Balls

$$x_{k+1} \in \operatorname{Argmin}\{f(y) \;:\; \|y - x_k\|_\infty \le t_k\}$$

Example: $f(x) = \frac{1}{2}\left(|\langle a, x\rangle| - 2\right)_+^2 + 8\langle b, x\rangle^2, \qquad a = (\cos 20^\circ, \sin 20^\circ),\ \ b = (-\sin 20^\circ, \cos 20^\circ)$

BPM run

$d_k = \operatorname{dist}_\infty(x_k, \operatorname{Argmin} f)$


$f_k = f(x_k) - f_\star$

$g_k = \|\nabla f(x_k)\|_1$ (dual norm)

	$d_0 = 3.57$	$f_0 = 49.53$	$g_0 = 43.60$
$t_0 = 1.0$	$d_1 = 4.22$ ↑	$f_1 = 19.25$	$g_1 = 16.96$
$t_1 = 1.3$	$d_2 = 2.92$	$f_2 = 5.35$	$g_2 = 5.91$

$\operatorname{Argmin} f = [-2a,\ 2a]$

$f_\star := \min f = 0$



BPM with ℓ_∞ Balls

$$x_{k+1} \in \operatorname{Argmin}\{f(y) \;:\; \|y - x_k\|_\infty \le t_k\}$$

Example: $f(x) = \frac{1}{2}\left(|\langle a, x\rangle| - 2\right)_+^2 + 8\langle b, x\rangle^2, \qquad a = (\cos 20^\circ, \sin 20^\circ),\ \ b = (-\sin 20^\circ, \cos 20^\circ)$

BPM run

$d_k = \operatorname{dist}_\infty(x_k, \operatorname{Argmin} f)$


$f_k = f(x_k) - f_\star$

$g_k = \|\nabla f(x_k)\|_1$ (dual norm)

	$d_0 = 3.57$	$f_0 = 49.53$	$g_0 = 43.60$
$t_0 = 1.0$	$d_1 = 4.22$ ↑	$f_1 = 19.25$	$g_1 = 16.96$
$t_1 = 1.3$	$d_2 = 2.92$	$f_2 = 5.35$	$g_2 = 5.91$
$t_2 = 1.6$	$d_3 = 1.32$	$f_3 = 0.98$	$g_3 = 1.49$

$\operatorname{Argmin} f = [-2a,\ 2a]$

$f_\star := \min f = 0$



BPM with ℓ_∞ Balls

$$x_{k+1} \in \operatorname{Argmin}\{f(y) \;:\; \|y - x_k\|_\infty \le t_k\}$$

Example: $f(x) = \frac{1}{2}\left(|\langle a, x\rangle| - 2\right)_+^2 + 8\langle b, x\rangle^2, \qquad a = (\cos 20^\circ, \sin 20^\circ),\ \ b = (-\sin 20^\circ, \cos 20^\circ)$

BPM run

$d_k = \operatorname{dist}_\infty(x_k, \operatorname{Argmin} f)$


$f_k = f(x_k) - f_\star$

$g_k = \|\nabla f(x_k)\|_1$ (dual norm)

	$d_0 = 3.57$	$f_0 = 49.53$	$g_0 = 43.60$
$t_0 = 1.0$	$d_1 = 4.22$ ↑	$f_1 = 19.25$	$g_1 = 16.96$
$t_1 = 1.3$	$d_2 = 2.92$	$f_2 = 5.35$	$g_2 = 5.91$
$t_2 = 1.6$	$d_3 = 1.32$	$f_3 = 0.98$	$g_3 = 1.49$

$\operatorname{Argmin} f = [-2a,\ 2a]$

$f_\star := \min f = 0$



BPM with ℓ_∞ Balls

$$x_{k+1} \in \operatorname{Argmin}\{f(y) \;:\; \|y - x_k\|_\infty \le t_k\}$$

Example: $f(x) = \frac{1}{2}\left(|\langle a, x\rangle| - 2\right)_+^2 + 8\langle b, x\rangle^2, \qquad a = (\cos 20^\circ, \sin 20^\circ),\ b = (-\sin 20^\circ, \cos 20^\circ)$

BPM run

$d_k = \operatorname{dist}_\infty(x_k, \operatorname{Argmin} f)$


$f_k = f(x_k) - f_\star$

$g_k = \|\nabla f(x_k)\|_1$ (dual norm)

	$d_0 = 3.57$	$f_0 = 49.53$	$g_0 = 43.60$
$t_0 = 1.0$	$d_1 = 4.22$ ↑	$f_1 = 19.25$	$g_1 = 16.96$
$t_1 = 1.3$	$d_2 = 2.92$	$f_2 = 5.35$	$g_2 = 5.91$
$t_2 = 1.6$	$d_3 = 1.32$	$f_3 = 0.98$	$g_3 = 1.49$
$t_3 = 1.8$	$d_4 = 0.00$	$f_4 = 0.00$	$g_4 = 0.00$

$\operatorname{Argmin} f = [-2a,\ 2a]$

$f_\star := \min f = 0$



BPM with ℓ_∞ Balls

$$x_{k+1} \in \operatorname{Argmin}\{f(y) \;:\; \|y - x_k\|_\infty \le t_k\}$$

Example: $f(x) = \frac{1}{2}\big(|\langle a, x\rangle| - 2\big)_+^2 + 8\langle b, x\rangle^2, \qquad a = (\cos 20^\circ, \sin 20^\circ),\ \ b = (-\sin 20^\circ, \cos 20^\circ)$

BPM run

$d_k = \operatorname{dist}_\infty(x_k, \operatorname{Argmin} f)$

$f_k = f(x_k) - f_\star$


$g_k = \|\nabla f(x_k)\|_1$ (dual norm)

	$d_0 = 3.57$	$f_0 = 49.53$	$g_0 = 43.60$
$t_0 = 1.0$	$d_1 = 4.22$ ↑	$f_1 = 19.25$	$g_1 = 16.96$
$t_1 = 1.3$	$d_2 = 2.92$	$f_2 = 5.35$	$g_2 = 5.91$
$t_2 = 1.6$	$d_3 = 1.32$	$f_3 = 0.98$	$g_3 = 1.49$
$t_3 = 1.8$	$d_4 = 0.00$	$f_4 = 0.00$	$g_4 = 0.00$

f_k and g_k decrease at every step, but $d_1 > d_0$!

$\operatorname{Argmin} f = [-2a,\ 2a]$

$f_\star := \min f = 0$



Linear Programming via BPM with ℓ_∞ Balls

$$x_{k+1} \in \operatorname{Argmin} \{ \langle c, y \rangle \ : \ y \in P, \ \ \|y - x_k\|_\infty \leq t \}$$

Example: $f(x) = \langle c, x \rangle + \delta_P(x), \quad P = \{x \in \mathbb{R}^2 : \ \langle a_i, x \rangle \leq b_i, \ i = 1, \ldots, m\}, \quad m = 12, \ c = (1,2), \ t = 1.4$




P.R.
Rethinking Gradient Descent for Linear Programming: A Screened Broximal Approach
Manuscript, 2026

Linear Programming via BPM with ℓ_∞ Balls

$$x_{k+1} \in \operatorname{Argmin} \{ \langle c, y \rangle \ : \ y \in P, \ \ \|y - x_k\|_\infty \leq t \}$$

Example: $f(x) = \langle c, x \rangle + \delta_P(x), \quad P = \{x \in \mathbb{R}^2 : \ \langle a_i, x \rangle \leq b_i, \ i = 1, \ldots, m\}, \quad m = 12, \ c = (1,2), \ t = 1.4$




level sets of f
(darker = lower f)

P.R.
Rethinking Gradient Descent for Linear Programming: A Screened Broximal Approach
Manuscript, 2026

Linear Programming via BPM with ℓ_∞ Balls

$$x_{k+1} \in \operatorname{Argmin} \{ \langle c, y \rangle \ : \ y \in P, \ \ \|y - x_k\|_\infty \leq t \}$$

Example: $f(x) = \langle c, x \rangle + \delta_P(x), \quad P = \{x \in \mathbb{R}^2 : \ \langle a_i, x \rangle \leq b_i, \ i = 1, \ldots, m\}, \quad m = 12, \ c = (1,2), \ t = 1.4$



level sets of f
(darker = lower f)

P.R.
Rethinking Gradient Descent for Linear Programming: A Screened Broximal Approach
Manuscript, 2026

Linear Programming via BPM with ℓ_∞ Balls

$$x_{k+1} \in \operatorname{Argmin}\{\langle c, y\rangle \;:\; y \in P,\ \|y - x_k\|_\infty \le t\}$$

Example: $f(x) = \langle c, x\rangle + \delta_P(x)$, $P = \{x \in \mathbb{R}^2 :\ \langle a_i, x\rangle \le b_i,\ i = 1, \dots, m\}$, $m = 12$, $c = (1,2)$, $t = 1.4$


BPM run

$d_k = \operatorname{dist}_\infty(x_k, \operatorname{Argmin} f)$

$f_k = \langle c, x_k\rangle - f_\star$

$\mathcal{I}_k = \{i :\ b_i - \langle a_i, x_k\rangle < t\,\|a_i\|_1\}$

$d_0 = 5.05$ $f_0 = 11.45$



level sets of f
(darker = lower f)

P.R.
Rethinking Gradient Descent for Linear Programming: A Screened Broximal Approach
Manuscript, 2026

Linear Programming via BPM with ℓ_∞ Balls

$$x_{k+1} \in \operatorname{Argmin} \{ \langle c, y \rangle \ : \ y \in P, \ \ \|y - x_k\|_\infty \le t \}$$

Example: $f(x) = \langle c, x \rangle + \delta_P(x), \quad P = \{x \in \mathbb{R}^2 : \ \langle a_i, x \rangle \le b_i, \ i = 1, \dots, m\}, \quad m = 12, \ c = (1,2), \ t = 1.4$


BPM run

$d_k = \operatorname{dist}_\infty(x_k, \operatorname{Argmin} f)$

$f_k = \langle c, x_k \rangle - f_\star$

$\mathcal{I}_k = \{i : \ b_i - \langle a_i, x_k \rangle < t \, \|a_i\|_1\}$

$d_0 = 5.05 \quad f_0 = 11.45 \quad \mathcal{I}_0 = \emptyset$



level sets of f
(darker = lower f)

P.R.
Rethinking Gradient Descent for Linear Programming: A Screened Broximal Approach
Manuscript, 2026

Linear Programming via BPM with ℓ_∞ Balls

$$x_{k+1} \in \operatorname{Argmin}\{\langle c, y\rangle \;:\; y \in P,\;\; \|y - x_k\|_\infty \le t\}$$

Example: $f(x) = \langle c, x\rangle + \delta_P(x), \quad P = \{x \in \mathbb{R}^2 :\; \langle a_i, x\rangle \le b_i,\; i = 1, \dots, m\}, \quad m = 12,\; c = (1,2),\; t = 1.4$

BPM run


$d_k = \operatorname{dist}_\infty(x_k, \operatorname{Argmin} f)$

$f_k = \langle c, x_k\rangle - f_\star$

$\mathcal{I}_k = \{i :\; b_i - \langle a_i, x_k\rangle < t\,\|a_i\|_1\}$

$d_0 = 5.05 \quad f_0 = 11.45 \quad \mathcal{I}_0 = \emptyset$

$d_1 = 3.65 \quad f_1 = \;\; 7.25$



level sets of f
(darker = lower f)

P.R.
Rethinking Gradient Descent for Linear Programming: A Screened Broximal Approach
Manuscript, 2026

Linear Programming via BPM with ℓ_∞ Balls

$$x_{k+1} \in \operatorname{Argmin}\{\langle c, y\rangle \;:\; y \in P,\;\; \|y - x_k\|_\infty \le t\}$$

Example: $f(x) = \langle c, x\rangle + \delta_P(x), \quad P = \{x \in \mathbb{R}^2 :\; \langle a_i, x\rangle \le b_i,\; i = 1, \dots, m\}, \quad m = 12,\; c = (1,2),\; t = 1.4$

BPM run


$d_k = \operatorname{dist}_\infty(x_k, \operatorname{Argmin} f)$

$f_k = \langle c, x_k\rangle - f_\star$

$\mathcal{I}_k = \{i :\; b_i - \langle a_i, x_k\rangle < t\,\|a_i\|_1\}$

$d_0 = 5.05 \quad f_0 = 11.45 \quad \mathcal{I}_0 = \emptyset$

$d_1 = 3.65 \quad f_1 = \;\; 7.25 \quad \mathcal{I}_1 = \emptyset$



level sets of f
(darker = lower f)

P.R.
Rethinking Gradient Descent for Linear Programming: A Screened Broximal Approach
Manuscript, 2026

Linear Programming via BPM with ℓ_∞ Balls

$$x_{k+1} \in \operatorname{Argmin}\{\langle c, y\rangle \ : \ y \in P, \ \ \|y - x_k\|_\infty \le t\}$$

Example: $f(x) = \langle c, x\rangle + \delta_P(x), \quad P = \{x \in \mathbb{R}^2 : \ \langle a_i, x\rangle \le b_i, \ i = 1, \dots, m\}, \quad m = 12, \ c = (1,2), \ t = 1.4$


BPM run

$d_k = \operatorname{dist}_\infty(x_k, \operatorname{Argmin} f)$

$f_k = \langle c, x_k\rangle - f_\star$

$\mathcal{I}_k = \{i : \ b_i - \langle a_i, x_k\rangle < t\,\|a_i\|_1\}$

$d_0 = 5.05 \quad f_0 = 11.45 \quad \mathcal{I}_0 = \emptyset$
$d_1 = 3.65 \quad f_1 = \ 7.25 \quad \mathcal{I}_1 = \emptyset$
$d_2 = 2.25 \quad f_2 = \ 3.05$



level sets of f
(darker = lower f)

P.R.
Rethinking Gradient Descent for Linear Programming: A Screened Broximal Approach
Manuscript, 2026

Linear Programming via BPM with ℓ_∞ Balls

$$x_{k+1} \in \operatorname{Argmin}\{\langle c, y\rangle \ : \ y \in P, \ \ \|y - x_k\|_\infty \le t\}$$
$$= \operatorname{Argmin}\{\langle c, y\rangle \ : \ \langle a_i, y\rangle \le b_i \ \ \forall i \in \mathcal{I}_k, \ \ \|y - x_k\|_\infty \le t\}$$

Example: $f(x) = \langle c, x\rangle + \delta_P(x), \quad P = \{x \in \mathbb{R}^2 : \ \langle a_i, x\rangle \le b_i, \ i = 1, \dots, m\}, \quad m = 12, \ c = (1,2), \ t = 1.4$


BPM run

$d_k = \operatorname{dist}_\infty(x_k, \operatorname{Argmin} f)$

$f_k = \langle c, x_k\rangle - f_\star$

$\mathcal{I}_k = \{i : \ b_i - \langle a_i, x_k\rangle < t\,\|a_i\|_1\}$

$d_0 = 5.05 \quad f_0 = 11.45 \quad \mathcal{I}_0 = \emptyset$
$d_1 = 3.65 \quad f_1 = \ \ 7.25 \quad \mathcal{I}_1 = \emptyset$
$d_2 = 2.25 \quad f_2 = \ \ 3.05 \quad \mathcal{I}_2 = \{1\}$



level sets of f
(darker = lower f)

P.R.
Rethinking Gradient Descent for Linear Programming: A Screened Broximal Approach
Manuscript, 2026

Linear Programming via BPM with ℓ_∞ Balls

$$x_{k+1} \in \operatorname{Argmin}\{\langle c, y\rangle \;:\; y \in P,\ \|y - x_k\|_\infty \le t\}$$
$$= \operatorname{Argmin}\{\langle c, y\rangle \;:\; \langle a_i, y\rangle \le b_i\ \ \forall i \in \mathcal{I}_k,\ \ \|y - x_k\|_\infty \le t\}$$

Example: $f(x) = \langle c, x\rangle + \delta_P(x), \quad P = \{x \in \mathbb{R}^2 : \langle a_i, x\rangle \le b_i,\ i = 1, \dots, m\}, \quad m = 12,\ c = (1,2),\ t = 1.4$


BPM run

$d_k = \operatorname{dist}_\infty(x_k, \operatorname{Argmin} f)$

$f_k = \langle c, x_k\rangle - f_\star$

$\mathcal{I}_k = \{i :\ b_i - \langle a_i, x_k\rangle < t\,\|a_i\|_1\}$

$d_0 = 5.05 \quad f_0 = 11.45 \quad \mathcal{I}_0 = \emptyset$
$d_1 = 3.65 \quad f_1 = 7.25 \quad \mathcal{I}_1 = \emptyset$
$d_2 = 2.25 \quad f_2 = 3.05 \quad \mathcal{I}_2 = \{1\}$
$d_3 = 0.85 \quad f_3 = 0.69$



level sets of f
(darker = lower f)

P.R.
Rethinking Gradient Descent for Linear Programming: A Screened Broximal Approach
Manuscript, 2026

Linear Programming via BPM with ℓ_∞ Balls

$$x_{k+1} \in \operatorname{Argmin}\{\langle c, y\rangle \;:\; y \in P,\ \|y - x_k\|_\infty \le t\}$$
$$= \operatorname{Argmin}\{\langle c, y\rangle \;:\; \langle a_i, y\rangle \le b_i\ \forall i \in \mathcal{I}_k,\ \|y - x_k\|_\infty \le t\}$$

Example: $f(x) = \langle c, x\rangle + \delta_P(x)$, $P = \{x \in \mathbb{R}^2 : \langle a_i, x\rangle \le b_i,\ i = 1, \dots, m\}$, $m = 12$, $c = (1, 2)$, $t = 1.4$


BPM run

$d_k = \operatorname{dist}_\infty(x_k, \operatorname{Argmin} f)$

$f_k = \langle c, x_k\rangle - f_\star$

$\mathcal{I}_k = \{i :\ b_i - \langle a_i, x_k\rangle < t\,\|a_i\|_1\}$

$d_0 = 5.05$ $f_0 = 11.45$ $\mathcal{I}_0 = \emptyset$
$d_1 = 3.65$ $f_1 = 7.25$ $\mathcal{I}_1 = \emptyset$
$d_2 = 2.25$ $f_2 = 3.05$ $\mathcal{I}_2 = \{1\}$
$d_3 = 0.85$ $f_3 = 0.69$ $\mathcal{I}_3 = \{1, 12\}$



level sets of f
(darker = lower f)

P.R.
Rethinking Gradient Descent for Linear Programming: A Screened Broximal Approach
Manuscript, 2026

Linear Programming via BPM with ℓ_∞ Balls

$$x_{k+1} \in \operatorname{Argmin}\{\langle c, y\rangle \;:\; y \in P,\ \|y - x_k\|_\infty \le t\}$$
$$= \operatorname{Argmin}\{\langle c, y\rangle \;:\; \langle a_i, y\rangle \le b_i\ \forall i \in \mathcal{I}_k,\ \|y - x_k\|_\infty \le t\}$$

Example: $f(x) = \langle c, x\rangle + \delta_P(x)$, $P = \{x \in \mathbb{R}^2 : \langle a_i, x\rangle \le b_i,\ i = 1, \dots, m\}$, $m = 12$, $c = (1, 2)$, $t = 1.4$


BPM run

$d_k = \operatorname{dist}_\infty(x_k, \operatorname{Argmin} f)$

$f_k = \langle c, x_k\rangle - f_\star$

$\mathcal{I}_k = \{i :\ b_i - \langle a_i, x_k\rangle < t\,\|a_i\|_1\}$

$d_0 = 5.05$	$f_0 = 11.45$	$\mathcal{I}_0 = \emptyset$
$d_1 = 3.65$	$f_1 = 7.25$	$\mathcal{I}_1 = \emptyset$
$d_2 = 2.25$	$f_2 = 3.05$	$\mathcal{I}_2 = \{1\}$
$d_3 = 0.85$	$f_3 = 0.69$	$\mathcal{I}_3 = \{1, 12\}$
$d_4 = 0.00$	$f_4 = 0.00$	$x_4 = x_\star$



level sets of f
(darker = lower f)

P.R.
Rethinking Gradient Descent for Linear Programming: A Screened Broximal Approach
Manuscript, 2026

Linear Programming via BPM with ℓ_∞ Balls

$$x_{k+1} \in \operatorname{Argmin}\{\langle c, y\rangle \;:\; y \in P,\ \|y - x_k\|_\infty \le t\}$$
$$= \operatorname{Argmin}\{\langle c, y\rangle \;:\; \langle a_i, y\rangle \le b_i\ \ \forall i \in \mathcal{I}_k,\ \ \|y - x_k\|_\infty \le t\}$$

Example: $f(x) = \langle c, x\rangle + \delta_P(x)$, $P = \{x \in \mathbb{R}^2 : \langle a_i, x\rangle \le b_i,\ i = 1, \dots, m\}$, $m = 12$, $c = (1, 2)$, $t = 1.4$


BPM run

$d_k = \operatorname{dist}_\infty(x_k, \operatorname{Argmin} f)$

$f_k = \langle c, x_k\rangle - f_\star$

$\mathcal{I}_k = \{i :\ b_i - \langle a_i, x_k\rangle < t\,\|a_i\|_1\}$

$d_0 = 5.05$	$f_0 = 11.45$	$\mathcal{I}_0 = \emptyset$
$d_1 = 3.65$	$f_1 = 7.25$	$\mathcal{I}_1 = \emptyset$
$d_2 = 2.25$	$f_2 = 3.05$	$\mathcal{I}_2 = \{1\}$
$d_3 = 0.85$	$f_3 = 0.69$	$\mathcal{I}_3 = \{1, 12\}$
$d_4 = 0.00$	$f_4 = 0.00$	$x_4 = x_\star$



level sets of f
(darker = lower f)

P.R.
Rethinking Gradient Descent for Linear Programming: A Screened Broximal Approach
Manuscript, 2026

LP $\longrightarrow$ 4 small LPs, one per BPM step;
each uses only $|\mathcal{I}_k| \le 2$ of the $m = 12$ constraints

Part 2
Broximal Point Method: Convergence Theory



Kaja Gruntkowska, Hanmin Li, Aadi Rane, P.R.
The ball-proximal (="broximal") point method: a new algorithm, convergence theory, and applications
arXiv:2502.02002, 2025



P.R., Hanmin Li
A sharper theory of ball-proximal optimization: convergence and radius selection
arXiv:2609.35147, 2026

Problem Formulation & Assumptions

$$\min_{x \in \mathbb{R}^d} f(x)$$



The Ball-Proximal (="Broximal") Point Method:
a New Algorithm, Convergence Theory, and Applications

Kaja Gruntkowska [1] Hanmin Li [1] Aadi Rane [1,2] Peter Richtárik [1]

Abstract

Non-smooth and non-convex global optimization poses significant challenges across various applications, where standard gradient-based methods often struggle. We propose the *Ball-Proximal Point Method*, *Broximal Point Method*, or *Ball Point Method* (BPM) for short – a novel algorithmic framework inspired by the classical Proximal Point Method (PPM) (Rockafellar, 1976), which, as we show, sheds new light on several foundational optimization paradigms and phenomena, including non-convex and non-smooth optimization, acceleration, smoothing, adaptive stepsize selection, and trust-region methods. At the core of BPM lies the *ball-proximal ("broximal") operator*, which arises from the classical proximal operator by replacing the quadratic distance penalty with a hard constraint. While BPM can be seen contrast with the sublinear rate of PPM in the nonsmooth convex regime, we prove that BPM converges linearly and in a finite number of steps in the same regime. Furthermore, by introducing the concept of ball-convexity, we prove that BPM retains the same global convergence guarantees under weaker assumptions, making it a powerful tool for a broader class of potentially non-convex optimization problems. Just like PPM plays the role of a conceptual method inspiring the development of practically efficient algorithms and algorithmic elements, e.g., gradient descent, adaptive step sizes, acceleration (Alv & Sra, 2020), and "W" in AdamW (Zhuang et al., 2022), we believe that BPM should be understood in the same manner: as a blueprint and inspiration for further development.

[1]King Abdullah University of Science and Technology, Thuwal, Saudi Arabia [2]University of California, Berkeley, USA.

[*]The work of Aadi Rane was conducted during a VSRP internship at KAUST.

1. Introduction

The minimization of non-convex functions is a fundamental challenge across many fields, including machine learning, optimization, applied mathematics, signal processing and operations research. Solving such problems is integral to most machine learning algorithms arising in both training and inference, where non-convex objectives or constraints are often necessary to capture complex prediction tasks.

1.1. Global nonconvex optimization

In this paper, we propose a new meta-algorithm (see Section 1.3) capable of finding *global* minimizers for a specific (new) class of non-convex functions. In particular, we introduce an algorithmic framework designed to solve the (potentially non-convex) optimization problem

$$\min_{x \in \mathbb{R}^d} f(x), \tag{1}$$

where $f : \mathbb{R}^d \mapsto \mathbb{R} \cup \{+\infty\}$ is assumed to be proper (which means that the set $\text{dom} f := \{x \in \mathbb{R}^d : f(x) < +\infty\}$ is nonempty), closed, and have at least one minimizer. We let X_f be the set of all minimizers of f, and $f_* := \min_x f(x)$.

1.2. The ball-proximal operator

A key inspiration for our method stems from the well-known Proximal Point Method (PPM) (Rockafellar, 1976), which iteratively adds a quadratic penalty term to the objective (see Section 7 for more details) and solves a modified subproblem at each step. Building on this idea, we introduce the *ball-proximal ("broximal") operator*:

Definition 1.1 (Ball-Proximal Operator). *The ball-proximal ("broximal") operator* with radius $t > 0$ associated with a function $f : \mathbb{R}^d \mapsto \mathbb{R} \cup \{+\infty\}$ is given by

$$\text{brox}_t^f(x) := \operatorname*{argmin}_{y \in B_t(x)} f(y), \tag{2}$$

where $B_t(x) := \{x \in \mathbb{R}^d : \|x - x\| \leq t\}$ and $\|\cdot\|$ is the standard Euclidean norm.

According to this definition, for a given input point $x \in \mathbb{R}^d$, $\text{brox}_t^f(x)$ returns the minimizer(s) of f within the ball of radius t centered at x.

1

arXiv:2502.02002v1 [math.OC] 4 Feb 2025

Broximal Point Method (BPM) with ℓ_2 Balls

$$x_{k+1} \in \text{Argmin}\{f(y) \ : \ \|y - x_k\|_2 \leq t_k\}$$



A Sharper Theory of Ball-Proximal Optimization:
Convergence and Radius Selection

Peter Richtárik Hanmin Li

King Abdullah University of Science and Technology
Thuwal 23955-6900, Saudi Arabia

28 September 2026

Abstract

We study the exact Euclidean ball-proximal point method for proper, closed, convex functions, where each iteration minimizes the objective over a ball centered at the current point. Retaining the objective gap is the decrease of squared distance yields sharper bounds on objective values, stationarity, and the symmetric Bregman distance to a minimizer. For constant radius $t > 0$ and initial distance $R_0 > 0$ to the solution set, the objective gap after K iterations is at most its initial value multiplied by $\exp(-2R^2t^2/D_0^4)$.

We characterize convergence for arbitrary positive radius sequences. If their sum diverges, the method reaches a minimizer is finitely many iterations whenever one exists; otherwise, the objective values converge to the infimum and the iterates escape every bounded set. If the radii are summable, the iterates converge to a possibly nonoptimal point. Self-contraction gives finite trajectory length for bounded iterates and a constant-radius termination bound $O(d(1 + D_0/t))$, whose implicit constant depends only on the dimension. A polyhedral family with $\Theta(d)$ facets shows that this dependence on d cannot be improved in general.

We also identify a minimax successful geometric decay factor and analyze adaptive radius rules based on subgradients or objective gaps, epigraph reformulation, and relaxed updates. Together, these results strengthen the foundations and convergence guarantees of the method without assuming smoothness or strong convexity.

Keywords: ball-proximal point method; convex optimization; finite termination; self-contracted sequences; radius selection; oracle complexity.
Mathematics Subject Classification (2020): Primary 90C25; Secondary 90C60, 65K05.

1 Introduction

We study the convex optimization problem

$$\min_{x \in \mathbb{R}^d} f(x), \tag{1}$$

where $f : \mathbb{R}^d \to \mathbb{R} \cup \{+\infty\}$ is proper, closed, and convex. Unless stated otherwise, we assume neither that the infimum is attained nor that f is smooth. We focus on the exact ball-oracle iteration considered by Carmon et al. (2020, Appendix A) and subsequently developed as the ball-proximal (broximal) point framework by Gruntkowska et al. (2025). Starting from $x^0 \in \text{dom } f$, the ball-proximal point method (BPM) minimizes f over a Euclidean ball centered at the current point:

$$x^{k+1} \in \text{brox}_t^f(x^k) := \operatorname*{argmin}_{u \in \mathbb{R}^d : u \in \mathbb{R}^d} f(u), \qquad t_k > 0, \tag{BPM}$$

1

arXiv:2609.35147v1 [math.OC] 28 Sep 2026

Problem Formulation & Assumptions



The Ball-Proximal (="Broximal") Point Method:
a New Algorithm, Convergence Theory, and Applications

Kaja Gruntkowska [1] Hanmin Li [1] Aadi Rane [1,2] Peter Richtárik [1]

Abstract

Non-smooth and non-convex global optimization poses significant challenges across various applications, where standard gradient-based methods often struggle. We propose the *Ball-Proximal Point Method*, *Broximal Point Method*, or *Ball Point Method* (BPM) for short – a novel algorithmic framework inspired by the classical Proximal Point Method (PPM) (Rockafellar, 1976), which, as we show, sheds new light on several foundational optimization paradigms and phenomena, including non-convex and non-smooth optimization, acceleration, smoothing, adaptive stepsize selection, and trust-region methods. At the core of BPM lies the *ball-proximal ("broximal") operator*, which arises from the classical proximal operator by replacing the quadratic distance penalty with a hard constraint. While PPM can be seen to contract with the sublinear rate of $\mathcal{O}(1/k)$ in fully nonsmooth convex regime, we prove that BPM converges linearly and in a finite number of steps in the same regime. Furthermore, by introducing the concept of ball-convexity, we prove that BPM retains the same global convergence guarantees under weaker assumptions, making it a powerful tool for a broader class of potentially non-convex optimization problems. Just like PPM plays the role of a conceptual method inspiring the development of practically efficient algorithms and algorithmic elements, e.g., gradient descent, adaptive step sizes, acceleration (Alp & Sra, 2020), and "W" in AdamW (Zhuang et al., 2022), we believe that BPM should be understood in the same manner: as a blueprint and inspiration for further development.

[1]King Abdullah University of Science and Technology, Thuwal, Saudi Arabia [2]University of California, Berkeley, USA.

*The work of Aadi Rane was conducted during a VSRP internship at KAUST.

1. Introduction

The minimization of non-convex functions is a fundamental challenge across many fields, including machine learning, optimization, applied mathematics, signal processing and operations research. Solving such problems is integral to most machine learning algorithms arising in both training and inference, where non-convex objectives or constraints are often necessary to capture complex prediction tasks.

1.1. Global nonconvex optimization

In this paper, we propose a new meta-algorithm (see Section 1.3) capable of finding *global* minimizers for a specific (new) class of non-convex functions. In particular, we introduce an algorithmic framework designed to solve the (potentially non-convex) optimization problem

$$\min_{x\in\mathbb{R}^d} f(x), \tag{1}$$

where $f:\mathbb{R}^d\mapsto\mathbb{R}\cup\{+\infty\}$ is assumed to be proper (which means that the set $\operatorname{dom} f := \{x\in\mathbb{R}^d : f(x) < +\infty\}$ is nonempty), closed, and have at least one minimizer. We let X_f be the set of all minimizers of f, and $f_* := \min_x f(x)$.

1.2. The ball-proximal operator

A key inspiration for our method stems from the well-known Proximal Point Method (PPM) (Rockafellar, 1976), which iteratively adds a quadratic penalty term to the objective (see Section 7 for more details) and solves a modified subproblem at each step. Building on this idea, we introduce the *ball-proximal ("broximal") operator*:

Definition 1.1 (Ball-Proximal Operator). *The ball-proximal ("broximal") operator* with radius $t>0$ associated with a function $f:\mathbb{R}^d\mapsto\mathbb{R}\cup\{+\infty\}$ is given by

$$\operatorname{brox}_t^f(x) := \operatorname*{argmin}_{y\in B_t(x)} f(y), \tag{2}$$

where $B_t(x) := \{z\in\mathbb{R}^d : \|z-x\|\le t\}$ and $\|\cdot\|$ is the standard Euclidean norm.

According to this definition, for a given input point $x\in\mathbb{R}^d$, $\operatorname{brox}_t^f(x)$ returns the minimizer(s) of f within the ball of radius t centered at x.

1

arXiv:2502.02002v1 [math.OC] 4 Feb 2025

Broximal Point Method (BPM) with ℓ_2 Balls

$$x_{k+1} \in \operatorname{Argmin}\{f(y) \;:\; \|y - x_k\|_2 \le t_k\}$$



A Sharper Theory of Ball-Proximal Optimization:
Convergence and Radius Selection

Peter Richtárik Hanmin Li

King Abdullah University of Science and Technology
Thuwal 23955-6900, Saudi Arabia

28 September 2026

Abstract

We study the exact Euclidean ball-proximal point method for proper, closed, convex functions, where each iteration minimizes the objective over a ball centered at the current point. Retaining the objective gap is the decrease of squared distance yields sharper bounds on objective values, stationarity, and the symmetric Bregman distance to a minimizer. For constant radius $t>0$ and initial distance $R_0>0$ to the solution set, the objective gap after K iterations is at most its initial value multiplied by $\exp(-2R^2t^2/D_0^4)$.

We characterize convergence for arbitrary positive radius sequences. If their sum diverges, the method reaches a minimizer in finitely many iterations whenever one exists; otherwise, the objective values converge to the infimum and the iterates escape every bounded set. If the radii are summable, the iterates converge to a possibly nonoptimal point. Self-contraction gives finite trajectory length for bounded iterates and a constant-radius termination bound $O(t+D_0/t)$, whose implicit constant depends only on the dimension. A polyhedral family shows that this dimension dependence is unavoidable, with matching upper and lower bounds.

We also identify a minimum successful geometric decay factor and analyze adaptive radius rules based on subgradients or objective gaps, epigraph reformulation, and relaxed updates. Together, these results strengthen the foundations and convergence guarantees of the method without assuming smoothness or strong convexity.

Keywords: ball-proximal point method; convex optimization; finite termination; self-contracted sequences; radius selection; oracle complexity.
Mathematics Subject Classification (2020): Primary 90C25; Secondary 90C60, 65K05.

1 Introduction

We study the convex optimization problem

$$\min_{x\in\mathbb{R}^d} f(x), \tag{1}$$

where $f:\mathbb{R}^d\to\mathbb{R}\cup\{+\infty\}$ is proper, closed, and convex. Unless stated otherwise, we assume neither that the infimum is attained nor that f is smooth. We focus on the exact ball-oracle iteration considered by Carmon et al. (2020, Appendix A) and subsequently developed as the ball-proximal (broximal) point framework by Gruntkowska et al. (2025). Starting from $x^0\in\operatorname{dom} f$, the ball-proximal point method (BPM) minimizes f over a Euclidean ball centered at the current point:

$$x^{k+1}\in\operatorname{brox}_t^f(x^k) := \operatorname*{argmin}_{u\in\mathbb{R}^d,\, u\in B_t(x^k)} f(u), \qquad t_k>0, \tag{BPM}$$

1

arXiv:2609.35147v1 [math.OC] 28 Sep 2026

$$f:\mathbb{R}^d\to\mathbb{R}\cup\{+\infty\} \text{ is proper, closed \& convex}$$

Problem Formulation & Assumptions

$$\min_{x \in \mathbb{R}^d} f(x)$$



Broximal Point Method (BPM) with ℓ_2 Balls

$$x_{k+1} \in \operatorname{Argmin}\{f(y) \ : \ \|y - x_k\|_2 \leq t_k\}$$



$$f : \mathbb{R}^d \to \mathbb{R} \cup \{+\infty\} \text{ is proper, closed \& convex}$$

$$x_\star \in \operatorname{Argmin} f \neq \emptyset, \qquad f_\star := \min f$$

Problem Formulation & Assumptions

$$\min_{x \in \mathbb{R}^d} f(x)$$



The Ball-Proximal (="Broximal") Point Method: a New Algorithm, Convergence Theory, and Applications

Kaja Gruntkowska · Hanmin Li · Aadi Rane · Peter Richtárik

Abstract

Non-smooth and non-convex global optimization poses significant challenges across various applications, where standard gradient-based methods often struggle. We propose the *Ball-Proximal Point Method*, *Broximal Point Method*, or *Ball Point Method* (BPM) for short – a novel algorithmic framework inspired by the classical Proximal Point Method (PPM) (Rockafellar, 1976), which, as we show, sheds new light on several foundational optimization paradigms and phenomena, including non-convex and non-smooth optimization, acceleration, smoothing, adaptive stepsize selection, and trust-region methods. At the core of BPM lies the *ball-proximal* ("broximal") operator, which arises from the classical proximal operator by replacing the quadratic distance penalty with a hard constraint. While BPM can be seen contrast with the subtlesse rule of PPM in the nonsmooth convex regime, we prove that BPM converges linearly and in a finite number of steps in the same regime. Furthermore, by introducing the concept of ball-convexity, we prove that BPM retains the same global convergence guarantees under weaker assumptions, making it a powerful tool for a broader class of potentially non-convex optimization problems. Just like PPM plays the role of a conceptual method inspiring the development of practically efficient algorithms and algorithmic elements, e.g., gradient descent, adaptive step sizes, acceleration (Alv & Yu, 2020), and "R" in AdaGrad (Zhang et al., 2022), we believe that BPM should be understood in the same manner: as a blueprint and inspiration for further development.

1. Introduction

The minimization of non-convex functions is a fundamental challenge across many fields, including machine learning, optimization, applied mathematics, signal processing and operations research. Solving such problems is integral to most machine learning algorithms arising in both training and inference, where non-convex objectives or constraints are often necessary to capture complex prediction tasks.

1.1. Global nonconvex optimization

In this paper, we propose a new meta-algorithm (see Section 1.3) capable of finding *global* minimizers for a specific (new) class of non-convex functions. In particular, we introduce an algorithmic framework designed to solve the (potentially non-convex) optimization problem

$$\min_{x \in \mathbb{R}^d} f(x), \tag{1}$$

where $f : \mathbb{R}^d \to \mathbb{R} \cup \{+\infty\}$ is assumed to be proper (which means that the set $\mathrm{dom} f := \{x \in \mathbb{R}^d : f(x) < +\infty\}$ is nonempty), closed, and have at least one minimizer. We let $X_\star$ be the set of all minimizers of f, and $f_\star := \min_x f(x)$.

1.2. The ball-proximal operator

A key inspiration for our method stems from the well-known Proximal Point Method (PPM) (Rockafellar, 1976), which iteratively adds a quadratic penalty term to the objective (see Section 7 for more details) and solves a modified subproblem at each step. Building on this idea, we introduce the *ball-proximal* ("broximal") *operator*:

Definition 1.1 (Ball-Proximal Operator). *The ball-proximal ("broximal") operator* with radius $t > 0$ associated with a function $f : \mathbb{R}^d \to \mathbb{R} \cup \{+\infty\}$ is given by

$$\mathrm{brox}_t^f(x) := \operatorname*{argmin}_{z \in B_t(x)} f(z), \tag{2}$$

where $B_t(x) := \{z \in \mathbb{R}^d : \|z - x\| \le t\}$ and $\|\cdot\|$ is the standard Euclidean norm.

According to this definition, for a given input point $x \in \mathbb{R}^d$, $\mathrm{brox}_t^f(x)$ returns the minimizer(s) of f within the ball of radius t centered at x.

King Abdullah University of Science and Technology, Thuwal, Saudi Arabia. University of California, Berkeley, USA.

*The work of Aadi Rane was conducted during a VSRP internship at KAUST.

arXiv:2502.02002v1 [math.OC] 4 Feb 2025

Broximal Point Method (BPM) with ℓ_2 Balls

$$x_{k+1} \in \mathrm{Argmin}\{f(y) \ : \ \|y - x_k\|_2 \le t_k\}$$



A Sharper Theory of Ball-Proximal Optimization: Convergence and Radius Selection

Peter Richtárik · Hanmin Li

King Abdullah University of Science and Technology, Thuwal 23955-6900, Saudi Arabia

28 September 2026

Abstract

We study the exact Euclidean ball-proximal point method for proper, closed, convex functions, where each iteration minimizes the objective over a ball centered at the current point. Retaining the objective gap is the decrease of squared distance yields sharper bounds on objective values, stationarity, and the symmetric Bregman distance to a minimizer. For constant radius $t > 0$ and initial distance $R_0 > 0$ to the solution set, the objective gap after K iterations is at most its initial value multiplied by $\exp(-2K^2t^2/D_0^2)$.

We characterize convergence for arbitrary positive radius sequences. If their sum diverges, the method reaches a minimizer in finitely many iterations whenever one exists; otherwise, the objective values converge to the infimum and the iterates escape every bounded set. If the radii are summable, the iterates converge to a possibly nonoptimal point. Self-contraction gives finite trajectory length for bounded iterates and a constant-radius termination bound $O(\rho(1+D_0/t))$, whose implicit constant depends only on the dimension. A polyhedral family shows that this dimension dependence is unavoidable, with matching lower and upper bounds.

We also identify a minimax successful geometric decay factor and analyze adaptive radius rules based on subgradients or objective gaps, epigraph reformulation, and relaxed updates. Together, these results strengthen the foundations and convergence guarantees of the method without assuming smoothness or strong convexity.

Keywords: ball-proximal point method; convex optimization; finite termination; self-contracted sequences; radius selection; oracle complexity.
Mathematics Subject Classification (2020): Primary 90C25; Secondary 90C60, 65K05.

1 Introduction

We study the convex optimization problem

$$\min_{x \in \mathbb{R}^d} f(x), \tag{1}$$

where $f : \mathbb{R}^d \to \mathbb{R} \cup \{+\infty\}$ is proper, closed, and convex. Unless stated otherwise, we assume neither that the infimum is attained nor that f is smooth. We focus on the exact ball-oracle iteration considered by Carmon et al. (2020, Appendix A) and subsequently developed as the ball-proximal (broximal) point framework by Gruntkowska et al. (2025). Starting from $x^0 \in \mathrm{dom}\, f$, the ball-proximal point method (BPM) minimizes f over a Euclidean ball centered at the current point:

$$x^{k+1} \in \mathrm{brox}_t^f(x^k) := \operatorname*{argmin}_{u \in \mathbb{R}^d,\, x^k} f(u), \qquad t_k > 0, \tag{BPM}$$


arXiv:2609.35147v1 [math.OC] 28 Sep 2026

$$f : \mathbb{R}^d \to \mathbb{R} \cup \{+\infty\} \text{ is proper, closed \& convex}$$

$$x_\star \in \mathrm{Argmin}\, f \neq \emptyset, \qquad f_\star := \min f$$

$$f(x_0) \neq +\infty$$

The Condition Number κ



Fixed-Radius Theory: Distance



Kaja Gruntkowska, Hanmin Li, Aadi Rane, P.R.
The ball-proximal (="broximal") point method: a new algorithm, convergence theory, and applications
arXiv:2502.02002, 2025

Fixed-Radius Theory: Distance

$$d_k^2 \leq \max\left\{d_0^2 - kt^2, 0\right\}$$



Kaja Gruntkowska, Hanmin Li, Aadi Rane, P.R.
The ball-proximal (="broximal") point method: a new algorithm, convergence theory, and applications
arXiv:2502.02002, 2025

Fixed-Radius Theory: Distance

$$d_k^2 \leq \max\left\{d_0^2 - kt^2, 0\right\}$$

$$d_k = \mathrm{dist}_2(x_k, \mathrm{Argmin}\, f)$$
$$f_k = f(x_k) - f_\star$$
$$g_k = \mathrm{dist}_2(0, \partial f(x_k))$$



Kaja Gruntkowska, Hanmin Li, Aadi Rane, P.R.
The ball-proximal (="broximal") point method: a new algorithm, convergence theory, and applications
arXiv:2502.02002, 2025

Fixed-Radius Theory: Distance

$$d_k = \mathrm{dist}_2(x_k, \mathrm{Argmin}\, f)$$
$$f_k = f(x_k) - f_\star$$
$$g_k = \mathrm{dist}_2(0, \partial f(x_k))$$

$$d_k^2 \leq \max\left\{d_0^2 - kt^2, 0\right\}$$

$$k \geq \frac{d_0^2}{t^2} \quad \Longrightarrow \quad d_k = 0$$



Kaja Gruntkowska, Hanmin Li, Aadi Rane, P.R.
The ball-proximal (="broximal") point method: a new algorithm, convergence theory, and applications
arXiv:2502.02002, 2025

Fixed-Radius Theory: Distance

$$d_k = \mathrm{dist}_2(x_k, \mathrm{Argmin}\, f)$$
$$f_k = f(x_k) - f_\star$$
$$g_k = \mathrm{dist}_2(0, \partial f(x_k))$$

$$d_k^2 \leq \max\left\{d_0^2 - kt^2, 0\right\}$$

$$k \geq \frac{d_0^2}{t^2} \quad \Longrightarrow \quad d_k = 0$$

No dependence on a condition number!

Kaja Gruntkowska, Hanmin Li, Aadi Rane, P.R.
The ball-proximal (="broximal") point method: a new algorithm, convergence theory, and applications
arXiv:2502.02002, 2025

Fixed-Radius Theory: Function



P.R., Hanmin Li
A sharper theory of ball-proximal optimization: convergence and radius selection
arXiv:2609.35147, 2026

Fixed-Radius Theory: Function

$$f_k \leq f_0 \cdot e^{-2k^2 t^2 / d_0^2}$$



P.R., Hanmin Li
A sharper theory of ball-proximal optimization: convergence and radius selection
arXiv:2609.35147, 2026

Fixed-Radius Theory: Function

$$f_k \leq f_0 \cdot e^{-2k^2 t^2 / d_0^2}$$

$$d_k = \mathrm{dist}_2(x_k, \mathrm{Argmin}\, f)$$
$$f_k = f(x_k) - f_\star$$
$$g_k = \mathrm{dist}_2(0, \partial f(x_k))$$



P.R., Hanmin Li
A sharper theory of ball-proximal optimization: convergence and radius selection
arXiv:2609.35147, 2026

Fixed-Radius Theory: Function

$$d_k = \mathrm{dist}_2(x_k, \mathrm{Argmin}\, f)$$
$$f_k = f(x_k) - f_\star$$
$$g_k = \mathrm{dist}_2(0, \partial f(x_k))$$

$$f_k \le f_0 \cdot e^{-2k^2 t^2 / d_0^2}$$

$$k \ge \frac{d_0}{\sqrt{2}\, t} \sqrt{\ln \frac{f_0}{\varepsilon}} \quad \Longrightarrow \quad f_k \le \varepsilon$$



P.R., Hanmin Li
A sharper theory of ball-proximal optimization: convergence and radius selection
arXiv:2609.35147, 2026

Fixed-Radius Theory: Function

$$d_k = \mathrm{dist}_2(x_k, \mathrm{Argmin}\, f)$$
$$f_k = f(x_k) - f_\star$$
$$g_k = \mathrm{dist}_2(0, \partial f(x_k))$$

$$f_k \le f_0 \cdot e^{-2k^2 t^2 / d_0^2}$$

$$k \ge \frac{d_0}{\sqrt{2}\, t} \sqrt{\ln \frac{f_0}{\varepsilon}} \quad \Longrightarrow \quad f_k \le \varepsilon$$

No dependence on a condition number!



P.R., Hanmin Li
A sharper theory of ball-proximal optimization: convergence and radius selection
arXiv:2609.35147, 2026

Fixed-Radius Theory: Gradient



P.R., Hanmin Li
A sharper theory of ball-proximal optimization: convergence and radius selection
arXiv:2609.35147, 2026

Fixed-Radius Theory: Gradient

$$g_{k+1} \leq g_k$$



P.R., Hanmin Li
A sharper theory of ball-proximal optimization: convergence and radius selection
arXiv:2609.35147, 2026

Fixed-Radius Theory: Gradient

$$g_{k+1} \le g_k$$

$$g_k \le \frac{f_0}{t} \cdot e^{-2(k-1)^2 t^2 / d_0^2}$$

P.R., Hanmin Li
A sharper theory of ball-proximal optimization: convergence and radius selection
arXiv:2609.35147, 2026

Fixed-Radius Theory: Gradient

$$d_k = \mathrm{dist}_2(x_k, \mathrm{Argmin}\, f)$$
$$f_k = f(x_k) - f_\star$$
$$g_k = \mathrm{dist}_2(0, \partial f(x_k))$$

$$g_{k+1} \leq g_k$$

$$g_k \leq \frac{f_0}{t} \cdot e^{-2(k-1)^2 t^2 / d_0^2}$$



P.R., Hanmin Li
A sharper theory of ball-proximal optimization: convergence and radius selection
arXiv:2609.35147, 2026

Fixed-Radius Theory: Gradient

$$d_k = \mathrm{dist}_2(x_k, \mathrm{Argmin}\, f)$$
$$f_k = f(x_k) - f_\star$$
$$g_k = \mathrm{dist}_2(0, \partial f(x_k))$$

$$g_{k+1} \leq g_k$$

$$g_k \leq \frac{f_0}{t} \cdot e^{-2(k-1)^2 t^2 / d_0^2}$$

$$k \geq 1 + \frac{d_0}{\sqrt{2}\, t} \sqrt{\ln \frac{f_0}{t\, \varepsilon}} \quad \Longrightarrow \quad g_k \leq \varepsilon$$



P.R., Hanmin Li
A sharper theory of ball-proximal optimization: convergence and radius selection
arXiv:2609.35147, 2026

Fixed-Radius Theory: Gradient

$$d_k = \mathrm{dist}_2(x_k, \mathrm{Argmin}\, f)$$
$$f_k = f(x_k) - f_\star$$
$$g_k = \mathrm{dist}_2(0, \partial f(x_k))$$

$$g_{k+1} \le g_k$$

$$g_k \le \frac{f_0}{t} \cdot e^{-2(k-1)^2 t^2 / d_0^2}$$

$$k \ge 1 + \frac{d_0}{\sqrt{2}\, t} \sqrt{\ln \frac{f_0}{t\, \varepsilon}} \quad \Longrightarrow \quad g_k \le \varepsilon$$

No dependence on a condition number!



P.R., Hanmin Li
A sharper theory of ball-proximal optimization: convergence and radius selection
arXiv:2609.35147, 2026

Arbitrary-Radius Theory: Convergence


Theorem. f proper, closed, convex, $x_0 \in \operatorname{dom} f$, any radii $t_k > 0$. Exactly one case applies:

P.R., Hanmin Li
A sharper theory of ball-proximal optimization: convergence and radius selection
arXiv:2609.35147, 2026 (*Theorem 4.1*)

Arbitrary-Radius Theory: Convergence

Theorem. f proper, closed, convex, $x_0 \in \operatorname{dom} f$, any radii $t_k > 0$. Exactly one case applies:

(i) $\sum_k t_k < \infty$:
$x_k \to \bar{x}$, $f(x_k) \downarrow f(\bar{x})$
$\bar{x}$ need *not* be a minimizer




$$f(x) = |x|, \quad x_0 = 2$$
$$t_k = 0.6 \cdot 2^{-k} \Rightarrow x_k \to 0.8$$

P.R., Hanmin Li
A sharper theory of ball-proximal optimization: convergence and radius selection
arXiv:2609.35147, 2026 (*Theorem 4.1*)

Arbitrary-Radius Theory: Convergence


Theorem. f proper, closed, convex, $x_0 \in \operatorname{dom} f$, any radii $t_k > 0$. Exactly one case applies:

(i) $\sum_k t_k < \infty$:
$x_k \to \bar{x}$, $f(x_k) \downarrow f(\bar{x})$
$\bar{x}$ need *not* be a minimizer



$f(x) = |x|$, $x_0 = 2$
$t_k = 0.6 \cdot 2^{-k} \Rightarrow x_k \to 0.8$

(ii) $\sum_k t_k = \infty$, $\operatorname{Argmin} f \neq \emptyset$:
$x_K \in \operatorname{Argmin} f$ for some $K < \infty$
finite termination




$f(x) = |x|$, $x_0 = 2$
$t_k = \frac{1}{k+1} \Rightarrow x_4 = 0$

P.R., Hanmin Li
A sharper theory of ball-proximal optimization: convergence and radius selection
arXiv:2609.35147, 2026 (*Theorem 4.1*)

Arbitrary-Radius Theory: Convergence


Theorem. f proper, closed, convex, $x_0 \in \operatorname{dom} f$, any radii $t_k > 0$. Exactly one case applies:

(i) $\sum_k t_k < \infty$:
$x_k \to \bar{x}$, $f(x_k) \downarrow f(\bar{x})$
$\bar{x}$ need *not* be a minimizer




$f(x) = |x|$, $x_0 = 2$
$t_k = 0.6 \cdot 2^{-k} \Rightarrow x_k \to 0.8$

(ii) $\sum_k t_k = \infty$, $\operatorname{Argmin} f \neq \emptyset$:
$x_K \in \operatorname{Argmin} f$ for some $K < \infty$
finite termination



$f(x) = |x|$, $x_0 = 2$
$t_k = \frac{1}{k+1} \Rightarrow x_4 = 0$

(iii) $\sum_k t_k = \infty$, $\operatorname{Argmin} f = \emptyset$:
$f(x_k) \downarrow \inf f$, $\|x_k\|_2 \to \infty$
also when $\inf f = -\infty$




$f(x) = e^x$, $x_0 = 1$
$t_k = \frac{1}{k+1} \Rightarrow x_k \to -\infty$

P.R., Hanmin Li
A sharper theory of ball-proximal optimization: convergence and radius selection
arXiv:2609.35147, 2026 (*Theorem 4.1*)

Arbitrary-Radius Theory: Convergence


Theorem. f proper, closed, convex, $x_0 \in \operatorname{dom} f$, any radii $t_k > 0$. Exactly one case applies:

(i) $\sum_k t_k < \infty$:
$x_k \to \bar{x}$, $f(x_k) \downarrow f(\bar{x})$
$\bar{x}$ need *not* be a minimizer




$f(x) = |x|$, $x_0 = 2$
$t_k = 0.6 \cdot 2^{-k} \Rightarrow x_k \to 0.8$

(ii) $\sum_k t_k = \infty$, $\operatorname{Argmin} f \neq \emptyset$:
$x_K \in \operatorname{Argmin} f$ for some $K < \infty$
finite termination



$f(x) = |x|$, $x_0 = 2$
$t_k = \frac{1}{k+1} \Rightarrow x_4 = 0$

(iii) $\sum_k t_k = \infty$, $\operatorname{Argmin} f = \emptyset$:
$f(x_k) \downarrow \inf f$, $\|x_k\|_2 \to \infty$
also when $\inf f = -\infty$



$f(x) = e^x$, $x_0 = 1$
$t_k = \frac{1}{k+1} \Rightarrow x_k \to -\infty$

In particular, $\sum_k t_k = \infty \implies f(x_k) \downarrow \inf f$; radii may shrink, e.g. $t_k = 1/(k+1)$.
Conversely, $\sum_k t_k = \infty$ is *necessary* for this, for every proper closed convex f and x_0 (see (i)).

P.R., Hanmin Li
A sharper theory of ball-proximal optimization: convergence and radius selection
arXiv:2609.35147, 2026 (*Theorem 4.1*)

Part 3
Broximal Point Method: Nonconvex Regime



Kaja Gruntkowska, Hanmin Li, Xun Qian, P.R.
Broximal alignment for global non-convex optimization
arXiv:2604.13483, 2026

2D Example 1: GD Gets Stuck


$$f(x) = \|x\|_2 + 10\sin\xi_1 + 10\sin\xi_2, \qquad x = (\xi_1, \xi_2) \in \mathbb{R}^2$$

2D version of the 1D example $|x| + 10\sin x$ (Gruntkowska et al., 2026)

2D Example 1: GD Gets Stuck

$$f(x) = \|x\|_2 + 10\sin\xi_1 + 10\sin\xi_2, \qquad x = (\xi_1, \xi_2) \in \mathbb{R}^2$$

2D version of the 1D example $|x| + 10\sin x$ (Gruntkowska et al., 2026)




level sets of f (darker = lower f)

2D Example 1: GD Gets Stuck

$$f(x) = \|x\|_2 + 10\sin\xi_1 + 10\sin\xi_2, \qquad x = (\xi_1, \xi_2) \in \mathbb{R}^2$$

2D version of the 1D example $|x| + 10\sin x$ (Gruntkowska et al., 2026)

non-optimal local minima



level sets of f (darker = lower f)


2D Example 1: GD Gets Stuck

$$f(x) = \|x\|_2 + 10\sin\xi_1 + 10\sin\xi_2, \qquad x = (\xi_1, \xi_2) \in \mathbb{R}^2$$

2D version of the 1D example $|x| + 10\sin x$ (Gruntkowska et al., 2026)

non-optimal local minima

Argmin f



level sets of f (darker = lower f)


2D Example 1: GD Gets Stuck

$$f(x) = \|x\|_2 + 10\sin\xi_1 + 10\sin\xi_2, \qquad x = (\xi_1, \xi_2) \in \mathbb{R}^2$$

2D version of the 1D example $|x| + 10\sin x$ (Gruntkowska et al., 2026)

non-optimal local minima

Argmin f




level sets of f (darker = lower f)

2D Example 1: GD Gets Stuck

$$f(x) = \|x\|_2 + 10\sin\xi_1 + 10\sin\xi_2, \qquad x = (\xi_1, \xi_2) \in \mathbb{R}^2$$


2D version of the 1D example $|x| + 10\sin x$ (Gruntkowska et al., 2026)



2D Example 1: GD Gets Stuck

$$f(x) = \|x\|_2 + 10\sin\xi_1 + 10\sin\xi_2, \qquad x = (\xi_1, \xi_2) \in \mathbb{R}^2$$


2D version of the 1D example $|x| + 10\sin x$ (Gruntkowska et al., 2026)



2D Example 1: GD Gets Stuck

$$f(x) = \|x\|_2 + 10\sin\xi_1 + 10\sin\xi_2, \qquad x = (\xi_1, \xi_2) \in \mathbb{R}^2$$

2D version of the 1D example $|x| + 10\sin x$ (Gruntkowska et al., 2026)




2D Example 1: BPM Jumps Over Local Minima

$$x_{k+1} \in \mathrm{brox}_f^t(x_k) = \mathrm{Argmin}\,\{f(y) \;:\; \|y - x_k\|_2 \le t\}, \qquad t = 2\pi$$

BPM function values

$f(x_0) = \quad 24.90$

Argmin f



level sets of f (darker = lower f)


2D Example 1: BPM Jumps Over Local Minima

$$x_{k+1} \in \mathrm{brox}_f^t(x_k) = \mathrm{Argmin}\,\{f(y) \;:\; \|y - x_k\|_2 \le t\}, \qquad t = 2\pi$$

BPM function values

$f(x_0) = \quad 24.90$

Argmin f



level sets of f (darker = lower f)

2D Example 1: BPM Jumps Over Local Minima


$$x_{k+1} \in \mathrm{brox}_f^t(x_k) = \mathrm{Argmin}\,\{f(y) \;:\; \|y - x_k\|_2 \le t\}, \qquad t = 2\pi$$

BPM function values

$f(x_0) = \quad 24.90$

$f(x_1) = -8.09$

Argmin f



level sets of f (darker = lower f)

2D Example 1: BPM Jumps Over Local Minima


$$x_{k+1} \in \mathrm{brox}_f^t(x_k) = \mathrm{Argmin}\,\{f(y) \;:\; \|y - x_k\|_2 \le t\}, \qquad t = 2\pi$$

BPM function values

$f(x_0) = \quad 24.90$

$f(x_1) = -8.09$

Argmin f



level sets of f (darker = lower f)


2D Example 1: BPM Jumps Over Local Minima

$$x_{k+1} \in \mathrm{brox}_f^t(x_k) = \mathrm{Argmin}\,\{f(y) \;:\; \|y - x_k\|_2 \le t\}, \qquad t = 2\pi$$

BPM function values

$f(x_0) = \quad 24.90$
$f(x_1) = -8.09$
$f(x_2) = -13.39$

Argmin f



level sets of f (darker = lower f)


2D Example 1: BPM Jumps Over Local Minima

$$x_{k+1} \in \mathrm{brox}_f^t(x_k) = \mathrm{Argmin}\,\{f(y) \;:\; \|y - x_k\|_2 \le t\}, \qquad t = 2\pi$$

BPM function values

$f(x_0) = \quad 24.90$
$f(x_1) = -8.09$
$f(x_2) = -13.39$

Argmin f



level sets of f (darker = lower f)


2D Example 1: BPM Jumps Over Local Minima

$$x_{k+1} \in \mathrm{brox}_f^t(x_k) = \mathrm{Argmin}\,\{f(y) \;:\; \|y - x_k\|_2 \le t\}, \qquad t = 2\pi$$

BPM function values

$f(x_0) = 24.90$
$f(x_1) = -8.09$
$f(x_2) = -13.39$
$f(x_3) = -15.08$

Argmin f



level sets of f (darker = lower f)


2D Example 1: BPM Jumps Over Local Minima

$$x_{k+1} \in \mathrm{brox}_f^t(x_k) = \mathrm{Argmin}\,\{f(y) \;:\; \|y - x_k\|_2 \le t\}, \qquad t = 2\pi$$

BPM function values

$f(x_0) = 24.90$
$f(x_1) = -8.09$
$f(x_2) = -13.39$
$f(x_3) = -15.08$

Argmin f



level sets of f (darker = lower f)


2D Example 1: BPM Jumps Over Local Minima

$$x_{k+1} \in \mathrm{brox}_f^t(x_k) = \mathrm{Argmin}\,\{f(y) \;:\; \|y - x_k\|_2 \le t\}, \qquad t = 2\pi$$

BPM function values

$f(x_0) = 24.90$
$f(x_1) = -8.09$
$f(x_2) = -13.39$
$f(x_3) = -15.08$
$f(x_4) = -17.83 = f_\star$

Argmin f



level sets of f (darker = lower f)


2D Example 1: BPM Jumps Over Local Minima

$$x_{k+1} \in \mathrm{brox}_f^t(x_k) = \mathrm{Argmin}\,\{f(y) \;:\; \|y - x_k\|_2 \le t\}, \qquad t = 2\pi$$

BPM function values

$f(x_0) = 24.90$
$f(x_1) = -8.09$
$f(x_2) = -13.39$
$f(x_3) = -15.08$
$f(x_4) = -17.83 = f_\star$

Argmin f



Global minimum in 4 steps!

level sets of f (darker = lower f)

2D Example 2: GD Gets Stuck


$$f(x) = \varphi(\|x\|_2) + \eta\,\mathrm{ripple}(x), \qquad \varphi(r) = r + 20\,w(r)\sin^2(r/2)$$

radial analogue of the 1D example $|x| + 10\sin x$; $w(r) = 1 - e^{-r^2/64}$; $0 \le \mathrm{ripple} \le 1$, $\mathrm{ripple}(0) = 0$; $\eta = 5$

2D Example 2: GD Gets Stuck

$$f(x) = \varphi(\|x\|_2) + \eta\,\mathrm{ripple}(x), \qquad \varphi(r) = r + 20\,w(r)\sin^2(r/2)$$

radial analogue of the 1D example $|x| + 10\sin x$; $w(r) = 1 - e^{-r^2/64}$; $0 \le \mathrm{ripple} \le 1$, $\mathrm{ripple}(0) = 0$; $\eta = 5$




level sets of f
(darker = lower f)

2D Example 2: GD Gets Stuck

$$f(x) = \varphi(\|x\|_2) + \eta\,\mathrm{ripple}(x), \qquad \varphi(r) = r + 20\,w(r)\sin^2(r/2)$$

radial analogue of the 1D example $|x| + 10\sin x$; $w(r) = 1 - e^{-r^2/64}$; $0 \le \mathrm{ripple} \le 1$, $\mathrm{ripple}(0) = 0$; $\eta = 5$

60 non-optimal local minima



level sets of f
(darker = lower f)


2D Example 2: GD Gets Stuck

$$f(x) = \varphi(\|x\|_2) + \eta\,\mathrm{ripple}(x), \qquad \varphi(r) = r + 20\,w(r)\sin^2(r/2)$$

radial analogue of the 1D example $|x| + 10\sin x$; $w(r) = 1 - e^{-r^2/64}$; $0 \le \mathrm{ripple} \le 1$, $\mathrm{ripple}(0) = 0$; $\eta = 5$

60 non-optimal local minima

Argmin f



level sets of f
(darker = lower f)


2D Example 2: GD Gets Stuck

$$f(x) = \varphi(\|x\|_2) + \eta\,\mathrm{ripple}(x), \qquad \varphi(r) = r + 20\,w(r)\sin^2(r/2)$$

radial analogue of the 1D example $|x| + 10\sin x$; $w(r) = 1 - e^{-r^2/64}$; $0 \le \mathrm{ripple} \le 1$, $\mathrm{ripple}(0) = 0$; $\eta = 5$

60 non-optimal local minima

Argmin f



level sets of f
(darker = lower f)


2D Example 2: GD Gets Stuck

$$f(x) = \varphi(\|x\|_2) + \eta\,\mathrm{ripple}(x), \qquad \varphi(r) = r + 20\,w(r)\sin^2(r/2)$$

radial analogue of the 1D example $|x| + 10\sin x$; $w(r) = 1 - e^{-r^2/64}$; $0 \le \mathrm{ripple} \le 1$, $\mathrm{ripple}(0) = 0$; $\eta = 5$

60 non-optimal local minima

Argmin f



level sets of f
(darker = lower f)


2D Example 2: GD Gets Stuck

$$f(x) = \varphi(\|x\|_2) + \eta\,\mathrm{ripple}(x), \qquad \varphi(r) = r + 20\,w(r)\sin^2(r/2)$$

radial analogue of the 1D example $|x| + 10\sin x$; $w(r) = 1 - e^{-r^2/64}$; $0 \le \mathrm{ripple} \le 1$, $\mathrm{ripple}(0) = 0$; $\eta = 5$

60 non-optimal local minima

Argmin f



GD gets stuck

$\nabla f(\bar{x}) = 0$

$f(\bar{x}) = 26.30$

level sets of f
(darker = lower f)


2D Example 2: GD Gets Stuck

$$f(x) = \varphi(\|x\|_2) + \eta\,\mathrm{ripple}(x), \qquad \varphi(r) = r + 20\,w(r)\sin^2(r/2)$$

radial analogue of the 1D example $|x| + 10\sin x$; $w(r) = 1 - e^{-r^2/64}$; $0 \le \mathrm{ripple} \le 1$, $\mathrm{ripple}(0) = 0$; $\eta = 5$

60 non-optimal local minima

Argmin f



GD gets stuck

$\nabla f(\bar{x}) = 0$

$f(\bar{x}) = 26.30$

level sets of f
(darker = lower f)


Stationary, but far from optimal!

2D Example 2: BPM Jumps Over Local Minima

$$x_{k+1} \in \mathrm{brox}_f^t(x_k) = \mathrm{Argmin}\,\{f(y) \;:\; \|y - x_k\|_2 \le t\}, \qquad t = 2.5\pi$$

BPM function values
$f(x_0) = 41.64$

Argmin f



level sets of f
(darker = lower f)


2D Example 2: BPM Jumps Over Local Minima

$$x_{k+1} \in \mathrm{brox}_f^t(x_k) = \mathrm{Argmin}\,\{f(y) \;:\; \|y - x_k\|_2 \le t\}, \qquad t = 2.5\pi$$

BPM function values

$f(x_0) = 41.64$

Argmin f



level sets of f
(darker = lower f)

2D Example 2: BPM Jumps Over Local Minima


$$x_{k+1} \in \mathrm{brox}_f^t(x_k) = \mathrm{Argmin}\,\{f(y) \;:\; \|y - x_k\|_2 \le t\}, \qquad t = 2.5\pi$$

BPM function values

$f(x_0) = 41.64$

$f(x_1) = 19.89$

Argmin f



level sets of f
(darker = lower f)

2D Example 2: BPM Jumps Over Local Minima


$$x_{k+1} \in \mathrm{brox}_f^t(x_k) = \mathrm{Argmin}\,\{f(y) \;:\; \|y - x_k\|_2 \le t\}, \qquad t = 2.5\pi$$

BPM function values

$f(x_0) = 41.64$

$f(x_1) = 19.89$

Argmin f



level sets of f
(darker = lower f)


2D Example 2: BPM Jumps Over Local Minima

$$x_{k+1} \in \mathrm{brox}_f^t(x_k) = \mathrm{Argmin}\,\{f(y) \;:\; \|y - x_k\|_2 \le t\}, \qquad t = 2.5\pi$$

BPM function values

$f(x_0) = 41.64$
$f(x_1) = 19.89$
$f(x_2) = 13.32$

Argmin f



level sets of f
(darker = lower f)


2D Example 2: BPM Jumps Over Local Minima

$$x_{k+1} \in \mathrm{brox}_f^t(x_k) = \mathrm{Argmin}\,\{f(y) \;:\; \|y - x_k\|_2 \le t\}, \qquad t = 2.5\pi$$

BPM function values

$f(x_0) = 41.64$
$f(x_1) = 19.89$
$f(x_2) = 13.32$

Argmin f



level sets of f
(darker = lower f)


2D Example 2: BPM Jumps Over Local Minima

$$x_{k+1} \in \mathrm{brox}_f^t(x_k) = \mathrm{Argmin}\,\{f(y) \;:\; \|y - x_k\|_2 \le t\}, \qquad t = 2.5\pi$$

BPM function values

$f(x_0) = 41.64$
$f(x_1) = 19.89$
$f(x_2) = 13.32$
$f(x_3) = \;\; 6.41$

Argmin f



level sets of f
(darker = lower f)


2D Example 2: BPM Jumps Over Local Minima

$$x_{k+1} \in \mathrm{brox}_f^t(x_k) = \mathrm{Argmin}\,\{f(y) \;:\; \|y - x_k\|_2 \le t\}, \qquad t = 2.5\pi$$

BPM function values

$f(x_0) = 41.64$
$f(x_1) = 19.89$
$f(x_2) = 13.32$
$f(x_3) = \;\; 6.41$

Argmin f



level sets of f
(darker = lower f)


2D Example 2: BPM Jumps Over Local Minima

$$x_{k+1} \in \mathrm{brox}_f^t(x_k) = \mathrm{Argmin}\,\{f(y) \;:\; \|y - x_k\|_2 \le t\}, \qquad t = 2.5\pi$$

BPM function values

$f(x_0) = 41.64$
$f(x_1) = 19.89$
$f(x_2) = 13.32$
$f(x_3) = 6.41$
$f(x_4) = 0.00 = f_\star$

Argmin f



level sets of f
(darker = lower f)


2D Example 2: BPM Jumps Over Local Minima

$$x_{k+1} \in \mathrm{brox}_f^t(x_k) = \mathrm{Argmin}\,\{f(y) \;:\; \|y - x_k\|_2 \le t\}, \qquad t = 2.5\pi$$

BPM function values

$f(x_0) = 41.64$
$f(x_1) = 19.89$
$f(x_2) = 13.32$
$f(x_3) = 6.41$
$f(x_4) = 0.00 = f_\star$

Argmin f



level sets of f
(darker = lower f)

Global minimum
in 4 steps!


2D Example 2: BPM Jumps Over Local Minima

$$x_{k+1} \in \operatorname{brox}_f^t(x_k) = \operatorname{Argmin}\{f(y) \;:\; \|y - x_k\|_2 \le t\}, \qquad t = 2.5\pi$$

BPM function values

$f(x_0) = 41.64$
$f(x_1) = 19.89$
$f(x_2) = 13.32$
$f(x_3) = 6.41$
$f(x_4) = 0.00 = f_\star$

Argmin f



level sets of f
(darker = lower f)

Global minimum
in 4 steps!

$f \in \mathcal{F}_{\mathrm{I}}^{\mathrm{BA}}(t) \;\Rightarrow$ Theorem 4.1 applies
Brox-aligned: each step lands in the ball with diameter $[x_k, x_\star]$ (dashed: $k = 0$)

Brox-Aligned Nonconvex Functions

Broximal Alignment for Global Non-Convex Optimization

Table 1. **Comparison of convex and non-convex function classes. Non-Convex** = does not require f to be convex; **Non-Differentiable** = does not require f to be differentiable; **Stationary pts** = allows existence of stationary points that are not global minima; **Non-unique min** = allows existence of multiple global minima; **Disconnected** $\mathcal{X}_f$: the set of minimizers need not be connected.

Function class	Non-Convex	Non-Differentiable	Stationary pts	Non-unique min	Disconnected $\mathcal{X}_f$
Strongly convex	✗	✓	✗	✗	✗
Convex	✗	✓	✗	✓	✗
Pseudoconvex (Mangasarian, 1975)	✓	✗	✗	✓	✗
Quasiconvex (Arrow & Enthoven, 1961)	✓	✓	✓	✓	✗
Star-convex (Nesterov & Polyak, 2006)	✓	✗	✗	✓	✗
Quasar convex (Hardt et al., 2018)	✓	✗	✗	✓	✗
Aiming (Liu et al., 2023)	✓	✗	✗	✓	✗
Polyak-Łojasiewicz (Polyak, 1963; Łojasiewicz, 1963)	✓	✗	✗	✗	✗
Quadratic growth (Bonnans & Ioffe, 1995)	✓	✓	✓	✓	✓
$\mathcal{F}_{\mathbf{X}}^{\mathrm{BA}}(t)$ (ours)	✓	✓	✓	✓	✓




Figure 3. **Hierarchy of assumptions.** Solid arrows encode one-way implications, dashed lines denote mutual non-implications, and red arrows indicate conditions guaranteeing Broximal Alignment for any $t > 0$. For simplicity, we assume f is L-smooth (i.e., $\|\nabla f(x) - \nabla f(y)\| \leq L \|x - y\| \; \forall x, y \in \mathbb{R}^d$) with domain $\operatorname{dom} f = \mathbb{R}^d$. See the appendix and Karimi et al. (2016); Hinder et al. (2019); Khanh & Lara (2025) for proofs.

Brox-Aligned Nonconvex Functions

Broximal Alignment for Global Non-Convex Optimization

Table 1. **Comparison of convex and non-convex function classes. Non-Convex** = does not require f to be convex; **Non-Differentiable** = does not require f to be differentiable; **Stationary pts** = allows existence of stationary points that are not global minima; **Non-unique min** = allows existence of multiple global minima; **Disconnected** $\mathcal{X}_f$: the set of minimizers need not be connected.

Function class	Non-Convex	Non-Differentiable	Stationary pts	Non-unique min	Disconnected $\mathcal{X}_f$
Strongly convex	✗	✓	✗	✗	✗
Convex	✗	✓	✗	✓	✗
Pseudoconvex (Mangasarian, 1975)	✓	✗	✗	✓	✗
Quasiconvex (Arrow & Enthoven, 1961)	✓	✓	✓	✓	✗
Star-convex (Nesterov & Polyak, 2006)	✓	✗	✗	✓	✗
Quasar convex (Hardt et al., 2018)	✓	✗	✗	✓	✗
Aiming (Liu et al., 2023)	✓	✗	✗	✓	✗
Polyak-Łojasiewicz (Polyak, 1963; Łojasiewicz, 1963)	✓	✗	✗	✗	✗
Quadratic growth (Bonnans & Ioffe, 1995)	✓	✓	✓	✓	✓
$\mathcal{F}_{\mathbf{X}}^{\mathrm{BA}}(t)$ (ours)	✓	✓	✓	✓	✓




Figure 3. **Hierarchy of assumptions.** Solid arrows encode one-way implications, dashed lines denote mutual non-implications, and red arrows indicate conditions guaranteeing Broximal Alignment for any $t > 0$. For simplicity, we assume f is L-smooth (i.e., $\|\nabla f(x) - \nabla f(y)\| \leq L \|x - y\| \; \forall x, y \in \mathbb{R}^d$) with domain $\mathrm{dom}\, f = \mathbb{R}^d$. See the appendix and Karimi et al. (2016); Hinder et al. (2019); Khanh & Lara (2025) for proofs.

Kaja Gruntkowska, Hanmin Li, Xun Qian, P.R.

Broximal alignment for global non-convex optimization

arXiv:2604.13483, 2026

Part 4
Broximal Point Method: Practical Approximations

GD = Linearized BPM


$$x_{k+1} = \operatorname{argmin}\left\{ \underbrace{f(x_k) + \langle \nabla f(x_k), y - x_k \rangle}_{\text{linearization of } f \text{ at } x_k} \;:\; \|y - x_k\|_2 \le t_k \right\} = x_k - \gamma \nabla f(x_k) \qquad \text{if } t_k = \gamma \|\nabla f(x_k)\|_2$$

Example: $f(x) = \frac{1}{2}\langle a, x\rangle^2 + 2\langle b, x\rangle^2, \qquad a = (\cos 20^\circ, \sin 20^\circ), \;\; b = (-\sin 20^\circ, \cos 20^\circ)$

GD = Linearized BPM

$$x_{k+1} = \operatorname{argmin} \Big\{ \underbrace{f(x_k) + \langle \nabla f(x_k), y - x_k \rangle}_{\text{linearization of } f \text{ at } x_k} \; : \; \|y - x_k\|_2 \le t_k \Big\} = x_k - \gamma \nabla f(x_k) \qquad \text{if } t_k = \gamma \|\nabla f(x_k)\|_2$$

Example: $f(x) = \frac{1}{2}\langle a, x\rangle^2 + 2\langle b, x\rangle^2, \qquad a = (\cos 20^\circ, \sin 20^\circ), \;\; b = (-\sin 20^\circ, \cos 20^\circ)$




GD = Linearized BPM

$$x_{k+1} = \operatorname{argmin} \Big\{ \underbrace{f(x_k) + \langle \nabla f(x_k), y - x_k \rangle}_{\text{linearization of } f \text{ at } x_k} \; : \; \|y - x_k\|_2 \le t_k \Big\} = x_k - \gamma \nabla f(x_k) \qquad \text{if } t_k = \gamma \|\nabla f(x_k)\|_2$$

Example: $f(x) = \frac{1}{2}\langle a, x\rangle^2 + 2\langle b, x\rangle^2, \qquad a = (\cos 20^\circ, \sin 20^\circ), \;\; b = (-\sin 20^\circ, \cos 20^\circ)$

Linearize the objective inside the ball




GD = Linearized BPM

$$x_{k+1} = \operatorname{argmin}\left\{ \underbrace{f(x_k) + \langle \nabla f(x_k), y - x_k \rangle}_{\text{linearization of } f \text{ at } x_k} \;:\; \|y - x_k\|_2 \le t_k \right\} = x_k - \gamma \nabla f(x_k) \qquad \text{if } t_k = \gamma \|\nabla f(x_k)\|_2$$

Example: $f(x) = \frac{1}{2}\langle a, x\rangle^2 + 2\langle b, x\rangle^2, \qquad a = (\cos 20^\circ, \sin 20^\circ), \;\; b = (-\sin 20^\circ, \cos 20^\circ)$

Linearize the objective inside the ball

linearize at x_k: $f(y) \approx f(x_k) + \langle \nabla f(x_k), y - x_k \rangle$



GD = Linearized BPM

$$x_{k+1} = \operatorname{argmin}\Big\{ \underbrace{f(x_k) + \langle \nabla f(x_k), y - x_k \rangle}_{\text{linearization of } f \text{ at } x_k} \;:\; \|y - x_k\|_2 \le t_k \Big\} = x_k - \gamma \nabla f(x_k) \qquad \text{if } t_k = \gamma \|\nabla f(x_k)\|_2$$


Example: $f(x) = \frac{1}{2}\langle a, x\rangle^2 + 2\langle b, x\rangle^2, \qquad a = (\cos 20^\circ, \sin 20^\circ), \;\; b = (-\sin 20^\circ, \cos 20^\circ)$

Linearize the objective inside the ball

linearize at x_k: $f(y) \approx f(x_k) + \langle \nabla f(x_k), y - x_k \rangle$

$$\operatorname{argmin}\{\langle \nabla f(x_k), y\rangle : \|y - x_k\|_2 \le t_k\} = x_k - t_k \frac{\nabla f(x_k)}{\|\nabla f(x_k)\|_2}$$

$$t_k = \gamma \|\nabla f(x_k)\|_2 \implies x_{k+1} = x_k - \gamma \nabla f(x_k) \quad \text{(GD)}$$



GD = Linearized BPM

$$x_{k+1} = \operatorname{argmin}\left\{ \underbrace{f(x_k) + \langle \nabla f(x_k), y - x_k \rangle}_{\text{linearization of } f \text{ at } x_k} \;:\; \|y - x_k\|_2 \le t_k \right\} = x_k - \gamma \nabla f(x_k) \qquad \text{if } t_k = \gamma \|\nabla f(x_k)\|_2$$


Example: $f(x) = \frac{1}{2}\langle a, x\rangle^2 + 2\langle b, x\rangle^2, \qquad a = (\cos 20^\circ, \sin 20^\circ), \;\; b = (-\sin 20^\circ, \cos 20^\circ)$

Linearize the objective inside the ball

linearize at x_k: $f(y) \approx f(x_k) + \langle \nabla f(x_k), y - x_k \rangle$

$$\operatorname{argmin}\left\{ \langle \nabla f(x_k), y \rangle : \|y - x_k\|_2 \le t_k \right\} = x_k - t_k \frac{\nabla f(x_k)}{\|\nabla f(x_k)\|_2}$$

$$t_k = \gamma \|\nabla f(x_k)\|_2 \implies x_{k+1} = x_k - \gamma \nabla f(x_k) \quad \text{(GD)}$$



GD = Linearized BPM

$$x_{k+1} = \operatorname{argmin}\Big\{ \underbrace{f(x_k) + \langle \nabla f(x_k), y - x_k\rangle}_{\text{linearization of } f \text{ at } x_k} \;:\; \|y - x_k\|_2 \le t_k \Big\} = x_k - \gamma \nabla f(x_k) \qquad \text{if } t_k = \gamma \|\nabla f(x_k)\|_2$$


Example: $f(x) = \frac{1}{2}\langle a, x\rangle^2 + 2\langle b, x\rangle^2, \qquad a = (\cos 20^\circ, \sin 20^\circ),\ \ b = (-\sin 20^\circ, \cos 20^\circ)$

Linearize the objective inside the ball

linearize at x_k: $f(y) \approx f(x_k) + \langle \nabla f(x_k), y - x_k\rangle$

$$\operatorname{argmin}\{\langle \nabla f(x_k), y\rangle : \|y - x_k\|_2 \le t_k\} = x_k - t_k \frac{\nabla f(x_k)}{\|\nabla f(x_k)\|_2}$$

$$t_k = \gamma \|\nabla f(x_k)\|_2 \implies x_{k+1} = x_k - \gamma \nabla f(x_k) \quad \text{(GD)}$$



GD = Linearized BPM

$$x_{k+1} = \operatorname{argmin}\Big\{ \underbrace{f(x_k) + \langle \nabla f(x_k), y - x_k \rangle}_{\text{linearization of } f \text{ at } x_k} \;:\; \|y - x_k\|_2 \le t_k \Big\} = x_k - \gamma \nabla f(x_k) \qquad \text{if } t_k = \gamma \|\nabla f(x_k)\|_2$$


Example: $f(x) = \frac{1}{2}\langle a, x\rangle^2 + 2\langle b, x\rangle^2, \qquad a = (\cos 20^\circ, \sin 20^\circ),\ \ b = (-\sin 20^\circ, \cos 20^\circ)$

Linearize the objective inside the ball

linearize at x_k: $f(y) \approx f(x_k) + \langle \nabla f(x_k), y - x_k \rangle$

$$\operatorname{argmin}\{\langle \nabla f(x_k), y\rangle : \|y - x_k\|_2 \le t_k\} = x_k - t_k \frac{\nabla f(x_k)}{\|\nabla f(x_k)\|_2}$$

$$t_k = \gamma \|\nabla f(x_k)\|_2 \implies x_{k+1} = x_k - \gamma \nabla f(x_k) \quad \text{(GD)}$$



GD = Linearized BPM

$$x_{k+1} = \operatorname{argmin}\Big\{ \underbrace{f(x_k) + \langle \nabla f(x_k), y - x_k\rangle}_{\text{linearization of } f \text{ at } x_k} \;:\; \|y - x_k\|_2 \le t_k \Big\} = x_k - \gamma \nabla f(x_k) \qquad \text{if } t_k = \gamma \|\nabla f(x_k)\|_2$$


Example: $f(x) = \frac{1}{2}\langle a, x\rangle^2 + 2\langle b, x\rangle^2, \qquad a = (\cos 20^\circ, \sin 20^\circ),\ \ b = (-\sin 20^\circ, \cos 20^\circ)$

Linearize the objective inside the ball

linearize at x_k: $f(y) \approx f(x_k) + \langle \nabla f(x_k), y - x_k\rangle$

$$\operatorname{argmin}\{\langle \nabla f(x_k), y\rangle : \|y - x_k\|_2 \le t_k\} = x_k - t_k \frac{\nabla f(x_k)}{\|\nabla f(x_k)\|_2}$$

$$t_k = \gamma \|\nabla f(x_k)\|_2 \implies x_{k+1} = x_k - \gamma \nabla f(x_k) \quad \text{(GD)}$$



GD = Linearized BPM

$$x_{k+1} = \operatorname{argmin}\Big\{ \underbrace{f(x_k) + \langle \nabla f(x_k), y - x_k \rangle}_{\text{linearization of } f \text{ at } x_k} \;:\; \|y - x_k\|_2 \le t_k \Big\} = x_k - \gamma \nabla f(x_k) \qquad \text{if } t_k = \gamma \|\nabla f(x_k)\|_2$$


Example: $f(x) = \frac{1}{2}\langle a, x\rangle^2 + 2\langle b, x\rangle^2, \qquad a = (\cos 20^\circ, \sin 20^\circ), \;\; b = (-\sin 20^\circ, \cos 20^\circ)$

Linearize the objective inside the ball

linearize at x_k: $f(y) \approx f(x_k) + \langle \nabla f(x_k), y - x_k \rangle$

$$\operatorname{argmin}\{\langle \nabla f(x_k), y\rangle : \|y - x_k\|_2 \le t_k\} = x_k - t_k \frac{\nabla f(x_k)}{\|\nabla f(x_k)\|_2}$$

$$t_k = \gamma \|\nabla f(x_k)\|_2 \implies x_{k+1} = x_k - \gamma \nabla f(x_k) \quad \text{(GD)}$$



GD = Linearized BPM

$$x_{k+1} = \operatorname{argmin}\Big\{ \underbrace{f(x_k) + \langle \nabla f(x_k), y - x_k \rangle}_{\text{linearization of } f \text{ at } x_k} \;:\; \|y - x_k\|_2 \le t_k \Big\} = x_k - \gamma \nabla f(x_k) \qquad \text{if } t_k = \gamma \|\nabla f(x_k)\|_2$$


Example: $f(x) = \frac{1}{2}\langle a, x\rangle^2 + 2\langle b, x\rangle^2, \qquad a = (\cos 20^\circ, \sin 20^\circ),\ \ b = (-\sin 20^\circ, \cos 20^\circ)$

Linearize the objective inside the ball

linearize at x_k: $f(y) \approx f(x_k) + \langle \nabla f(x_k), y - x_k \rangle$

$$\operatorname{argmin}\{\langle \nabla f(x_k), y\rangle : \|y - x_k\|_2 \le t_k\} = x_k - t_k \frac{\nabla f(x_k)}{\|\nabla f(x_k)\|_2}$$

$$t_k = \gamma \|\nabla f(x_k)\|_2 \implies x_{k+1} = x_k - \gamma \nabla f(x_k) \quad \text{(GD)}$$



GD = Linearized BPM

$$x_{k+1} = \operatorname{argmin}\Big\{ \underbrace{f(x_k) + \langle \nabla f(x_k), y - x_k \rangle}_{\text{linearization of } f \text{ at } x_k} \;:\; \|y - x_k\|_2 \le t_k \Big\} = x_k - \gamma \nabla f(x_k) \qquad \text{if } t_k = \gamma \|\nabla f(x_k)\|_2$$

Example: $f(x) = \frac{1}{2}\langle a, x\rangle^2 + 2\langle b, x\rangle^2, \qquad a = (\cos 20^\circ, \sin 20^\circ),\ \ b = (-\sin 20^\circ, \cos 20^\circ)$


Linearize the objective inside the ball

linearize at x_k: $f(y) \approx f(x_k) + \langle \nabla f(x_k), y - x_k \rangle$

$$\operatorname{argmin}\{\langle \nabla f(x_k), y\rangle : \|y - x_k\|_2 \le t_k\} = x_k - t_k \frac{\nabla f(x_k)}{\|\nabla f(x_k)\|_2}$$

$$t_k = \gamma \|\nabla f(x_k)\|_2 \implies x_{k+1} = x_k - \gamma \nabla f(x_k) \quad \text{(GD)}$$

GD = BPM on the linearization of f,
with radius $t_k = \gamma \|\nabla f(x_k)\|_2$



SGD = Stochastically Linearized BPM


$$x_{k+1} = \operatorname{argmin}\Big\{ \underbrace{f_{i_k}(x_k) + \langle \nabla f_{i_k}(x_k), y - x_k \rangle}_{\text{stochastic linearization of } f \text{ at } x_k} \;:\; \|y - x_k\|_2 \le t_k \Big\} = x_k - \gamma \nabla f_{i_k}(x_k) \qquad \text{if } t_k = \gamma \|\nabla f_{i_k}(x_k)\|_2$$

Example: $f = \frac{1}{4}\sum_{i=1}^{4} f_i, \quad f_i(x) = \frac{1}{2}\langle a, x - c_i\rangle^2 + 2\langle b, x - c_i\rangle^2$ (a, b as on the GD slide)

SGD = Stochastically Linearized BPM

$$x_{k+1} = \operatorname{argmin}\Big\{ \underbrace{f_{i_k}(x_k) + \langle \nabla f_{i_k}(x_k), y - x_k \rangle}_{\text{stochastic linearization of } f \text{ at } x_k} \;:\; \|y - x_k\|_2 \le t_k \Big\} = x_k - \gamma \nabla f_{i_k}(x_k) \qquad \text{if } t_k = \gamma \|\nabla f_{i_k}(x_k)\|_2$$

Example: $f = \frac{1}{4}\sum_{i=1}^{4} f_i$, $\quad f_i(x) = \frac{1}{2}\langle a, x - c_i\rangle^2 + 2\langle b, x - c_i\rangle^2 \quad$ (a, b as on the GD slide)



SGD = Stochastically Linearized BPM


$$x_{k+1} = \operatorname{argmin}\Big\{ \underbrace{f_{i_k}(x_k) + \langle \nabla f_{i_k}(x_k), y - x_k\rangle}_{\text{stochastic linearization of } f \text{ at } x_k} \;:\; \|y - x_k\|_2 \le t_k \Big\} = x_k - \gamma \nabla f_{i_k}(x_k) \qquad \text{if } t_k = \gamma \|\nabla f_{i_k}(x_k)\|_2$$

Example: $f = \frac{1}{4}\sum_{i=1}^4 f_i$, $f_i(x) = \frac{1}{2}\langle a, x - c_i\rangle^2 + 2\langle b, x - c_i\rangle^2$ (a, b as on the GD slide)

Linearize a random summand inside the ball

$$\ell_k(y) := f_{i_k}(x_k) + \langle \nabla f_{i_k}(x_k), y - x_k\rangle$$

$$\mathbb{E}_{i_k}\big[\ell_k(y)\big] = f(x_k) + \langle \nabla f(x_k), y - x_k\rangle \quad \text{(unbiased)}$$



SGD = Stochastically Linearized BPM


$$x_{k+1} = \operatorname{argmin}\Big\{ \underbrace{f_{i_k}(x_k) + \langle \nabla f_{i_k}(x_k), y - x_k\rangle}_{\text{stochastic linearization of } f \text{ at } x_k} \;:\; \|y - x_k\|_2 \le t_k \Big\} = x_k - \gamma \nabla f_{i_k}(x_k) \qquad \text{if } t_k = \gamma \|\nabla f_{i_k}(x_k)\|_2$$

Example: $f = \frac{1}{4}\sum_{i=1}^4 f_i$, $f_i(x) = \frac{1}{2}\langle a, x - c_i\rangle^2 + 2\langle b, x - c_i\rangle^2$ (a, b as on the GD slide)

Linearize a random summand inside the ball

$$\ell_k(y) := f_{i_k}(x_k) + \langle \nabla f_{i_k}(x_k), y - x_k\rangle$$

$$\mathbb{E}_{i_k}\big[\ell_k(y)\big] = f(x_k) + \langle \nabla f(x_k), y - x_k\rangle \quad \text{(unbiased)}$$



orange: stochastic linearization

♦ $c_i = \operatorname{argmin} f_i$, ● $x_\star = \operatorname{argmin} f$

○ possible x_{k+1} (one per i), ◇ GD step

SGD = Stochastically Linearized BPM

$$x_{k+1} = \operatorname{argmin}\Big\{ \underbrace{f_{i_k}(x_k) + \langle \nabla f_{i_k}(x_k), y - x_k\rangle}_{\text{stochastic linearization of } f \text{ at } x_k} \;:\; \|y - x_k\|_2 \le t_k \Big\} = x_k - \gamma \nabla f_{i_k}(x_k) \qquad \text{if } t_k = \gamma \|\nabla f_{i_k}(x_k)\|_2$$


Example: $f = \frac{1}{4}\sum_{i=1}^4 f_i, \quad f_i(x) = \frac{1}{2}\langle a, x - c_i\rangle^2 + 2\langle b, x - c_i\rangle^2$ (a, b as on the GD slide)

Linearize a random summand inside the ball

$$\ell_k(y) := f_{i_k}(x_k) + \langle \nabla f_{i_k}(x_k), y - x_k\rangle$$

$$\mathbb{E}_{i_k}\big[\ell_k(y)\big] = f(x_k) + \langle \nabla f(x_k), y - x_k\rangle \quad \text{(unbiased)}$$

$$t_k = \gamma\|\nabla f_{i_k}(x_k)\|_2 \implies x_{k+1} = x_k - \gamma \nabla f_{i_k}(x_k) \quad \text{(SGD)}$$



SGD = Stochastically Linearized BPM

$$x_{k+1} = \operatorname{argmin}\Big\{ \underbrace{f_{i_k}(x_k) + \langle \nabla f_{i_k}(x_k), y - x_k\rangle}_{\text{stochastic linearization of } f \text{ at } x_k} \;:\; \|y - x_k\|_2 \le t_k \Big\} = x_k - \gamma \nabla f_{i_k}(x_k) \qquad \text{if } t_k = \gamma \|\nabla f_{i_k}(x_k)\|_2$$


Example: $f = \frac{1}{4}\sum_{i=1}^4 f_i$, $\quad f_i(x) = \frac{1}{2}\langle a, x - c_i\rangle^2 + 2\langle b, x - c_i\rangle^2$ $\quad$ (a, b as on the GD slide)

Linearize a random summand inside the ball

$$\ell_k(y) := f_{i_k}(x_k) + \langle \nabla f_{i_k}(x_k), y - x_k\rangle$$

$$\mathbb{E}_{i_k}\big[\ell_k(y)\big] = f(x_k) + \langle \nabla f(x_k), y - x_k\rangle \quad \text{(unbiased)}$$

$$t_k = \gamma \|\nabla f_{i_k}(x_k)\|_2 \implies x_{k+1} = x_k - \gamma \nabla f_{i_k}(x_k) \quad \text{(SGD)}$$



SGD = Stochastically Linearized BPM

$$x_{k+1} = \operatorname{argmin}\Big\{ \underbrace{f_{i_k}(x_k) + \langle \nabla f_{i_k}(x_k), y - x_k\rangle}_{\text{stochastic linearization of } f \text{ at } x_k} \;:\; \|y - x_k\|_2 \le t_k \Big\} = x_k - \gamma \nabla f_{i_k}(x_k) \qquad \text{if } t_k = \gamma \|\nabla f_{i_k}(x_k)\|_2$$


Example: $f = \frac{1}{4}\sum_{i=1}^4 f_i$, $\quad f_i(x) = \frac{1}{2}\langle a, x - c_i\rangle^2 + 2\langle b, x - c_i\rangle^2 \quad (a, b \text{ as on the GD slide})$

Linearize a random summand inside the ball

$$\ell_k(y) := f_{i_k}(x_k) + \langle \nabla f_{i_k}(x_k), y - x_k\rangle$$

$$\mathbb{E}_{i_k}\big[\ell_k(y)\big] = f(x_k) + \langle \nabla f(x_k), y - x_k\rangle \quad \text{(unbiased)}$$

$$t_k = \gamma \|\nabla f_{i_k}(x_k)\|_2 \implies x_{k+1} = x_k - \gamma \nabla f_{i_k}(x_k) \quad \text{(SGD)}$$



orange: stochastic linearization

♦ $c_i = \operatorname{argmin} f_i$, ● $x_\star = \operatorname{argmin} f$

○ possible x_{k+1} (one per i), ◇ GD step

SGD = Stochastically Linearized BPM

$$x_{k+1} = \operatorname{argmin}\Big\{ \underbrace{f_{i_k}(x_k) + \langle \nabla f_{i_k}(x_k), y - x_k\rangle}_{\text{stochastic linearization of } f \text{ at } x_k} \;:\; \|y - x_k\|_2 \le t_k \Big\} = x_k - \gamma \nabla f_{i_k}(x_k) \qquad \text{if } t_k = \gamma \|\nabla f_{i_k}(x_k)\|_2$$


Example: $f = \frac{1}{4}\sum_{i=1}^4 f_i$, $\quad f_i(x) = \frac{1}{2}\langle a, x - c_i\rangle^2 + 2\langle b, x - c_i\rangle^2$ $\quad$ (a, b as on the GD slide)

Linearize a random summand inside the ball

$$\ell_k(y) := f_{i_k}(x_k) + \langle \nabla f_{i_k}(x_k), y - x_k\rangle$$

$$\mathbb{E}_{i_k}\big[\ell_k(y)\big] = f(x_k) + \langle \nabla f(x_k), y - x_k\rangle \quad \text{(unbiased)}$$

$$t_k = \gamma\|\nabla f_{i_k}(x_k)\|_2 \implies x_{k+1} = x_k - \gamma \nabla f_{i_k}(x_k) \quad \text{(SGD)}$$



SGD = Stochastically Linearized BPM

$$x_{k+1} = \operatorname{argmin}\Big\{ \underbrace{f_{i_k}(x_k) + \langle \nabla f_{i_k}(x_k), y - x_k\rangle}_{\text{stochastic linearization of } f \text{ at } x_k} \;:\; \|y - x_k\|_2 \le t_k \Big\} = x_k - \gamma \nabla f_{i_k}(x_k) \qquad \text{if } t_k = \gamma \|\nabla f_{i_k}(x_k)\|_2$$


Example: $f = \frac{1}{4}\sum_{i=1}^4 f_i$, $f_i(x) = \frac{1}{2}\langle a, x - c_i\rangle^2 + 2\langle b, x - c_i\rangle^2$ (a, b as on the GD slide)

Linearize a random summand inside the ball

$$\ell_k(y) := f_{i_k}(x_k) + \langle \nabla f_{i_k}(x_k), y - x_k\rangle$$

$$\mathbb{E}_{i_k}\big[\ell_k(y)\big] = f(x_k) + \langle \nabla f(x_k), y - x_k\rangle \quad \text{(unbiased)}$$

$$t_k = \gamma\|\nabla f_{i_k}(x_k)\|_2 \implies x_{k+1} = x_k - \gamma \nabla f_{i_k}(x_k) \quad \text{(SGD)}$$



SGD = Stochastically Linearized BPM

$$x_{k+1} = \operatorname{argmin}\Big\{ \underbrace{f_{i_k}(x_k) + \langle \nabla f_{i_k}(x_k), y - x_k\rangle}_{\text{stochastic linearization of } f \text{ at } x_k} \;:\; \|y - x_k\|_2 \le t_k \Big\} = x_k - \gamma \nabla f_{i_k}(x_k) \qquad \text{if } t_k = \gamma \|\nabla f_{i_k}(x_k)\|_2$$


Example: $f = \frac{1}{4}\sum_{i=1}^4 f_i$, $\quad f_i(x) = \frac{1}{2}\langle a, x - c_i\rangle^2 + 2\langle b, x - c_i\rangle^2 \quad (a, b \text{ as on the GD slide})$

Linearize a random summand inside the ball

$$\ell_k(y) := f_{i_k}(x_k) + \langle \nabla f_{i_k}(x_k), y - x_k\rangle$$

$$\mathbb{E}_{i_k}\big[\ell_k(y)\big] = f(x_k) + \langle \nabla f(x_k), y - x_k\rangle \quad \text{(unbiased)}$$

$$t_k = \gamma \|\nabla f_{i_k}(x_k)\|_2 \implies x_{k+1} = x_k - \gamma \nabla f_{i_k}(x_k) \quad \text{(SGD)}$$



SGD = Stochastically Linearized BPM

$$x_{k+1} = \operatorname{argmin}\Big\{ \underbrace{f_{i_k}(x_k) + \langle \nabla f_{i_k}(x_k), y - x_k\rangle}_{\text{stochastic linearization of } f \text{ at } x_k} \;:\; \|y - x_k\|_2 \le t_k \Big\} = x_k - \gamma \nabla f_{i_k}(x_k) \qquad \text{if } t_k = \gamma \|\nabla f_{i_k}(x_k)\|_2$$


Example: $f = \frac{1}{4}\sum_{i=1}^4 f_i$, $\quad f_i(x) = \frac{1}{2}\langle a, x - c_i\rangle^2 + 2\langle b, x - c_i\rangle^2$ $\quad (a, b$ as on the GD slide$)$

Linearize a random summand inside the ball

$$\ell_k(y) := f_{i_k}(x_k) + \langle \nabla f_{i_k}(x_k), y - x_k\rangle$$

$$\mathbb{E}_{i_k}\big[\ell_k(y)\big] = f(x_k) + \langle \nabla f(x_k), y - x_k\rangle \quad \text{(unbiased)}$$

$$t_k = \gamma\|\nabla f_{i_k}(x_k)\|_2 \implies x_{k+1} = x_k - \gamma \nabla f_{i_k}(x_k) \quad \text{(SGD)}$$



SGD = Stochastically Linearized BPM

$$x_{k+1} = \operatorname{argmin}\Big\{ \underbrace{f_{i_k}(x_k) + \langle \nabla f_{i_k}(x_k), y - x_k\rangle}_{\text{stochastic linearization of } f \text{ at } x_k} \;:\; \|y - x_k\|_2 \le t_k \Big\} = x_k - \gamma \nabla f_{i_k}(x_k) \qquad \text{if } t_k = \gamma \|\nabla f_{i_k}(x_k)\|_2$$

Example: $f = \frac{1}{4}\sum_{i=1}^4 f_i$, $\quad f_i(x) = \frac{1}{2}\langle a, x - c_i\rangle^2 + 2\langle b, x - c_i\rangle^2$ $\quad (a, b$ as on the GD slide$)$


Linearize a random summand inside the ball

$$\ell_k(y) := f_{i_k}(x_k) + \langle \nabla f_{i_k}(x_k), y - x_k\rangle$$

$$\mathbb{E}_{i_k}\big[\ell_k(y)\big] = f(x_k) + \langle \nabla f(x_k), y - x_k\rangle \quad \text{(unbiased)}$$

$$t_k = \gamma\|\nabla f_{i_k}(x_k)\|_2 \implies x_{k+1} = x_k - \gamma \nabla f_{i_k}(x_k) \quad \text{(SGD)}$$

SGD = BPM on a random, unbiased linearization of f.
The radius does not vanish at $x_\star$: SGD hovers around $x_\star$.



orange: stochastic linearization
♦ $c_i = \operatorname{argmin} f_i$, ● $x_\star = \operatorname{argmin} f$
○ possible x_{k+1} (one per i), ◇ GD step

SGD with Spectral-Norm Balls

$$X_{k+1} \in \operatorname{Argmin}\left\{ \underbrace{f_{i_k}(X_k) + \langle G_k, Y - X_k \rangle}_{\text{stochastic linearization of } f \text{ at } X_k} \;:\; \|Y - X_k\|_{2\to 2} \le t_k \right\} \ni X_k - t_k U_k V_k^\top$$

Setting: $X \in \mathbb{R}^{m\times n}$, $\langle A, B\rangle = \operatorname{tr}(A^\top B)$, $G_k := \nabla f_{i_k}(X_k) = U_k \Sigma_k V_k^\top$ (reduced SVD)



Kaja Gruntkowska, P.R.
Non-Euclidean broximal point method: a blueprint for geometry-aware optimization
arXiv:2510.00823, 2025

SGD with Spectral-Norm Balls


$$X_{k+1} \in \operatorname{Argmin}\left\{ \underbrace{f_{i_k}(X_k) + \langle G_k, Y - X_k\rangle}_{\text{stochastic linearization of } f \text{ at } X_k} \;:\; \|Y - X_k\|_{2\to 2} \le t_k \right\} \ni X_k - t_k U_k V_k^\top$$

Setting: $X \in \mathbb{R}^{m\times n}$, $\langle A, B\rangle = \operatorname{tr}(A^\top B)$, $G_k := \nabla f_{i_k}(X_k) = U_k \Sigma_k V_k^\top$ (reduced SVD)

Linearize, then minimize over a spectral ball

$\|D\|_F = \|\sigma(D)\|_2$, $\|D\|_{2\to 2} = \|\sigma(D)\|_\infty$

$\|D\|_* = \|\sigma(D)\|_1$ (nuclear norm)



green: $\|D\|_{2\to 2} \le t_k$
dashed: $\|D\|_F \le t_k$
orange: $\langle G_k, \cdot\rangle$
$\sigma(G_k) = (3, 0.6)$

Kaja Gruntkowska, P.R.
Non-Euclidean broximal point method: a blueprint for geometry-aware optimization
arXiv:2510.00823, 2025

SGD with Spectral-Norm Balls

$$X_{k+1} \in \operatorname{Argmin}\left\{ \underbrace{f_{i_k}(X_k) + \langle G_k, Y - X_k\rangle}_{\text{stochastic linearization of } f \text{ at } X_k} \;:\; \|Y - X_k\|_{2\to 2} \le t_k \right\} \ni X_k - t_k U_k V_k^\top$$


Setting: $X \in \mathbb{R}^{m\times n}$, $\langle A, B\rangle = \operatorname{tr}(A^\top B)$, $G_k := \nabla f_{i_k}(X_k) = U_k \Sigma_k V_k^\top$ (reduced SVD)

Linearize, then minimize over a spectral ball

$\|D\|_F = \|\sigma(D)\|_2, \quad \|D\|_{2\to 2} = \|\sigma(D)\|_\infty$

$\|D\|_* = \|\sigma(D)\|_1$ (nuclear norm)

Frobenius ball: $D = -t_k G_k / \|G_k\|_F$ (SGD)



green: $\|D\|_{2\to 2} \le t_k$
dashed: $\|D\|_F \le t_k$
orange: $\langle G_k, \cdot\rangle$
$\sigma(G_k) = (3, 0.6)$

Kaja Gruntkowska, P.R.
Non-Euclidean broximal point method: a blueprint for geometry-aware optimization
arXiv:2510.00823, 2025

SGD with Spectral-Norm Balls

$$X_{k+1} \in \operatorname{Argmin}\left\{ \underbrace{f_{i_k}(X_k) + \langle G_k, Y - X_k\rangle}_{\text{stochastic linearization of } f \text{ at } X_k} \;:\; \|Y - X_k\|_{2\to 2} \le t_k \right\} \ni X_k - t_k U_k V_k^\top$$

Setting: $X \in \mathbb{R}^{m\times n}$, $\langle A, B\rangle = \operatorname{tr}(A^\top B)$, $G_k := \nabla f_{i_k}(X_k) = U_k \Sigma_k V_k^\top$ (reduced SVD)


Linearize, then minimize over a spectral ball

$\|D\|_F = \|\sigma(D)\|_2, \quad \|D\|_{2\to 2} = \|\sigma(D)\|_\infty$

$\|D\|_* = \|\sigma(D)\|_1$ (nuclear norm)

Frobenius ball: $D = -t_k\, G_k / \|G_k\|_F$ (SGD)

spectral ball: $D = -t_k\, U_k V_k^\top$ (all $\sigma_i(D) = t_k$)



green: $\|D\|_{2\to 2} \le t_k$
dashed: $\|D\|_F \le t_k$
orange: $\langle G_k, \cdot\rangle$
$\sigma(G_k) = (3, 0.6)$

Kaja Gruntkowska, P.R.
Non-Euclidean broximal point method: a blueprint for geometry-aware optimization
arXiv:2510.00823, 2025

SGD with Spectral-Norm Balls

$$X_{k+1} \in \operatorname{Argmin}\left\{ \underbrace{f_{i_k}(X_k) + \langle G_k, Y - X_k\rangle}_{\text{stochastic linearization of } f \text{ at } X_k} \;:\; \|Y - X_k\|_{2\to 2} \le t_k \right\} \ni X_k - t_k U_k V_k^\top$$

Setting: $X \in \mathbb{R}^{m\times n}$, $\langle A, B\rangle = \operatorname{tr}(A^\top B)$, $G_k := \nabla f_{i_k}(X_k) = U_k \Sigma_k V_k^\top$ (reduced SVD)

Linearize, then minimize over a spectral ball


$\|D\|_F = \|\sigma(D)\|_2, \quad \|D\|_{2\to 2} = \|\sigma(D)\|_\infty$

$\|D\|_* = \|\sigma(D)\|_1$ (nuclear norm)

Frobenius ball: $D = -t_k\, G_k / \|G_k\|_F$ (SGD)

spectral ball: $D = -t_k\, U_k V_k^\top$ (all $\sigma_i(D) = t_k$)

$t_k = \gamma \|G_k\|_* \implies X_{k+1} = X_k - \gamma \|G_k\|_*\, U_k V_k^\top$



green: $\|D\|_{2\to 2} \le t_k$
dashed: $\|D\|_F \le t_k$
orange: $\langle G_k, \cdot\rangle$
$\sigma(G_k) = (3, 0.6)$

Kaja Gruntkowska, P.R.
Non-Euclidean broximal point method: a blueprint for geometry-aware optimization
arXiv:2510.00823, 2025

SGD with Spectral-Norm Balls

$$X_{k+1} \in \operatorname{Argmin}\left\{ \underbrace{f_{i_k}(X_k) + \langle G_k, Y - X_k\rangle}_{\text{stochastic linearization of } f \text{ at } X_k} \;:\; \|Y - X_k\|_{2\to 2} \le t_k \right\} \ni X_k - t_k U_k V_k^\top$$

Setting: $X \in \mathbb{R}^{m\times n}$, $\langle A, B\rangle = \operatorname{tr}(A^\top B)$, $G_k := \nabla f_{i_k}(X_k) = U_k \Sigma_k V_k^\top$ (reduced SVD)

Linearize, then minimize over a spectral ball

$\|D\|_F = \|\sigma(D)\|_2, \quad \|D\|_{2\to 2} = \|\sigma(D)\|_\infty$


$\|D\|_* = \|\sigma(D)\|_1$ (nuclear norm)

Frobenius ball: $D = -t_k\, G_k / \|G_k\|_F$ (SGD)

spectral ball: $D = -t_k\, U_k V_k^\top$ (all $\sigma_i(D) = t_k$)

$t_k = \gamma \|G_k\|_* \implies X_{k+1} = X_k - \gamma \|G_k\|_*\, U_k V_k^\top$

Same recipe as SGD, only the ball changes:
the step is the orthogonalized gradient $-t_k U_k V_k^\top$.



green: $\|D\|_{2\to 2} \le t_k$
dashed: $\|D\|_F \le t_k$
orange: $\langle G_k, \cdot\rangle$
$\sigma(G_k) = (3, 0.6)$

Kaja Gruntkowska, P.R.
Non-Euclidean broximal point method: a blueprint for geometry-aware optimization
arXiv:2510.00823, 2025

Muon: Adding Momentum and Newton–Schulz

$$M_k = \beta M_{k-1} + (1-\beta)\, G_k, \qquad X_{k+1} = X_k - \eta\, \mathrm{NS}_5(M_k) \;\approx\; X_k - \eta\, U_k V_k^\top \qquad \text{(Jordan et al., 2024)}$$

$$M_k = U_k \Sigma_k V_k^\top \text{ (reduced SVD)}: \qquad X_k - \eta\, U_k V_k^\top \;\in\; \mathrm{Argmin}\left\{ \langle M_k, Y - X_k \rangle \;:\; \|Y - X_k\|_{2\to 2} \le \eta \right\}$$



Keller Jordan, Yuchen Jin, Vlado Boza, Jiacheng You, Franz Cesista, Laker Newhouse, Jeremy Bernstein
Muon: An optimizer for hidden layers in neural networks
kellerjordan.github.io/posts/muon, 2024


Muon: Adding Momentum and Newton–Schulz

$$M_k = \beta M_{k-1} + (1-\beta)\, G_k, \qquad X_{k+1} = X_k - \eta\, \mathrm{NS}_5(M_k) \;\approx\; X_k - \eta\, U_k V_k^\top \qquad \text{(Jordan et al., 2024)}$$

$$M_k = U_k \Sigma_k V_k^\top \text{ (reduced SVD)}: \qquad X_k - \eta\, U_k V_k^\top \;\in\; \mathrm{Argmin}\left\{ \langle M_k, Y - X_k \rangle \;:\; \|Y - X_k\|_{2\to 2} \le \eta \right\}$$

Spectral SGD (previous slide) + three changes

1. momentum: linearize with M_k, not G_k



green: $\|D\|_{2\to 2} \le \eta$
dashed: $\|D\|_F \le \eta$
orange: $\langle M_k, \cdot \rangle$
purple: $-\eta Y_j$
$\sigma(M_k) = (3, 0.6)$

Keller Jordan, Yuchen Jin, Vlado Boza, Jiacheng You, Franz Cesista, Laker Newhouse, Jeremy Bernstein
Muon: An optimizer for hidden layers in neural networks
kellerjordan.github.io/posts/muon, 2024


Muon: Adding Momentum and Newton–Schulz

$$M_k = \beta M_{k-1} + (1-\beta)\, G_k, \qquad X_{k+1} = X_k - \eta\, \mathrm{NS}_5(M_k) \;\approx\; X_k - \eta\, U_k V_k^\top \qquad \text{(Jordan et al., 2024)}$$

$$M_k = U_k \Sigma_k V_k^\top \text{ (reduced SVD)}: \qquad X_k - \eta\, U_k V_k^\top \;\in\; \mathrm{Argmin}\left\{ \langle M_k, Y - X_k \rangle \;:\; \|Y - X_k\|_{2\to 2} \le \eta \right\}$$

Spectral SGD (previous slide) + three changes

1. momentum: linearize with M_k, not G_k
2. fixed radius: $t_k \equiv \eta$ (fixed-radius theory)



Keller Jordan, Yuchen Jin, Vlado Boza, Jiacheng You, Franz Cesista, Laker Newhouse, Jeremy Bernstein
Muon: An optimizer for hidden layers in neural networks
kellerjordan.github.io/posts/muon, 2024


Muon: Adding Momentum and Newton–Schulz

$$M_k = \beta M_{k-1} + (1-\beta)\, G_k, \qquad X_{k+1} = X_k - \eta\, \mathrm{NS}_5(M_k) \;\approx\; X_k - \eta\, U_k V_k^\top \qquad \text{(Jordan et al., 2024)}$$

$$M_k = U_k \Sigma_k V_k^\top \text{ (reduced SVD)}: \qquad X_k - \eta\, U_k V_k^\top \;\in\; \mathrm{Argmin}\left\{ \langle M_k, Y - X_k \rangle \;:\; \|Y - X_k\|_{2\to 2} \le \eta \right\}$$

Spectral SGD (previous slide) + three changes

1. momentum: linearize with M_k, not G_k
2. fixed radius: $t_k \equiv \eta$ (fixed-radius theory)
3. inexact ball step: $U_k V_k^\top \approx \mathrm{NS}_5(M_k)$



green: $\|D\|_{2\to 2} \le \eta$
dashed: $\|D\|_F \le \eta$
orange: $\langle M_k, \cdot \rangle$
purple: $-\eta Y_j$
$\sigma(M_k) = (3, 0.6)$

Keller Jordan, Yuchen Jin, Vlado Boza, Jiacheng You, Franz Cesista, Laker Newhouse, Jeremy Bernstein
Muon: An optimizer for hidden layers in neural networks
kellerjordan.github.io/posts/muon, 2024

Muon: Adding Momentum and Newton–Schulz


$$M_k = \beta M_{k-1} + (1-\beta)\, G_k, \qquad X_{k+1} = X_k - \eta\, \mathrm{NS}_5(M_k) \;\approx\; X_k - \eta\, U_k V_k^\top \qquad \text{(Jordan et al., 2024)}$$

$$M_k = U_k \Sigma_k V_k^\top \text{ (reduced SVD)}: \qquad X_k - \eta\, U_k V_k^\top \;\in\; \mathrm{Argmin}\left\{ \langle M_k, Y - X_k \rangle \;:\; \|Y - X_k\|_{2\to 2} \le \eta \right\}$$

Spectral SGD (previous slide) + three changes

1. momentum: linearize with M_k, not G_k
2. fixed radius: $t_k \equiv \eta$ (fixed-radius theory)
3. inexact ball step: $U_k V_k^\top \approx \mathrm{NS}_5(M_k)$

NS_5: 5 Newton–Schulz steps, $Y_0 \to Y_5 \approx U_k V_k^\top$



green: $\|D\|_{2\to 2} \le \eta$
dashed: $\|D\|_F \le \eta$
orange: $\langle M_k, \cdot \rangle$
purple: $-\eta Y_j$
$\sigma(M_k) = (3, 0.6)$

Keller Jordan, Yuchen Jin, Vlado Boza, Jiacheng You, Franz Cesista, Laker Newhouse, Jeremy Bernstein
Muon: An optimizer for hidden layers in neural networks
kellerjordan.github.io/posts/muon, 2024

Muon: Adding Momentum and Newton–Schulz

$$M_k = \beta M_{k-1} + (1-\beta)\, G_k, \qquad X_{k+1} = X_k - \eta\, \mathrm{NS}_5(M_k) \;\approx\; X_k - \eta\, U_k V_k^\top \qquad \text{(Jordan et al., 2024)}$$


$$M_k = U_k \Sigma_k V_k^\top \text{ (reduced SVD)}: \qquad X_k - \eta\, U_k V_k^\top \;\in\; \mathrm{Argmin}\left\{ \langle M_k, Y - X_k \rangle \;:\; \|Y - X_k\|_{2\to 2} \le \eta \right\}$$

Spectral SGD (previous slide) + three changes

1. momentum: linearize with M_k, not G_k
2. fixed radius: $t_k \equiv \eta$ (fixed-radius theory)
3. inexact ball step: $U_k V_k^\top \approx \mathrm{NS}_5(M_k)$

NS_5: 5 Newton–Schulz steps, $Y_0 \to Y_5 \approx U_k V_k^\top$

Muon ≈ BPM: *momentum* linearization, spectral ball of *fixed* radius η, ball step computed *inexactly*.



green: $\|D\|_{2\to 2} \le \eta$
dashed: $\|D\|_F \le \eta$
orange: $\langle M_k, \cdot \rangle$
purple: $-\eta Y_j$
$\sigma(M_k) = (3, 0.6)$

Keller Jordan, Yuchen Jin, Vlado Boza, Jiacheng You, Franz Cesista, Laker Newhouse, Jeremy Bernstein
Muon: An optimizer for hidden layers in neural networks
kellerjordan.github.io/posts/muon, 2024

BroxGD: Linearized BPM with Constraints

$$x_{k+1} \in \operatorname{Argmin} \Big\{ \underbrace{f(x_k) + \langle \nabla f(x_k), y - x_k \rangle}_{\text{linearization of } f \text{ at } x_k} : \quad \underbrace{y \in \mathcal{X}}_{\text{kept exactly}} \ , \ \|y - x_k\|_2 \le t_k \Big\}$$

Problem: $\min_{x \in \mathcal{X}} f(x)$, $\mathcal{X}$ closed, convex $\iff$ $\min_x f(x) + \iota_{\mathcal{X}}(x)$: BPM with only f linearized



P.R., Kaja Gruntkowska, Hanmin Li
Broximal gradient descent: a projection-free sister of projected gradient descent
arXiv:2605.08850, 2026



P.R., Ammar Mahran
Local LMO is Secretly a Projection Method!
arXiv:2609.33383, 2026


BroxGD: Linearized BPM with Constraints

$$x_{k+1} \in \operatorname{Argmin} \Big\{ \underbrace{f(x_k) + \langle \nabla f(x_k), y - x_k \rangle}_{\text{linearization of } f \text{ at } x_k} : \quad \underbrace{y \in \mathcal{X}}_{\text{kept exactly}} , \ \|y - x_k\|_2 \le t_k \Big\}$$

Problem: $\min_{x \in \mathcal{X}} f(x)$, $\mathcal{X}$ closed, convex $\iff$ $\min_x f(x) + \iota_{\mathcal{X}}(x)$: BPM with only f linearized

Between projected GD and Frank–Wolfe

local LMO on $\mathcal{X} \cap \mathcal{B}(x_k, t_k)$: no projection



P.R., Kaja Gruntkowska, Hanmin Li
Broximal gradient descent: a projection-free sister of projected gradient descent
arXiv:2605.08850, 2026

P.R., Ammar Mahran
Local LMO is Secretly a Projection Method!
arXiv:2609.33383, 2026

BroxGD: Linearized BPM with Constraints

$$x_{k+1} \in \operatorname{Argmin}\Big\{ \underbrace{f(x_k) + \langle \nabla f(x_k), y - x_k \rangle}_{\text{linearization of } f \text{ at } x_k} : \quad \underbrace{y \in \mathcal{X}}_{\text{kept exactly}} \ , \ \|y - x_k\|_2 \le t_k \Big\}$$


Problem: $\min_{x \in \mathcal{X}} f(x)$, $\mathcal{X}$ closed, convex $\iff$ $\min_x f(x) + \iota_{\mathcal{X}}(x)$: BPM with only f linearized

Between projected GD and Frank–Wolfe

local LMO on $\mathcal{X} \cap \mathcal{B}(x_k, t_k)$: no projection

ProjGD: soft ball, needs $\mathrm{Proj}_{\mathcal{X}}$

Frank–Wolfe: global LMO over $\mathcal{X}$



P.R., Kaja Gruntkowska, Hanmin Li
Broximal gradient descent: a projection-free sister of projected gradient descent
arXiv:2605.08850, 2026

P.R., Ammar Mahran
Local LMO is Secretly a Projection Method!
arXiv:2609.33383, 2026

BroxGD: Linearized BPM with Constraints

$$x_{k+1} \in \operatorname{Argmin}\Big\{ \underbrace{f(x_k) + \langle \nabla f(x_k), y - x_k \rangle}_{\text{linearization of } f \text{ at } x_k} : \quad \underbrace{y \in \mathcal{X}}_{\text{kept exactly}} \ , \ \|y - x_k\|_2 \le t_k \Big\}$$


Problem: $\min_{x \in \mathcal{X}} f(x)$, $\mathcal{X}$ closed, convex $\iff$ $\min_x f(x) + \iota_{\mathcal{X}}(x)$: BPM with only f linearized

Between projected GD and Frank–Wolfe

local LMO on $\mathcal{X} \cap \mathcal{B}(x_k, t_k)$: no projection

ProjGD: soft ball, needs $\mathrm{Proj}_{\mathcal{X}}$

Frank–Wolfe: global LMO over $\mathcal{X}$



P.R., Kaja Gruntkowska, Hanmin Li
Broximal gradient descent: a projection-free sister of projected gradient descent
arXiv:2605.08850, 2026

P.R., Ammar Mahran
Local LMO is Secretly a Projection Method!
arXiv:2609.33383, 2026

BroxGD: Linearized BPM with Constraints

$$x_{k+1} \in \operatorname{Argmin} \Big\{ \underbrace{f(x_k) + \langle \nabla f(x_k), y - x_k \rangle}_{\text{linearization of } f \text{ at } x_k} : \quad \underbrace{y \in \mathcal{X}}_{\text{kept exactly}} , \ \|y - x_k\|_2 \le t_k \Big\}$$

Problem: $\min_{x \in \mathcal{X}} f(x)$, $\mathcal{X}$ closed, convex $\iff$ $\min_x f(x) + \iota_{\mathcal{X}}(x)$: BPM with only f linearized


Between projected GD and Frank–Wolfe

local LMO on $\mathcal{X} \cap \mathcal{B}(x_k, t_k)$: no projection

ProjGD: soft ball, needs $\operatorname{Proj}_{\mathcal{X}}$

Frank–Wolfe: global LMO over $\mathcal{X}$

$$t_k \le \frac{\langle \nabla f(x_k), x_k - x_\star \rangle}{\|\nabla f(x_k)\|_2} \implies d_{k+1}^2 \le d_k^2 - {t_k}^2$$



P.R., Kaja Gruntkowska, Hanmin Li
Broximal gradient descent: a projection-free sister of projected gradient descent
arXiv:2605.08850, 2026

P.R., Ammar Mahran
Local LMO is Secretly a Projection Method!
arXiv:2609.33383, 2026

BroxGD: Linearized BPM with Constraints

$$x_{k+1} \in \operatorname{Argmin} \Big\{ \underbrace{f(x_k) + \langle \nabla f(x_k), y - x_k \rangle}_{\text{linearization of } f \text{ at } x_k} : \quad \underbrace{y \in \mathcal{X}}_{\text{kept exactly}} , \ \|y - x_k\|_2 \le t_k \Big\}$$

Problem: $\min_{x \in \mathcal{X}} f(x)$, $\mathcal{X}$ closed, convex $\iff$ $\min_x f(x) + \iota_{\mathcal{X}}(x)$: BPM with only f linearized

Between projected GD and Frank–Wolfe


local LMO on $\mathcal{X} \cap \mathcal{B}(x_k, t_k)$: no projection

ProjGD: soft ball, needs $\mathrm{Proj}_{\mathcal{X}}$

Frank–Wolfe: global LMO over $\mathcal{X}$

$$t_k \le \frac{\langle \nabla f(x_k), x_k - x_\star \rangle}{\|\nabla f(x_k)\|_2} \implies d_{k+1}^2 \le d_k^2 - t_k^2$$

orange: linearization
$t_k = f_k / \|\nabla f(x_k)\|_2$
(Polyak radius)



P.R., Kaja Gruntkowska, Hanmin Li
Broximal gradient descent: a projection-free sister of projected gradient descent
arXiv:2605.08850, 2026

P.R., Ammar Mahran
Local LMO is Secretly a Projection Method!
arXiv:2609.33383, 2026

BroxGD: Linearized BPM with Constraints

$$x_{k+1} \in \operatorname{Argmin} \Big\{ \underbrace{f(x_k) + \langle \nabla f(x_k), y - x_k \rangle}_{\text{linearization of } f \text{ at } x_k} : \quad \underbrace{y \in \mathcal{X}}_{\text{kept exactly}} , \ \|y - x_k\|_2 \le t_k \Big\}$$

Problem: $\min_{x \in \mathcal{X}} f(x)$, $\mathcal{X}$ closed, convex $\iff$ $\min_x f(x) + \iota_{\mathcal{X}}(x)$: BPM with only f linearized

Between projected GD and Frank–Wolfe


local LMO on $\mathcal{X} \cap \mathcal{B}(x_k, t_k)$: no projection

ProjGD: soft ball, needs $\mathrm{Proj}_{\mathcal{X}}$

Frank–Wolfe: global LMO over $\mathcal{X}$

$$t_k \le \frac{\langle \nabla f(x_k), x_k - x_\star \rangle}{\|\nabla f(x_k)\|_2} \implies d_{k+1}^2 \le d_k^2 - {t_k}^2$$

orange: linearization
$t_k = f_k / \|\nabla f(x_k)\|_2$
(Polyak radius)



P.R., Kaja Gruntkowska, Hanmin Li
Broximal gradient descent: a projection-free sister of projected gradient descent
arXiv:2605.08850, 2026

P.R., Ammar Mahran
Local LMO is Secretly a Projection Method!
arXiv:2609.33383, 2026

BroxGD: Linearized BPM with Constraints

$$x_{k+1} \in \operatorname{Argmin} \Big\{ \underbrace{f(x_k) + \langle \nabla f(x_k), y - x_k \rangle}_{\text{linearization of } f \text{ at } x_k} : \quad \underbrace{y \in \mathcal{X}}_{\text{kept exactly}} , \ \|y - x_k\|_2 \le t_k \Big\}$$

Problem: $\min_{x \in \mathcal{X}} f(x)$, $\mathcal{X}$ closed, convex $\iff$ $\min_x f(x) + \iota_{\mathcal{X}}(x)$: BPM with only f linearized

Between projected GD and Frank–Wolfe


local LMO on $\mathcal{X} \cap \mathcal{B}(x_k, t_k)$: no projection

ProjGD: soft ball, needs $\mathrm{Proj}_{\mathcal{X}}$

Frank–Wolfe: global LMO over $\mathcal{X}$

$$t_k \le \frac{\langle \nabla f(x_k), x_k - x_\star \rangle}{\|\nabla f(x_k)\|_2} \implies d_{k+1}^2 \le d_k^2 - {t_k}^2$$

orange: linearization
$t_k = f_k / \|\nabla f(x_k)\|_2$
(Polyak radius)



P.R., Kaja Gruntkowska, Hanmin Li
Broximal gradient descent: a projection-free sister of projected gradient descent
arXiv:2605.08850, 2026

P.R., Ammar Mahran
Local LMO is Secretly a Projection Method!
arXiv:2609.33383, 2026

BroxGD: Linearized BPM with Constraints

$$x_{k+1} \in \operatorname{Argmin} \Big\{ \underbrace{f(x_k) + \langle \nabla f(x_k), y - x_k \rangle}_{\text{linearization of } f \text{ at } x_k} : \quad \underbrace{y \in \mathcal{X}}_{\text{kept exactly}} , \ \|y - x_k\|_2 \le t_k \Big\}$$

Problem: $\min_{x \in \mathcal{X}} f(x)$, $\mathcal{X}$ closed, convex $\iff$ $\min_x f(x) + \iota_{\mathcal{X}}(x)$: BPM with only f linearized

Between projected GD and Frank–Wolfe


local LMO on $\mathcal{X} \cap \mathcal{B}(x_k, t_k)$: no projection

ProjGD: soft ball, needs $\mathrm{Proj}_{\mathcal{X}}$

Frank–Wolfe: global LMO over $\mathcal{X}$

$$t_k \le \frac{\langle \nabla f(x_k), x_k - x_\star \rangle}{\|\nabla f(x_k)\|_2} \implies d_{k+1}^2 \le d_k^2 - {t_k}^2$$

orange: linearization
$t_k = f_k / \|\nabla f(x_k)\|_2$
(Polyak radius)



P.R., Kaja Gruntkowska, Hanmin Li
Broximal gradient descent: a projection-free sister of projected gradient descent
arXiv:2605.08850, 2026

P.R., Ammar Mahran
Local LMO is Secretly a Projection Method!
arXiv:2609.33383, 2026

BroxGD: Linearized BPM with Constraints

$$x_{k+1} \in \operatorname{Argmin} \Big\{ \underbrace{f(x_k) + \langle \nabla f(x_k), y - x_k \rangle}_{\text{linearization of } f \text{ at } x_k} : \quad \underbrace{y \in \mathcal{X}}_{\text{kept exactly}} , \ \|y - x_k\|_2 \le t_k \Big\}$$

Problem: $\min_{x \in \mathcal{X}} f(x)$, $\mathcal{X}$ closed, convex $\iff$ $\min_x f(x) + \iota_{\mathcal{X}}(x)$: BPM with only f linearized

Between projected GD and Frank–Wolfe

local LMO on $\mathcal{X} \cap \mathcal{B}(x_k, t_k)$: no projection


ProjGD: soft ball, needs $\mathrm{Proj}_{\mathcal{X}}$

Frank–Wolfe: global LMO over $\mathcal{X}$

$$t_k \le \frac{\langle \nabla f(x_k), x_k - x_\star \rangle}{\|\nabla f(x_k)\|_2} \implies d_{k+1}^2 \le d_k^2 - t_k^2$$

BroxGD = BPM on $f + \iota_{\mathcal{X}}$ with only f linearized.

Smooth convex, $t_k = R/\sqrt{K}$: $f(\hat{x}_K) - f_\star \le LR^2/(2K)$.



P.R., Kaja Gruntkowska, Hanmin Li
Broximal gradient descent: a projection-free sister of projected gradient descent
arXiv:2605.08850, 2026

P.R., Ammar Mahran
Local LMO is Secretly a Projection Method!
arXiv:2609.33383, 2026

Part 5
Broximal Point Method: Four Formulations




P.R., Hanmin Li
A sharper theory of ball-proximal optimization: convergence and radius selection
arXiv:2609.35147, 2026

BPM = Projection onto Sublevel Sets

$$x_{k+1} \in \mathrm{brox}_f^t(x_k) \implies x_{k+1} = \mathrm{Proj}_{C_k}(x_k), \qquad C_k := \{y : f(y) \le f(x_{k+1})\}$$

BPM = Projection onto Sublevel Sets

$$x_{k+1} \in \mathrm{brox}_f^t(x_k) \implies x_{k+1} = \mathrm{Proj}_{C_k}(x_k), \qquad C_k := \{y : \ f(y) \le f(x_{k+1})\}$$




BPM = Projection onto Sublevel Sets

$$x_{k+1} \in \mathrm{brox}_f^t(x_k) \implies x_{k+1} = \mathrm{Proj}_{C_k}(x_k), \qquad C_k := \{y : \ f(y) \le f(x_{k+1})\}$$

Sublevel projection

$$\min_{\|y-x_k\|_2 \le t} f(y) = \alpha_k$$

$$\iff \min_{f(y) \le \alpha_k} \|y - x_k\|_2 = t$$



Argmin f

sublevel set C_k (orange)
level sets of f (gray)
(darker = lower f)


BPM = Projection onto Sublevel Sets

$$x_{k+1} \in \mathrm{brox}_f^t(x_k) \implies x_{k+1} = \mathrm{Proj}_{C_k}(x_k), \qquad C_k := \{y : \ f(y) \le f(x_{k+1})\}$$

Sublevel projection

$$\min_{\|y-x_k\|_2 \le t} f(y) = \alpha_k$$

$$\iff \min_{f(y) \le \alpha_k} \|y - x_k\|_2 = t$$



Argmin f

sublevel set C_k (orange)
level sets of f (gray)
(darker = lower f)

BPM = Projection onto Sublevel Sets

$$x_{k+1} \in \mathrm{brox}_f^t(x_k) \implies x_{k+1} = \mathrm{Proj}_{C_k}(x_k), \qquad C_k := \{y : \ f(y) \le f(x_{k+1})\}$$


Sublevel projection

$$\min_{\|y-x_k\|_2 \le t} f(y) = \alpha_k$$

$$\iff \min_{f(y) \le \alpha_k} \|y - x_k\|_2 = t$$

$$C_0 \supseteq C_1 \supseteq \cdots \supseteq \mathrm{Argmin}\, f$$

$$\implies d_{k+1}^2 \le d_k^2 - t^2$$



Argmin f

sublevel set C_k (orange)
level sets of f (gray)
(darker = lower f)

BPM = Projection onto Sublevel Sets

$$x_{k+1} \in \mathrm{brox}_f^t(x_k) \implies x_{k+1} = \mathrm{Proj}_{C_k}(x_k), \qquad C_k := \{y : \ f(y) \le f(x_{k+1})\}$$


Sublevel projection

$$\min_{\|y-x_k\|_2 \le t} f(y) = \alpha_k$$

$$\iff \min_{f(y) \le \alpha_k} \|y - x_k\|_2 = t$$

$$C_0 \supseteq C_1 \supseteq \cdots \supseteq \mathrm{Argmin}\, f$$

$$\implies d_{k+1}^2 \le d_k^2 - t^2$$



Argmin f

sublevel set C_k (orange)
level sets of f (gray)
(darker = lower f)

BPM = Projection onto Sublevel Sets

$$x_{k+1} \in \mathrm{brox}_f^t(x_k) \implies x_{k+1} = \mathrm{Proj}_{C_k}(x_k), \qquad C_k := \{y : \ f(y) \le f(x_{k+1})\}$$


Sublevel projection

$$\min_{\|y-x_k\|_2 \le t} f(y) = \alpha_k$$

$$\iff \min_{f(y) \le \alpha_k} \|y - x_k\|_2 = t$$

$$C_0 \supseteq C_1 \supseteq \cdots \supseteq \mathrm{Argmin}\, f$$

$$\implies d_{k+1}^2 \le d_k^2 - t^2$$



Argmin f

sublevel set C_k (orange)
level sets of f (gray)
(darker = lower f)

BPM = Projection onto Sublevel Sets

$$x_{k+1} \in \mathrm{brox}_f^t(x_k) \implies x_{k+1} = \mathrm{Proj}_{C_k}(x_k), \qquad C_k := \{y : \ f(y) \le f(x_{k+1})\}$$


Sublevel projection

$$\min_{\|y-x_k\|_2 \le t} f(y) = \alpha_k$$

$$\iff \min_{f(y) \le \alpha_k} \|y - x_k\|_2 = t$$

$$C_0 \supseteq C_1 \supseteq \cdots \supseteq \mathrm{Argmin}\, f$$

$$\implies d_{k+1}^2 \le d_k^2 - t^2$$



Argmin f

sublevel set C_k (orange)
level sets of f (gray)
(darker = lower f)

BPM = Projection onto Sublevel Sets

$$x_{k+1} \in \mathrm{brox}_f^t(x_k) \implies x_{k+1} = \mathrm{Proj}_{C_k}(x_k), \qquad C_k := \{y : \ f(y) \le f(x_{k+1})\}$$


Sublevel projection

$$\min_{\|y-x_k\|_2 \le t} f(y) = \alpha_k$$

$$\iff \min_{f(y) \le \alpha_k} \|y - x_k\|_2 = t$$

$$C_0 \supseteq C_1 \supseteq \cdots \supseteq \mathrm{Argmin}\, f$$

$$\implies d_{k+1}^2 \le d_k^2 - t^2$$



Argmin f

sublevel set C_k (orange)
level sets of f (gray)
(darker = lower f)

BPM = Projection onto Sublevel Sets

$$x_{k+1} \in \mathrm{brox}_f^t(x_k) \implies x_{k+1} = \mathrm{Proj}_{C_k}(x_k), \qquad C_k := \{y : \ f(y) \le f(x_{k+1})\}$$


Sublevel projection

$$\min_{\|y-x_k\|_2 \le t} f(y) = \alpha_k$$

$$\iff \min_{f(y) \le \alpha_k} \|y - x_k\|_2 = t$$

$$C_0 \supseteq C_1 \supseteq \cdots \supseteq \mathrm{Argmin}\, f$$

$$\implies d_{k+1}^2 \le d_k^2 - t^2$$



Argmin f

sublevel set C_k (orange)
level sets of f (gray)
(darker = lower f)

BPM = Projection onto Sublevel Sets

$$x_{k+1} \in \mathrm{brox}_f^t(x_k) \implies x_{k+1} = \mathrm{Proj}_{C_k}(x_k), \qquad C_k := \{y : \ f(y) \le f(x_{k+1})\}$$

Sublevel projection


$$\min_{\|y-x_k\|_2 \le t} f(y) = \alpha_k$$

$$\iff \min_{f(y) \le \alpha_k} \|y - x_k\|_2 = t$$

$$C_0 \supseteq C_1 \supseteq \cdots \supseteq \mathrm{Argmin}\, f$$

$$\implies d_{k+1}^2 \le d_k^2 - t^2$$

$$\alpha_3 = 0 = f_\star : \quad x_4 = \mathrm{Proj}_{\mathrm{Argmin}\, f}(x_3)$$



Argmin f


sublevel set C_k (orange)
level sets of f (gray)
(darker = lower f)

BPM = Implicit Normalized GD

$$x_{k+1} \in \operatorname{brox}_f^t(x_k) \implies x_{k+1} = x_k - t\,\frac{\nabla f(x_{k+1})}{\|\nabla f(x_{k+1})\|_2}$$

BPM = Implicit Normalized GD


$$x_{k+1} \in \operatorname{brox}_f^t(x_k) \implies x_{k+1} = x_k - t\,\frac{\nabla f(x_{k+1})}{\|\nabla f(x_{k+1})\|_2}$$



BPM = Implicit Normalized GD

$$x_{k+1} \in \mathrm{brox}_f^t(x_k) \implies x_{k+1} = x_k - t\,\frac{\nabla f(x_{k+1})}{\|\nabla f(x_{k+1})\|_2}$$

Implicit (BPM) vs explicit




BPM = Implicit Normalized GD

$$x_{k+1} \in \operatorname{brox}_f^t(x_k) \implies x_{k+1} = x_k - t\,\frac{\nabla f(x_{k+1})}{\|\nabla f(x_{k+1})\|_2}$$

Implicit (BPM) vs explicit

$$\hat{x}_{k+1} = x_k - t\,\frac{\nabla f(x_k)}{\|\nabla f(x_k)\|_2} \quad \text{(explicit)}$$



Argmin f

explicit normalized GD (orange)
level set through x_{k+1} (purple)
(darker = lower f)


BPM = Implicit Normalized GD

$$x_{k+1} \in \operatorname{brox}_f^t(x_k) \implies x_{k+1} = x_k - t\,\frac{\nabla f(x_{k+1})}{\|\nabla f(x_{k+1})\|_2}$$

Implicit (BPM) vs explicit

$$\hat{x}_{k+1} = x_k - t\,\frac{\nabla f(x_k)}{\|\nabla f(x_k)\|_2} \quad \text{(explicit)}$$

$$x_{k+1} = \operatorname{prox}_{\gamma_k f}(x_k), \quad \gamma_k = \frac{t}{\|\nabla f(x_{k+1})\|_2}$$



Argmin f

explicit normalized GD (orange)
level set through x_{k+1} (purple)
(darker = lower f)


BPM = Implicit Normalized GD

$$x_{k+1} \in \operatorname{brox}_f^t(x_k) \implies x_{k+1} = x_k - t\,\frac{\nabla f(x_{k+1})}{\|\nabla f(x_{k+1})\|_2}$$

Implicit (BPM) vs explicit

$$\hat{x}_{k+1} = x_k - t\,\frac{\nabla f(x_k)}{\|\nabla f(x_k)\|_2} \quad \text{(explicit)}$$

$$x_{k+1} = \operatorname{prox}_{\gamma_k f}(x_k), \quad \gamma_k = \frac{t}{\|\nabla f(x_{k+1})\|_2}$$



explicit normalized GD (orange)
level set through x_{k+1} (purple)
(darker = lower f)


BPM = Implicit Normalized GD

$$x_{k+1} \in \operatorname{brox}_f^t(x_k) \implies x_{k+1} = x_k - t\,\frac{\nabla f(x_{k+1})}{\|\nabla f(x_{k+1})\|_2}$$

Implicit (BPM) vs explicit

$$\hat{x}_{k+1} = x_k - t\,\frac{\nabla f(x_k)}{\|\nabla f(x_k)\|_2} \quad \text{(explicit)}$$

$$x_{k+1} = \operatorname{prox}_{\gamma_k f}(x_k), \quad \gamma_k = \frac{t}{\|\nabla f(x_{k+1})\|_2}$$



Argmin f

explicit normalized GD (orange)
level set through x_{k+1} (purple)
(darker = lower f)


BPM = Implicit Normalized GD

$$x_{k+1} \in \operatorname{brox}_f^t(x_k) \implies x_{k+1} = x_k - t\,\frac{\nabla f(x_{k+1})}{\|\nabla f(x_{k+1})\|_2}$$

Implicit (BPM) vs explicit

$$\hat{x}_{k+1} = x_k - t\,\frac{\nabla f(x_k)}{\|\nabla f(x_k)\|_2} \quad \text{(explicit)}$$

$$x_{k+1} = \operatorname{prox}_{\gamma_k f}(x_k), \quad \gamma_k = \frac{t}{\|\nabla f(x_{k+1})\|_2}$$



explicit normalized GD (orange)
level set through x_{k+1} (purple)
(darker = lower f)


BPM = Implicit Normalized GD

$$x_{k+1} \in \operatorname{brox}_f^t(x_k) \implies x_{k+1} = x_k - t\,\frac{\nabla f(x_{k+1})}{\|\nabla f(x_{k+1})\|_2}$$

Implicit (BPM) vs explicit

$$\hat{x}_{k+1} = x_k - t\,\frac{\nabla f(x_k)}{\|\nabla f(x_k)\|_2} \quad \text{(explicit)}$$

$$x_{k+1} = \operatorname{prox}_{\gamma_k f}(x_k), \quad \gamma_k = \frac{t}{\|\nabla f(x_{k+1})\|_2}$$



Argmin f

explicit normalized GD (orange)
level set through x_{k+1} (purple)
(darker = lower f)


BPM = Implicit Normalized GD

$$x_{k+1} \in \operatorname{brox}_f^t(x_k) \implies x_{k+1} = x_k - t\,\frac{\nabla f(x_{k+1})}{\|\nabla f(x_{k+1})\|_2}$$

Implicit (BPM) vs explicit

$$\hat{x}_{k+1} = x_k - t\,\frac{\nabla f(x_k)}{\|\nabla f(x_k)\|_2} \quad \text{(explicit)}$$

$$x_{k+1} = \operatorname{prox}_{\gamma_k f}(x_k), \quad \gamma_k = \frac{t}{\|\nabla f(x_{k+1})\|_2}$$



explicit normalized GD (orange)
level set through x_{k+1} (purple)
(darker = lower f)


BPM = Implicit Normalized GD

$$x_{k+1} \in \operatorname{brox}_f^t(x_k) \implies x_{k+1} = x_k - t\,\frac{\nabla f(x_{k+1})}{\|\nabla f(x_{k+1})\|_2}$$

Implicit (BPM) vs explicit

$$\hat{x}_{k+1} = x_k - t\,\frac{\nabla f(x_k)}{\|\nabla f(x_k)\|_2} \quad \text{(explicit)}$$

$$x_{k+1} = \operatorname{prox}_{\gamma_k f}(x_k), \quad \gamma_k = \frac{t}{\|\nabla f(x_{k+1})\|_2}$$



explicit normalized GD (orange)
level set through x_{k+1} (purple)
(darker = lower f)

BPM = Implicit Normalized GD


$$x_{k+1} \in \operatorname{brox}_f^t(x_k) \implies x_{k+1} = x_k - t\,\frac{\nabla f(x_{k+1})}{\|\nabla f(x_{k+1})\|_2}$$

Implicit (BPM) vs explicit

$$\hat{x}_{k+1} = x_k - t\,\frac{\nabla f(x_k)}{\|\nabla f(x_k)\|_2} \quad \text{(explicit)}$$

$$x_{k+1} = \operatorname{prox}_{\gamma_k f}(x_k), \quad \gamma_k = \frac{t}{\|\nabla f(x_{k+1})\|_2}$$

BPM: $f(x_4) = f_\star$
normalized GD: $f(\hat{x}_{10}) = 5.48$



Argmin f


explicit normalized GD (orange)
level set through x_{k+1} (purple)
(darker = lower f)

BPM = Normalized GD on the Ball Envelope

$$x_{k+1} \in \mathrm{brox}_f^t(x_k) \implies x_{k+1} = x_k - t\,\frac{\nabla N_f^t(x_k)}{\|\nabla N_f^t(x_k)\|_2}$$

BPM = Normalized GD on the Ball Envelope

$$x_{k+1} \in \mathrm{brox}_f^t(x_k) \implies x_{k+1} = x_k - t\,\frac{\nabla N_f^t(x_k)}{\|\nabla N_f^t(x_k)\|_2}$$




BPM = Normalized GD on the Ball Envelope

$$x_{k+1} \in \operatorname{brox}_f^t(x_k) \implies x_{k+1} = x_k - t\,\frac{\nabla N_f^t(x_k)}{\|\nabla N_f^t(x_k)\|_2}$$

Ball envelope

$$N_f^t(x) := \min_{\|y-x\|_2 \le t} f(y)$$

$$\{N_f^t \le c\} = \{f \le c\} + \mathcal{B}(0,t)$$



$$\operatorname{Argmin} N_f^t = \operatorname{Argmin} f + \mathcal{B}(0,t)$$

level sets of f (gray)
level sets of N_f^t (dashed)
(darker = lower f)

BPM = Normalized GD on the Ball Envelope


$$x_{k+1} \in \mathrm{brox}_f^t(x_k) \implies x_{k+1} = x_k - t\,\frac{\nabla N_f^t(x_k)}{\|\nabla N_f^t(x_k)\|_2}$$

Ball envelope

$$N_f^t(x) := \min_{\|y-x\|_2 \le t} f(y)$$

$$\{N_f^t \le c\} = \{f \le c\} + \mathcal{B}(0,t)$$

$$\mathrm{Argmin}\, N_f^t = \mathrm{Argmin}\, f + \mathcal{B}(0,t)$$



BPM = Normalized GD on the Ball Envelope

$$x_{k+1} \in \mathrm{brox}_f^t(x_k) \implies x_{k+1} = x_k - t\,\frac{\nabla N_f^t(x_k)}{\|\nabla N_f^t(x_k)\|_2}$$


Ball envelope

$$N_f^t(x) := \min_{\|y-x\|_2 \le t} f(y)$$

$$\{N_f^t \le c\} = \{f \le c\} + \mathcal{B}(0,t)$$

$$N_f^t(x_k) = f(x_{k+1}),\quad \nabla N_f^t(x_k) = \nabla f(x_{k+1})$$

$$\mathrm{Argmin}\, N_f^t = \mathrm{Argmin}\, f + \mathcal{B}(0,t)$$



BPM = Normalized GD on the Ball Envelope

$$x_{k+1} \in \mathrm{brox}_f^t(x_k) \implies x_{k+1} = x_k - t\,\frac{\nabla N_f^t(x_k)}{\|\nabla N_f^t(x_k)\|_2}$$


Ball envelope

$$N_f^t(x) := \min_{\|y-x\|_2 \le t} f(y)$$

$$\{N_f^t \le c\} = \{f \le c\} + \mathcal{B}(0,t)$$

$$N_f^t(x_k) = f(x_{k+1}),\quad \nabla N_f^t(x_k) = \nabla f(x_{k+1})$$

$$\mathrm{Argmin}\, N_f^t = \mathrm{Argmin}\, f + \mathcal{B}(0,t)$$



level sets of f (gray)
level sets of N_f^t (dashed)
(darker = lower f)

BPM = Normalized GD on the Ball Envelope

$$x_{k+1} \in \mathrm{brox}_f^t(x_k) \implies x_{k+1} = x_k - t\,\frac{\nabla N_f^t(x_k)}{\|\nabla N_f^t(x_k)\|_2}$$


Ball envelope

$$N_f^t(x) := \min_{\|y-x\|_2 \le t} f(y)$$

$$\{N_f^t \le c\} = \{f \le c\} + \mathcal{B}(0,t)$$

$$N_f^t(x_k) = f(x_{k+1}),\quad \nabla N_f^t(x_k) = \nabla f(x_{k+1})$$

$$\mathrm{Argmin}\, N_f^t = \mathrm{Argmin}\, f + \mathcal{B}(0,t)$$



level sets of f (gray)
level sets of N_f^t (dashed)
(darker = lower f)

BPM = Normalized GD on the Ball Envelope

$$x_{k+1} \in \mathrm{brox}_f^t(x_k) \implies x_{k+1} = x_k - t\,\frac{\nabla N_f^t(x_k)}{\|\nabla N_f^t(x_k)\|_2}$$


Ball envelope

$$N_f^t(x) := \min_{\|y-x\|_2 \le t} f(y)$$

$$\{N_f^t \le c\} = \{f \le c\} + \mathcal{B}(0,t)$$

$$N_f^t(x_k) = f(x_{k+1}),\quad \nabla N_f^t(x_k) = \nabla f(x_{k+1})$$

$$\mathrm{Argmin}\, N_f^t = \mathrm{Argmin}\, f + \mathcal{B}(0,t)$$



level sets of f (gray)
level sets of N_f^t (dashed)
(darker = lower f)

BPM = Normalized GD on the Ball Envelope


$$x_{k+1} \in \operatorname{brox}_f^t(x_k) \implies x_{k+1} = x_k - t\,\frac{\nabla N_f^t(x_k)}{\|\nabla N_f^t(x_k)\|_2}$$

Ball envelope

$$N_f^t(x) := \min_{\|y-x\|_2 \le t} f(y)$$

$$\{N_f^t \le c\} = \{f \le c\} + \mathcal{B}(0,t)$$

$$N_f^t(x_k) = f(x_{k+1}),\quad \nabla N_f^t(x_k) = \nabla f(x_{k+1})$$



$\operatorname{Argmin} N_f^t = \operatorname{Argmin} f + \mathcal{B}(0,t)$

level sets of f (gray)
level sets of N_f^t (dashed)
(darker = lower f)

BPM = Normalized GD on the Ball Envelope

$$x_{k+1} \in \mathrm{brox}_f^t(x_k) \implies x_{k+1} = x_k - t\,\frac{\nabla N_f^t(x_k)}{\|\nabla N_f^t(x_k)\|_2}$$


Ball envelope

$$N_f^t(x) := \min_{\|y-x\|_2 \le t} f(y)$$

$$\{N_f^t \le c\} = \{f \le c\} + \mathcal{B}(0,t)$$

$$N_f^t(x_k) = f(x_{k+1}),\quad \nabla N_f^t(x_k) = \nabla f(x_{k+1})$$

$$\mathrm{Argmin}\, N_f^t = \mathrm{Argmin}\, f + \mathcal{B}(0,t)$$



level sets of f (gray)
level sets of N_f^t (dashed)
(darker = lower f)

BPM = Normalized GD on the Ball Envelope

$$x_{k+1} \in \mathrm{brox}_f^t(x_k) \implies x_{k+1} = x_k - t\,\frac{\nabla N_f^t(x_k)}{\|\nabla N_f^t(x_k)\|_2}$$


Ball envelope

$$N_f^t(x) := \min_{\|y-x\|_2 \le t} f(y)$$

$$\{N_f^t \le c\} = \{f \le c\} + \mathcal{B}(0,t)$$

$$N_f^t(x_k) = f(x_{k+1}),\quad \nabla N_f^t(x_k) = \nabla f(x_{k+1})$$

$\mathrm{Argmin}\, N_f^t = \mathrm{Argmin}\, f + \mathcal{B}(0,t)$



level sets of f (gray)
level sets of N_f^t (dashed)
(darker = lower f)

BPM = Normalized GD on the Ball Envelope

$$x_{k+1} \in \mathrm{brox}_f^t(x_k) \implies x_{k+1} = x_k - t\,\frac{\nabla N_f^t(x_k)}{\|\nabla N_f^t(x_k)\|_2}$$

Ball envelope


$$N_f^t(x) := \min_{\|y-x\|_2 \le t} f(y)$$

$$\{N_f^t \le c\} = \{f \le c\} + \mathcal{B}(0,t)$$

$$N_f^t(x_k) = f(x_{k+1}),\quad \nabla N_f^t(x_k) = \nabla f(x_{k+1})$$

$$N_f^t(x_3) = 0 \;\Rightarrow\; x_4 \in \mathrm{Argmin}\, f$$

$$\mathrm{Argmin}\, N_f^t = \mathrm{Argmin}\, f + \mathcal{B}(0,t)$$



BPM = Implicit Regularized Newton

$$x_{k+1} = x_k - \left(\bar{H}_k + \underbrace{\lambda_k I}_{\text{regularization}}\right)^{-1} \nabla f(x_k)$$

Example: $f(x) = \frac{1}{3}\left(|\langle a, x\rangle|^3 + 8|\langle b, x\rangle|^3\right), \quad t = 0.6$



P.R.
The Design and Analysis of Newton-Type Algorithms as Approximate Broximal Point Methods
Manuscript, 2026

BPM = Implicit Regularized Newton

averaged Hessian

$$\bar{H}_k := \int_0^1 \nabla^2 f\left(x_k + \tau\left(x_{k+1} - x_k\right)\right) d\tau$$

$$x_{k+1} = x_k - \left(\bar{H}_k + \underbrace{\lambda_k I}_{\text{regularization}}\right)^{-1} \nabla f(x_k)$$

Example: $f(x) = \frac{1}{3}\left(|\langle a, x\rangle|^3 + 8|\langle b, x\rangle|^3\right), \quad t = 0.6$



P.R.
The Design and Analysis of Newton-Type Algorithms as Approximate Broximal Point Methods
Manuscript, 2026


BPM = Implicit Regularized Newton

averaged Hessian

$$\bar{H}_k := \int_0^1 \nabla^2 f\left(x_k + \tau\left(x_{k+1} - x_k\right)\right) d\tau$$

$$x_{k+1} = x_k - \left(\bar{H}_k + \underbrace{\lambda_k I}_{\text{regularization}}\right)^{-1} \nabla f(x_k)$$

Example: $f(x) = \frac{1}{3}\left(|\langle a, x\rangle|^3 + 8|\langle b, x\rangle|^3\right), \quad t = 0.6$



P.R.
The Design and Analysis of Newton-Type Algorithms as Approximate Broximal Point Methods
Manuscript, 2026

BPM = Implicit Regularized Newton

averaged Hessian


$$\bar{H}_k := \int_0^1 \nabla^2 f\left(x_k + \tau\left(x_{k+1} - x_k\right)\right) d\tau$$

$$x_{k+1} = x_k - \left(\bar{H}_k + \underbrace{\lambda_k I}_{\text{regularization}}\right)^{-1} \nabla f(x_k)$$

Example: $f(x) = \frac{1}{3}\left(|\langle a, x\rangle|^3 + 8|\langle b, x\rangle|^3\right), \quad t = 0.6$

Why? (convex, twice-differentiable objective)

optimality: $\nabla f(x_{k+1}) + \lambda_k s_k = 0, \quad \lambda_k = \|\nabla f(x_{k+1})\|_2 / t$



P.R.
The Design and Analysis of Newton-Type Algorithms as Approximate Broximal Point Methods
Manuscript, 2026

BPM = Implicit Regularized Newton

averaged Hessian

$$\bar{H}_k := \int_0^1 \nabla^2 f\left(x_k + \tau\,(x_{k+1} - x_k)\right) d\tau$$

$$x_{k+1} = x_k - \left(\bar{H}_k + \underbrace{\lambda_k I}_{\text{regularization}}\right)^{-1} \nabla f(x_k)$$


Example: $f(x) = \frac{1}{3}\left(|\langle a, x\rangle|^3 + 8|\langle b, x\rangle|^3\right), \quad t = 0.6$

Why? (convex, twice-differentiable objective)

optimality: $\nabla f(x_{k+1}) + \lambda_k s_k = 0, \quad \lambda_k = \|\nabla f(x_{k+1})\|_2 / t$

mean value: $\nabla f(x_{k+1}) - \nabla f(x_k) = \bar{H}_k s_k, \quad s_k := x_{k+1} - x_k$

$\Longrightarrow \quad (\bar{H}_k + \lambda_k I)\, s_k = -\nabla f(x_k)$ (implicit: $\bar{H}_k, \lambda_k$ depend on x_{k+1})



P.R.
The Design and Analysis of Newton-Type Algorithms as Approximate Broximal Point Methods
Manuscript, 2026

BPM = Implicit Regularized Newton

averaged Hessian

$$\bar{H}_k := \int_0^1 \nabla^2 f\left(x_k + \tau\,(x_{k+1} - x_k)\right) d\tau$$

$$x_{k+1} = x_k - \left(\bar{H}_k + \underbrace{\lambda_k I}_{\text{regularization}}\right)^{-1} \nabla f(x_k)$$


Example: $f(x) = \frac{1}{3}\left(|\langle a, x\rangle|^3 + 8|\langle b, x\rangle|^3\right)$, $t = 0.6$

Why? (convex, twice-differentiable objective)

optimality: $\nabla f(x_{k+1}) + \lambda_k s_k = 0$, $\lambda_k = \|\nabla f(x_{k+1})\|_2 / t$

mean value: $\nabla f(x_{k+1}) - \nabla f(x_k) = \bar{H}_k s_k$, $s_k := x_{k+1} - x_k$

$\Longrightarrow$ $(\bar{H}_k + \lambda_k I)\, s_k = -\nabla f(x_k)$ (implicit: $\bar{H}_k$, λ_k depend on x_{k+1})



P.R.
The Design and Analysis of Newton-Type Algorithms as Approximate Broximal Point Methods
Manuscript, 2026

BPM = Implicit Regularized Newton

averaged Hessian

$$\bar{H}_k := \int_0^1 \nabla^2 f\left(x_k + \tau\,(x_{k+1} - x_k)\right) d\tau$$

$$x_{k+1} = x_k - \left(\bar{H}_k + \underbrace{\lambda_k I}_{\text{regularization}}\right)^{-1} \nabla f(x_k)$$


Example: $f(x) = \frac{1}{3}\left(|\langle a, x\rangle|^3 + 8|\langle b, x\rangle|^3\right)$, $t = 0.6$

Why? (convex, twice-differentiable objective)

optimality: $\nabla f(x_{k+1}) + \lambda_k s_k = 0, \quad \lambda_k = \|\nabla f(x_{k+1})\|_2 / t$

mean value: $\nabla f(x_{k+1}) - \nabla f(x_k) = \bar{H}_k s_k, \quad s_k := x_{k+1} - x_k$

$\Longrightarrow \quad (\bar{H}_k + \lambda_k I)\, s_k = -\nabla f(x_k)$ (implicit: $\bar{H}_k$, λ_k depend on x_{k+1})



P.R.
The Design and Analysis of Newton-Type Algorithms as Approximate Broximal Point Methods
Manuscript, 2026

BPM = Implicit Regularized Newton

averaged Hessian

$$\bar{H}_k := \int_0^1 \nabla^2 f\left(x_k + \tau\,(x_{k+1} - x_k)\right) d\tau$$

$$x_{k+1} = x_k - \left(\bar{H}_k + \underbrace{\lambda_k I}_{\text{regularization}}\right)^{-1} \nabla f(x_k)$$


Example: $f(x) = \frac{1}{3}\left(|\langle a, x\rangle|^3 + 8|\langle b, x\rangle|^3\right)$, $t = 0.6$

Why? (convex, twice-differentiable objective)

optimality: $\nabla f(x_{k+1}) + \lambda_k s_k = 0, \quad \lambda_k = \|\nabla f(x_{k+1})\|_2 / t$

mean value: $\nabla f(x_{k+1}) - \nabla f(x_k) = \bar{H}_k s_k, \quad s_k := x_{k+1} - x_k$

$\Longrightarrow \quad (\bar{H}_k + \lambda_k I)\, s_k = -\nabla f(x_k)$ (implicit: $\bar{H}_k$, λ_k depend on x_{k+1})



P.R.
The Design and Analysis of Newton-Type Algorithms as Approximate Broximal Point Methods
Manuscript, 2026

BPM = Implicit Regularized Newton

averaged Hessian

$$\bar{H}_k := \int_0^1 \nabla^2 f\left(x_k + \tau\,(x_{k+1} - x_k)\right) d\tau$$

$$x_{k+1} = x_k - \left(\bar{H}_k + \underbrace{\lambda_k I}_{\text{regularization}}\right)^{-1} \nabla f(x_k)$$


Example: $f(x) = \frac{1}{3}\left(|\langle a, x\rangle|^3 + 8|\langle b, x\rangle|^3\right)$, $t = 0.6$

Why? (convex, twice-differentiable objective)

optimality: $\nabla f(x_{k+1}) + \lambda_k s_k = 0, \quad \lambda_k = \|\nabla f(x_{k+1})\|_2 / t$

mean value: $\nabla f(x_{k+1}) - \nabla f(x_k) = \bar{H}_k s_k, \quad s_k := x_{k+1} - x_k$

$\Longrightarrow \quad (\bar{H}_k + \lambda_k I)\, s_k = -\nabla f(x_k)$ (implicit: $\bar{H}_k$, λ_k depend on x_{k+1})



P.R.
The Design and Analysis of Newton-Type Algorithms as Approximate Broximal Point Methods
Manuscript, 2026

BPM = Implicit Regularized Newton

averaged Hessian

$$\bar{H}_k := \int_0^1 \nabla^2 f\left(x_k + \tau\,(x_{k+1} - x_k)\right) d\tau$$

$$x_{k+1} = x_k - \left(\bar{H}_k + \underbrace{\lambda_k I}_{\text{regularization}}\right)^{-1} \nabla f(x_k)$$


Example: $f(x) = \frac{1}{3}\left(|\langle a, x\rangle|^3 + 8|\langle b, x\rangle|^3\right)$, $t = 0.6$

Why? (convex, twice-differentiable objective)

optimality: $\nabla f(x_{k+1}) + \lambda_k s_k = 0, \quad \lambda_k = \|\nabla f(x_{k+1})\|_2 / t$

mean value: $\nabla f(x_{k+1}) - \nabla f(x_k) = \bar{H}_k s_k, \quad s_k := x_{k+1} - x_k$

$\Longrightarrow \quad (\bar{H}_k + \lambda_k I)\, s_k = -\nabla f(x_k)$ (implicit: $\bar{H}_k$, λ_k depend on x_{k+1})



P.R.
The Design and Analysis of Newton-Type Algorithms as Approximate Broximal Point Methods
Manuscript, 2026

BPM = Implicit Regularized Newton

averaged Hessian

$$\bar{H}_k := \int_0^1 \nabla^2 f\left(x_k + \tau\,(x_{k+1} - x_k)\right) d\tau$$

$$x_{k+1} = x_k - \left(\bar{H}_k + \underbrace{\lambda_k I}_{\text{regularization}}\right)^{-1} \nabla f(x_k)$$


Example: $f(x) = \frac{1}{3}\left(|\langle a, x\rangle|^3 + 8|\langle b, x\rangle|^3\right)$, $t = 0.6$

Why? (convex, twice-differentiable objective)

optimality: $\nabla f(x_{k+1}) + \lambda_k s_k = 0, \quad \lambda_k = \|\nabla f(x_{k+1})\|_2 / t$

mean value: $\nabla f(x_{k+1}) - \nabla f(x_k) = \bar{H}_k s_k, \quad s_k := x_{k+1} - x_k$

$\Longrightarrow \quad (\bar{H}_k + \lambda_k I)\, s_k = -\nabla f(x_k)$ (implicit: $\bar{H}_k, \lambda_k$ depend on x_{k+1})



P.R.
The Design and Analysis of Newton-Type Algorithms as Approximate Broximal Point Methods
Manuscript, 2026

BPM = Implicit Regularized Newton

averaged Hessian

$$\bar{H}_k := \int_0^1 \nabla^2 f\left(x_k + \tau\,(x_{k+1} - x_k)\right) d\tau$$

$$x_{k+1} = x_k - \left(\bar{H}_k + \underbrace{\lambda_k I}_{\text{regularization}}\right)^{-1} \nabla f(x_k)$$


Example: $f(x) = \frac{1}{3}\left(|\langle a, x\rangle|^3 + 8|\langle b, x\rangle|^3\right)$, $t = 0.6$

Why? (convex, twice-differentiable objective)

optimality: $\nabla f(x_{k+1}) + \lambda_k s_k = 0$, $\lambda_k = \|\nabla f(x_{k+1})\|_2 / t$

mean value: $\nabla f(x_{k+1}) - \nabla f(x_k) = \bar{H}_k s_k$, $s_k := x_{k+1} - x_k$

$\Longrightarrow$ $(\bar{H}_k + \lambda_k I)\, s_k = -\nabla f(x_k)$ (implicit: $\bar{H}_k$, λ_k depend on x_{k+1})



P.R.
The Design and Analysis of Newton-Type Algorithms as Approximate Broximal Point Methods
Manuscript, 2026

BPM = Implicit Regularized Newton

averaged Hessian

$$\bar{H}_k := \int_0^1 \nabla^2 f\left(x_k + \tau\,(x_{k+1} - x_k)\right) d\tau$$

$$x_{k+1} = x_k - \left(\bar{H}_k + \underbrace{\lambda_k I}_{\text{regularization}}\right)^{-1} \nabla f(x_k)$$

Example: $f(x) = \frac{1}{3}\left(|\langle a, x\rangle|^3 + 8|\langle b, x\rangle|^3\right)$, $t = 0.6$

Why? (convex, twice-differentiable objective)


optimality: $\nabla f(x_{k+1}) + \lambda_k s_k = 0, \quad \lambda_k = \|\nabla f(x_{k+1})\|_2 / t$

mean value: $\nabla f(x_{k+1}) - \nabla f(x_k) = \bar{H}_k s_k, \quad s_k := x_{k+1} - x_k$

$\Longrightarrow \quad (\bar{H}_k + \lambda_k I)\, s_k = -\nabla f(x_k)$ (implicit: $\bar{H}_k$, λ_k depend on x_{k+1})

Computable: replace $\bar{H}_k \to \nabla^2 f(x_k)$ and $\lambda_k \to \sqrt{\sigma \|\nabla f(x_k)\|_2}$

$\Rightarrow$ regularized Newton (Mishchenko, 2023): orange, $\sigma = 8$



P.R.
The Design and Analysis of Newton-Type Algorithms as Approximate Broximal Point Methods
Manuscript, 2026

Conclusions


The BPM Family Tree

BPM


$$x_{k+1} \in \underset{y}{\text{Argmin}}\ f(y)$$

$$\text{s.t.}\ \ \|y - x_k\| \leq t_k$$


The BPM Family Tree




The BPM Family Tree




The BPM Family Tree




The BPM Family Tree



The BPM Family Tree



The BPM Family Tree



The BPM Family Tree

BPM

$$x_{k+1} \in \operatorname*{Argmin}_{y} f(y) \quad \text{s.t.} \quad \|y - x_k\| \le t_k$$

- linearize f → **Gradient descent**
- linearize a random summand f_i → **SGD**
- linearize, spectral-norm ball, momentum → **Spectral SGD, Muon**
- linearize f, keep $\mathcal{X}$ exactly → **BroxGD**
- $\nabla f(x_{k+1}) \to \nabla f(x_k)$ → **Normalized GD**
- $\bar{H}_k \to \nabla^2 f(x_k)$, $\lambda_k \to \sqrt{\sigma \|\nabla f(x_k)\|_2}$ → **Regularized Newton**
- ball → penalty $\|y - x_k\|_2^2/(2\gamma)$ → **Proximal point method**

Conclusions

BPM: $x_{k+1} \in \operatorname{Argmin} \{ f(y) \ : \ \|y - x_k\| \leq t_k \}$

Conclusions

$$\text{BPM:} \quad x_{k+1} \in \operatorname{Argmin} \{ f(y) \ : \ \|y - x_k\| \leq t_k \}$$

Part 1
Convex Regime

any norm ball (ℓ_2, ℓ_1, ℓ_∞): exact solution in a few steps
LP $\rightarrow$ a few small LPs, each using only a few constraints

Conclusions

BPM: $x_{k+1} \in \operatorname{Argmin} \{ f(y) \ : \ \|y - x_k\| \le t_k \}$

Part 1 **Convex Regime**	any norm ball (ℓ_2, ℓ_1, ℓ_∞): exact solution in a few steps LP $\to$ a few small LPs, each using only a few constraints
Part 2 **Convergence Theory**	$d_{k+1}^2 \le d_k^2 - t^2$: exact solution after $\lceil d_0^2 / t^2 \rceil$ steps; no condition number any radii: $\sum_k t_k = \infty \ \Rightarrow \ f(x_k) \downarrow \inf f$

Conclusions

$$\text{BPM:} \quad x_{k+1} \in \operatorname{Argmin} \{ f(y) \ : \ \|y - x_k\| \le t_k \}$$

Part 1 **Convex Regime**	any norm ball (ℓ_2, ℓ_1, ℓ_∞): exact solution in a few steps LP $\to$ a few small LPs, each using only a few constraints
Part 2 **Convergence Theory**	$d_{k+1}^2 \le d_k^2 - t^2$: exact solution after $\lceil d_0^2 / t^2 \rceil$ steps; no condition number any radii: $\sum_k t_k = \infty \ \Rightarrow \ f(x_k) \downarrow \inf f$
Part 3 **Nonconvex Regime**	BPM jumps over non-optimal local minima where GD gets stuck brox-aligned f: Theorem 4.1 still applies $\Rightarrow$ global minimum

Conclusions

$$\text{BPM:} \quad x_{k+1} \in \operatorname{Argmin}\{f(y) \ : \ \|y - x_k\| \le t_k\}$$

Part 1 **Convex Regime**	any norm ball (ℓ_2, ℓ_1, ℓ_∞): exact solution in a few steps LP $\to$ a few small LPs, each using only a few constraints
Part 2 **Convergence Theory**	$d_{k+1}^2 \le d_k^2 - t^2$: exact solution after $\lceil d_0^2/t^2 \rceil$ steps; no condition number any radii: $\sum_k t_k = \infty \ \Rightarrow \ f(x_k) \downarrow \inf f$
Part 3 **Nonconvex Regime**	BPM jumps over non-optimal local minima where GD gets stuck brox-aligned f: Theorem 4.1 still applies $\Rightarrow$ global minimum
Part 4 **Practical Approximations**	linearize f inside the ball: GD, SGD, spectral SGD, Muon, BroxGD the ball sets the geometry: spectral $\to$ Muon, ball $\cap\, \mathcal{X} \to$ BroxGD

Conclusions

$$\text{BPM:} \quad x_{k+1} \in \operatorname{Argmin}\{f(y) \ : \ \|y - x_k\| \le t_k\}$$

Part 1 **Convex Regime**	any norm ball (ℓ_2, ℓ_1, ℓ_∞): exact solution in a few steps LP $\to$ a few small LPs, each using only a few constraints
Part 2 **Convergence Theory**	$d_{k+1}^2 \le d_k^2 - t^2$: exact solution after $\lceil d_0^2/t^2 \rceil$ steps; no condition number any radii: $\sum_k t_k = \infty \ \Rightarrow \ f(x_k) \downarrow \inf f$
Part 3 **Nonconvex Regime**	BPM jumps over non-optimal local minima where GD gets stuck brox-aligned f: Theorem 4.1 still applies $\Rightarrow$ global minimum
Part 4 **Practical Approximations**	linearize f inside the ball: GD, SGD, spectral SGD, Muon, BroxGD the ball sets the geometry: spectral $\to$ Muon, ball $\cap\, \mathcal{X} \to$ BroxGD
Part 5 **Four Formulations**	BPM = projection onto sublevel sets = implicit normalized GD = normalized GD on the ball envelope = implicit regularized Newton

Conclusions

$$\text{BPM:} \quad x_{k+1} \in \operatorname{Argmin}\{f(y) \ : \ \|y - x_k\| \le t_k\}$$

Part 1 **Convex Regime**	any norm ball (ℓ_2, ℓ_1, ℓ_∞): exact solution in a few steps LP $\to$ a few small LPs, each using only a few constraints
Part 2 **Convergence Theory**	$d_{k+1}^2 \le d_k^2 - t^2$: exact solution after $\lceil d_0^2/t^2 \rceil$ steps; no condition number any radii: $\sum_k t_k = \infty \ \Rightarrow \ f(x_k) \downarrow \inf f$
Part 3 **Nonconvex Regime**	BPM jumps over non-optimal local minima where GD gets stuck brox-aligned f: Theorem 4.1 still applies $\Rightarrow$ global minimum
Part 4 **Practical Approximations**	linearize f inside the ball: GD, SGD, spectral SGD, Muon, BroxGD the ball sets the geometry: spectral $\to$ Muon, ball $\cap\, \mathcal{X} \to$ BroxGD
Part 5 **Four Formulations**	BPM = projection onto sublevel sets = implicit normalized GD = normalized GD on the ball envelope = implicit regularized Newton

How to think about optimization: *minimize f over a ball*, step after step.

Practical methods approximate this step: a ball + an approximation of f + a radius rule.

The End