

Residues and Duality

Robin Hartshorne

May 1966

Preface

In the spring of 1963 I suggested to Grothendieck the possibility of my running a seminar at Harvard on his theory of duality for coherent sheaves — a theory which had been hinted at in his talk to the Séminaire Bourbaki in 1957 [Gro57b], and in his talk to the International Congress of Mathematicians in 1958 [Gro60b], but had never been developed systematically. He agreed, saying that he would provide an outline of the material, if I would fill in the details and write up lecture notes of the seminar. During the summer of 1963, he wrote a series of “pré-notes” [Gro63] which were to be the basis for the seminar.

I quote from the preface of the pré-notes:

Les présentes notes donnent une esquisse assez détaillée d’une théorie cohomologique de la dualité des Modules cohérents sur les préschémas. Les idées principales de la théorie m’étaient connues dès 1959, mais le manque de fondements adéquats d’Algèbre Homologique m’avait empêché d’aborder une rédaction d’ensemble. Cette lacune de fondements est sur le point d’être comblée par la thèse de Verdier, ce qui rend en principe possible un exposé satisfaisant. Il est d’ailleurs apparu depuis qu’il existe des théories cohomologiques de dualité formellement très analogues à celle développée ici dans toutes sortes d’autres contextes: faisceaux cohérents sur les espaces analytiques, faisceaux abéliens sur les espaces topologiques (Verdier), modules galoisiens (Verdier, Tate), faisceaux de torsion sur les schémas munis de leur topologie étale, corps de classe en tous genres... Cela me semble une raison assez sérieuse pour se familiariser avec le yoga général de la dualité dans un cas type, comme la théorie cohomologique des résidus.

La théorie consiste pour l’essentiel dans des questions de variance: construction d’un foncteur $f^!$ et d’un homomorphisme-trace $\underline{R}f_*f^! \rightarrow \text{id}$. La construction donnée ici est compliquée et indirecte et n’est pas valable sous des conditions aussi générales qu’on est en droit de s’y attendre.

Il faudra sans doute une idée nouvelle pour apporter des simplifications substantielles.

The seminar took place in the fall and winter of 1963–64, with the assistance of David Mumford, John Tate, Stephen Lichtenbaum, John Fogarty, and others, and gave rise to a series of six exposés which were circulated to a limited audience under the title “Séminaire Hartshorne”. The present notes are a revised, expanded, and completed version of the previous notes.

I would like to take this opportunity to thank all those people who have helped in the course of this work, and in particular A. Grothendieck, who gave continual support and encouragement throughout the whole project.

R.H.
Cambridge, May 1966

Contents

1	Introduction	1
2	The Derived Category	11
2.1	Introduction	11
2.2	Triangulated Categories	11
2.2.1	Definition	11
2.3	$K(\mathcal{A})$ is triangulated	14
2.4	Localization of categories	16
2.5	Quasi-isomorphisms and the derived category	20
2.6	Derived functors	27
2.7	Examples. Ext and $\underline{\text{R}}\text{Hom}$	33
2.8	Way-out functors and Isomorphisms	37
3	Application to Preschemes	47
3.1	Categories of sheaves	47
3.2	The derived functors of f_* and Γ	47
3.3	The derived functor of $\mathcal{H}om$	47
3.4	The derived functors of $\otimes$ and f^*	47
3.5	Relations among the derived functors	47
3.6	Compatibilities among the relations of §3.5	47
3.7	Injective sheaves on a locally noetherian prescheme	47
4	Duality for Projective Morphisms	49
4.1	Differentials	50
4.2	$f^\#$ for a smooth morphism f	50
4.3	Recall of the explicit calculations	50
4.4	The trace map for projective space	50
4.5	The duality theorem for projective space	50
4.6	Duality for a finite morphism	50

4.7	The fundamental local isomorphism	50
4.8	$f^!$ for embeddable morphisms	50
4.9	The residue symbol	50
4.10	Trace for projective morphisms	50
4.11	Duality for projective morphisms	50
5	Local Cohomology	51
5.1	Local cohomology groups, sheaves, and complexes	51
5.2	Depth and the Cousin complex	51
5.3	Generalization to complexes	51
6	Dualizing Complexes and Local Duality	53
6.1	Introduction	54
6.2	Example: duality for abelian groups	54
6.3	Dualizing complexes	54
6.4	Uniqueness of the dualizing complex	54
6.5	Local cohomology on a prescheme	54
6.6	Dualizing functors on a local noetherian ring	54
6.7	Local duality	54
6.8	Application to dualizing complexes	54
6.9	Pointwise dualizing complexes and $f^\#$	54
6.10	Gorenstein preschemes	54
6.11	Existence of dualizing complexes	54
7	Residual Complexes	55
7.1	Introduction	55
7.2	Functorial properties	55
7.3	$f^!$ for residual complexes	55
7.4	Trace for residual complexes	55
7.5	Behavior with respect to certain base changes	55
8	The Duality Theorem	57
8.1	Curves over an Artin ring	57
8.2	The residue theorem	57
8.3	The duality theorem for proper morphisms	57
8.4	Smooth morphisms	57
	Index of Definitions	59

CONTENTS

v

Index of Definitions	59
Index of Notations	61
Index of Notations	63
Cohomologie à Support Propre, et Construction du Foncteur $f^!$	73
Errata	74

Chapter 1

Introduction

The way up and the way down are one and the same.

–Heraclitus

The main purpose of these notes is to prove a duality theorem for cohomology of quasi-coherent sheaves, with respect to a proper morphism of locally noetherian preschemes. Various such theorems are already known. Typical is the duality theorem for a non-singular complete curve X over an algebraically closed field k , which says that

$$h^0(D) = h^1(K - D),$$

where D is a divisor, K is the canonical divisor, and

$$h^i(D) = \dim_k H^i(X, \mathcal{O}_X(D))$$

for any i , and any divisor D . (See e.g. [Ch. II; Ser59] for a proof.)

Various attempts were made to generalize this theorem to varieties of higher dimension, and as Zariski points out in his report [Zar56], his generalization of a lemma of Enriques–Severi [Zar52] is equivalent to the statement that for a normal projective variety X of dimension n over k , for any divisor D ,

$$h^0(D) = h^n(K - D).$$

This is also equivalent to a theorem of Serre [Ser55a] §76, Thm. 4 on the vanishing of the cohomology group $H^1(X, \mathcal{O}_X(-m))$ for m large and $\mathcal{L}$ locally free. Using

a related theorem [Ser55a] §75, Thm. 3, Zariski shows how one can deduce on a non-singular projective variety the formula

$$h^i(D) = h^{n-i}(K - D)$$

for $0 \leq i \leq n$. In terms of sheaves, this result corresponds to the fact that the k -vector spaces

$$H^i(X, \mathcal{F}) \quad \text{and} \quad H^{n-i}(X, \mathcal{F}^\vee \otimes \omega)$$

are dual to each other, where $\mathcal{F}$ is a locally free sheaf, $\mathcal{F}^\vee$ is the dual sheaf $\underline{\text{Hom}}(\mathcal{F}, \mathcal{O}_X)$, and $\omega = \Omega_{X/k}^n$ is the sheaf of n -differentials on X . Serre [Ser55b] gives a proof of this same theorem by analytic methods for a compact complex analytic manifold X .

Grothendieck [Gro57b] gave some generalizations of these theorems for non-singular projective varieties and then in [Gro60b] announced the general theorem for schemes proper over a field, with arbitrary singularities, which is the subject of the present lecture notes.

To motivate the statement of our main theorem, let us consider the case of projective space $X = \mathbb{P}_k^n$ over an algebraically closed field k . Then there is a canonical isomorphism

$$H^n(X, \omega) \cong k \tag{1.1}$$

where $\omega = \Omega_{X/k}^n$ is the sheaf of n -differentials. Combining this with the Yoneda pairing

$$H^i(X, \mathcal{F}) \times \text{Ext}_X^{n-i}(\mathcal{F}, \omega) \rightarrow H^n(X, \omega) \tag{1.2}$$

we obtain a pairing

$$H^i(X, \mathcal{F}) \times \text{Ext}_X^{n-i}(\mathcal{F}, \omega) \rightarrow k \tag{1.3}$$

which one shows easily to be a perfect pairing [SGA 62; expose 12]. This generalizes the statements above, because for a locally free sheaf $\mathcal{F}$,

$$\text{Ext}^{n-i}_X(\mathcal{F}, \omega) = \text{Ext}^{n-i}_X(\mathcal{O}_X, \mathcal{F}^\vee \otimes \omega) = H^{n-i}(X, \mathcal{F}^\vee \otimes \omega).$$

Another way of looking at our duality pairing is as an isomorphism

$$\text{Ext}_X^{n-i}(\mathcal{F}, \omega) \cong \text{Hom}_k(H^i(X, \mathcal{F}), k). \tag{1.4}$$

Since everything is linear over k , we may introduce a k -vector space G , and have an isomorphism

$$\text{Ext}_X^{n-i}(\mathcal{F}, G \otimes_k \omega) \rightarrow \text{Hom}_k(H^i(X, \mathcal{F}), G). \tag{1.5}$$

Before proceeding further, we must introduce the derived category. It will be discussed in detail in Chapter I, but for the moment it will be sufficient to know

the following: For each abelian category $\mathcal{A}$, there is a category $D(\mathcal{A})$, called the derived category of $\mathcal{A}$, whose objects are complexes of objects of $\mathcal{A}$. If $F: \mathcal{A} \rightarrow \mathcal{B}$ is an additive functor from one abelian category to another, then under reasonable conditions there is a right derived functor $\underline{\underline{R}}F: D(\mathcal{A}) \rightarrow D(\mathcal{B})$ with the property that for any $X \in \text{Ob } \mathcal{A}$, if X denotes also the complex which is X in degree zero, and zero elsewhere, then

$$H^i(\underline{\underline{R}}F(X)) = R^iF(X),$$

where R^iF is the ordinary i^{th} right derived functor of F . Finally, if $F: \mathcal{A} \rightarrow \mathcal{B}$ and $G: \mathcal{B} \rightarrow \mathcal{C}$ are two functors then

$$\underline{\underline{R}}(G \circ F) = \underline{\underline{R}}G \circ \underline{\underline{R}}F.$$

This replaces the old-fashioned spectral sequence of a composite functor.

Now we can jazz up our duality for projective space as follows. We replace k by a prescheme Y , so that $X = \mathbb{P}_Y^n$. We consider the derived categories $D(X)$ and $D(Y)$ of the categories of $\mathcal{O}_X$ -modules and $\mathcal{O}_Y$ -modules, respectively. Then cohomology H^i becomes $\underline{\underline{R}}f_*$, the derived functor of the direct image functor f_* , where $f: X \rightarrow Y$ is the projection. Ext becomes the derived functor $\underline{\underline{R}}\text{Hom}$ of Hom . We define

$$f^!(\mathcal{G}) = f^*(\mathcal{G}) \otimes \omega$$

for $\mathcal{G} \in D(Y)$, and we replace $\mathcal{F}$ by a complex of sheaves $\mathcal{F} \in D(X)$. Then the isomorphism (1.1) gives us an isomorphism

$$\underline{\underline{R}}f_* f^! \mathcal{G} \xrightarrow{\sim} \mathcal{G} \tag{1.6}$$

which we call the trace map.

The Yoneda pairing reappears as a natural map

$$\underline{\underline{R}}\text{Hom}_X(\mathcal{F}, f^! \mathcal{G}) \rightarrow \underline{\underline{R}}\text{Hom}_Y(\underline{\underline{R}}f_* \mathcal{F}, \underline{\underline{R}}f_* f^! \mathcal{G}), \tag{1.7}$$

which, composed with the trace map (1.6) gives us the duality morphism

$$\underline{\underline{R}}\text{Hom}_X(\mathcal{F}, f^! \mathcal{G}) \rightarrow \underline{\underline{R}}\text{Hom}_Y(\underline{\underline{R}}f_* \mathcal{F}, \mathcal{G}) \tag{1.8}$$

which generalizes (1.5). This is easily proved to be an isomorphism ([III, 5.1] below) under suitable hypotheses on $Y, \mathcal{F}, \mathcal{G}$. In fact, the proof is nothing but “general nonsense” once one has the isomorphism (1.4).

Having examined the case of projective space, we can state the following ideal theorem, which is the primum mobile of these notes, although it may never appear explicitly in this form.

Ideal Theorem.

- (a) For every morphism $f: X \rightarrow Y$ of finite type of preschemes, there is a functor $f^!$ such that

- 1) if $g: Y \rightarrow Z$ is a second morphism of finite type, then

$$(gf)^! = f^! \circ g^!;$$

- 2) if f is a smooth morphism, then

$$f^!(\mathcal{G}) = f^*(\mathcal{G}) \otimes \omega,$$

where $\omega = \Omega_{X/Y}^n$ is the sheaf of highest order differentials;

- 3) if f is a finite morphism, then

$$f^!(\mathcal{G}) = \underline{\text{Hom}}_{\mathcal{O}_Y}(f_*\mathcal{O}_X, \mathcal{G})^\sim.$$

- (b) For every proper morphism $f: X \rightarrow Y$ of preschemes, there is a trace morphism

$$\text{Tr}_f: \underline{\text{R}}f_*f^! \rightarrow \text{id}$$

of functors from $\text{D}(Y)$ to $\text{D}(Y)$ such that

- 1) if $g: Y \rightarrow Z$ is a second proper morphism, then $\text{Tr}_{g \circ f} = \text{Tr}_g \circ \text{Tr}_f$.
 2) if $X = \mathbb{P}_Y^n$, then Tr_f is the map deduced from the canonical isomorphism $R^n f_*(\omega) \cong \mathcal{O}_Y$.
 3) if f is a finite morphism, then Tr_f is obtained from the natural map “evaluation at one”

$$\underline{\text{Hom}}_{\mathcal{O}_Y}(f_*\mathcal{O}_X, G) \rightarrow G.$$

- (c) If $f: X \rightarrow Y$ is a proper morphism, then the duality morphism

$$\theta_f: \underline{\text{R}}\text{Hom}_X(F, f^!G) \rightarrow \underline{\text{R}}\text{Hom}_Y(\underline{\text{R}}f_*F, G)$$

obtained by composing the natural map (7) above with Tr_f , is an isomorphism for $F \in \text{D}(X)$ and $G \in \text{D}(Y)$.

It should be noted that we have deliberately left certain technical details out of the above statement, for ease of reading. Thus it seems reasonable to make these statements only for complexes of quasi-coherent sheaves, or rather for complexes of arbitrary sheaves whose cohomology sheaves are quasi-coherent. (We denote this category by $D_{qc}(Y)$ or $D_{qc}(X)$.) In fact, we give an example in the case of a finite morphism to show that the duality theorem (c) fails if G is not quasi-coherent (see example following [III, 6.7]). Secondly one must expect certain boundedness conditions on the complexes involved, namely F should be bounded above (we write $F \in D_{qc}^-(X)$) and G should be bounded below ($G \in D_{qc}^+(Y)$) for the duality theorem. Finally, questions of variance will be very important in the proof, and so the equals signs in (a) and (b) must be taken with a grain of salt. We must preserve very carefully the distinction between “equals” and “is canonically isomorphic to”. Hence we will have more precise statements below.

As to proving the ideal theorem, we succeed only partially. Certainly the conditions under which we can prove it will suffice for most applications, but it is unsatisfying to have restrictive hypotheses which are apparently not essential to the truth of the theorem. I mention four sets of hypotheses under which the theorem can be proved.

- (i) For the category of noetherian preschemes of finite Krull dimension, and morphisms $f: X \rightarrow Y$ which can be factored through a suitable projective space $\mathbb{P}_Y^N$, we have (a1)–(a3) for $D_{qc}^+(Y)$ [I, 8.7], (b1)–(b3) for $D_{qc}^+(Y)$ [I, 10.5], and (c) for $F \in D_{qc}^-(X)$ and $G \in D_{qc}^+(Y)$ [III, 11.1].
- (ii) For the category of noetherian preschemes which admit dualizing complexes (see [V, §2]; this implies in particular that the preschemes have finite Krull dimension) and morphisms whose fibres are of bounded dimension, we have (a1)–(a3) and (b1)–(b3) for $D_c^+(Y)$, and (c) for $F \in D_{qc}^-(X)$, $G \in D_c^+(Y)$ [VII, 3.4]. Here the subscript c denotes complexes with coherent cohomology sheaves.
- (iii) For the category of noetherian preschemes of finite Krull dimension and smooth morphisms, we have (a1)–(a3) for $D_{qc}^+(Y)$, (b1)–(b3) for $G \in D_{qc}^b(Y)$, and (c) for $F \in D_{qc}^-(X)$ and $G \in D_{qc}^b(Y)$ [VII, 4.3]. Here the exponent “ b ” denotes complexes which are bounded in both directions, i.e., finite.
- (iv) Recently P. Deligne has shown (unpublished) that for the category of noetherian preschemes, one can construct $f^!$ and Tr_f satisfying (a), (b), and (c), working with $F \in D(\text{Qco}(X))$ and $G \in D^+(\text{Qco}(Y))$, the derived categories of the categories of quasi-coherent sheaves on X and Y , respectively.

The principal difficulty has been the lack of a suitable construction of the functor $f^!$: to define it locally, and then glue. Our procedure is to define $f^!$ for a finite or a smooth morphism, where we have it given to us by (a2) and (a3) [III, §6, §11]. Thus by composition we can obtain it for any morphism which can be factored into a finite morphism followed by a smooth morphism [III, §8]. However, the derived category is not a local object, so we cannot glue these local determinations of $f^!$ to obtain a global one. We resort to a clumsy, round-about procedure of defining $f^!$ for a special class of complexes, called residual complexes [VI, §1], and then pulling ourselves up by our bootstraps to get it for arbitrary complexes [VII, §3]. But our result has the unpleasant hypotheses of (ii) above.

Deligne's construction of $f^!$ is entirely different, and proceeds essentially by representing the functor (for $F \in D(X)$)

$$F \longmapsto \underline{\underline{R}}\mathrm{Hom}_Y(\underline{\underline{R}}f_*F, G).$$

This approach has the advantage of giving the duality theorem immediately. He also has a method for calculating $f^!G$ locally on X . However, it is not immediately clear from his construction that the properties (a2), (a3) and (b2), (b3) hold. In fact, their proof will probably require some knowledge of the duality theorem as we have proved it here. At least one can hope that some combination of the two approaches will provide substantial simplifications of the theory at a later date.

A second difficulty, which lurks on the fringes of these notes, is that the derived category seems to be a little too big when it comes to unbounded complexes. For example, one can have two unequal morphisms $f, g: X \rightarrow Y$ in $D^+(\mathcal{A})$ (where $\mathcal{A}$ is an abelian category) such that their restrictions to each truncation of the complex X to a bounded complex are equal. This gives rise to some trouble with unbounded complexes (see the problem after [II, 5.7], the boundedness hypotheses in [IV, 3.1] and [VII, 4.3a], and the remark following [VI, 1.1c]). Perhaps one will have to replace D^+ by the categories $\mathrm{ind} D^b$ and $\mathrm{pro} D^b$, which however may not be triangulated categories.

Thirdly, some discussion of our noetherian hypotheses is in order. In the present state of the theory, the noetherian hypotheses are well entrenched. We have used them in [II, §7] for the structure of the injective objects in the category of sheaves on a locally noetherian prescheme, and its consequence that $D^+(\mathrm{Qco}(X)) \rightarrow D_{\mathrm{qc}}^+(X)$ is an equivalence of categories. We have used them in the construction of the trace map for projective space [III, §4], in the construction of $f^!$ for a finite morphism [III, §6], in the whole theory of dualizing complexes [V, §2], in the definition of residual complexes [VI, §1], and so forth. Our finite Krull dimension hypotheses are often needed only to make possible the definition of $\mathbb{R}f_*$ and $f^!$ for unbounded complexes — a problem

which will disappear when the second difficulty above is solved, and the relation between $D(\mathbf{Qco}(X))$ and $D_{qc}(X)$ is better understood. It is certainly clear that our methods of proof rely heavily on noetherian hypotheses. I expect, however, that once a suitable statement is obtained, e.g., in case (iii) above, one could expect to prove the theorem without noetherian hypotheses, by reducing to the noetherian case. More satisfactory, of course, would be a treatment where noetherian hypotheses on the base were eliminated from the proofs as well as from the statements. When that is achieved, it will be reasonable to state the duality theorem for a proper morphism of ringed tops, whose fibres are noetherian schemes... At the present, however, this must remain a dream of things to come.

Now we will give the reader a brief description of the organization of these notes.

Chapter I gives the language of derived categories, which is used continually in the sequel. Sections 1–5, containing the definition of derived categories and derived functors, are essential, while sections 6 and 7 are refinements which are needed in proofs later on. This chapter is a self-contained treatment of the subject, and makes no reference to algebraic geometry, so can be used independently as an introduction to the notion of derived category. Sections 1–6 (except for the notion of localizing subcategory and the corresponding categories $K_{A'}(\mathcal{A})$, $D_{A'}(\mathcal{A})$, etc.) are taken almost without change from notes of Verdier, and should appear in his thesis [Ver63b].

Chapter II is a fairly systematic treatment of the applications of the language of derived categories to the category of sheaves on a prescheme. We consider the functors Γ , f_* , Hom , $\underline{\text{Hom}}$, $\otimes$, f^* , their derived functors, and relations between these derived functors, such as associativity formulae. There is really no new mathematics involved here, since it is merely a translation into a new language of known results. However, some care is taken to see what hypotheses are needed. Only section 7 is notable new material. Here we give the structure of injective $\mathcal{O}_X$ -modules on a locally noetherian prescheme X , showing that they are all direct sums of certain indecomposable injectives $J(x, x')$, for pairs of points x specializing to x' of X . This extends results of Matlis [Mat58] and Gabriel [Gab62] for the case of quasi-coherent sheaves.

Chapter III contains everything we can say about duality for projective morphisms, and if the reader cares only for them, he may stop at the end of this chapter. However, it should be noted that the following chapters, besides giving us the tools for the proof of the general duality theorem in Chapter VII, also give much new insight into the nature of duality for a projective morphism.

There are so many situations in which we have a functor $f^!$ like the one mentioned in the ideal theorem above, that we use different notations for them. Thus we have

$f^\#$ for a smooth morphism in section 2, $f^\flat$ for a finite morphism in section 6, and $f^!$ for an embeddable morphism in section 8. Later we will also have f^y , f^z [VI, §2] and f^Δ [VI, §3] for residual complexes.

In sections 3, 4, and 5 we recall the explicit calculations of cohomology for projective space, and give the old duality for projective space in the new language of derived categories. Sections 6 and 7 give the corresponding formalism for finite morphisms, and its relation to the case of smooth morphisms, so that in sections 8, 10, and 11 we can prove the ideal duality theorem for projective morphisms. Section 9 gives the formalism of a residue symbol which generalizes the classical residue of a differential on a curve. Even over the complex numbers, this important concept of residue for varieties of dimension greater than one was not known before.

In Chapter IV we study “local cohomology”, or cohomology with restricted supports, of abelian sheaves on a locally noetherian topological space, generalizing results of [LC, §16]. In particular, we discuss various cohomological properties which a sheaf or complex of sheaves may have with respect to certain families of supports. This gives rise to the notions of depth, Cousin complex, Cohen–Macaulay complex, and Gorenstein complex. The results of this chapter are independent of all other chapters of these notes, and so may be of use elsewhere, although their only application for the moment is to the theory of duality on preschemes.

Chapter V discusses dualizing complexes (read section 0 to find out what one is) and gives a duality theorem for modules over a local ring, generalizing results of [LC, §16]. In particular, the dualizing functor

$$\underline{D} = \underline{\mathbf{R}}\underline{\mathbf{Hom}}(\bullet, R^\bullet)$$

treated here will be useful for our bootstrap operation (construction of $f^!$) later, because it interchanges $\otimes$ and $\underline{\mathbf{Hom}}$, $f^!$ and $\underline{\mathbf{L}}f_*$, and commutes with $\underline{\mathbf{R}}f_*$ (duality theorem).

In Chapter VI we prepare for the final duality theorem by giving the construction of $f^!$ and Tr_f for residual complexes. This is accomplished by a delicate glueing procedure which is the most difficult part of the theory, so we have given it in some detail. Perhaps some day this type of construction will be done more elegantly using the language of fibred categories and results of Giraud’s thesis [Gir60].

Chapter VII contains two main results. The first is the residue theorem, which generalizes the classical theorem that the sum of the residues of a differential on a curve is zero, and which is proved by reduction to that case. The second is the proof of the duality theorem for a proper morphism, which, now that all the functorial machinery has been set up, is little more than putting together the pieces of a jigsaw puzzle.

It remains to give the reader some perspective by listing some topics which have a logical place in these notes, but which are not here.

1. The cohomology class associated to a cycle. Let X be proper and smooth of dimension n over a field k . In [8, §4] it is shown how to associate to each non-singular subvariety Y of codimension p of X , a cohomology class

$$P_X(Y) \in H^p(X, \Omega_{X/k}^p).$$

Now this can be done for an arbitrary subvariety of X , using the remarks in [9]. One defines

$$\omega_Y = H^{-n+p}(g^!k),$$

where $g: Y \rightarrow k$ is the projection. Then there is a canonical element

$$\eta \in \text{Ext}_{\mathcal{O}_Y}^{-n+p}(\omega_Y, g^!k).$$

One defines a natural map

$$\Omega_{Y/k}^{n-p} \rightarrow \omega_Y,$$

which, together with η and the construction of [8, §4], gives the cohomology class $P_X(Y)$. One proves the fundamental theorem that formation of the cohomology class associated to a cycle takes an intersection of cycles into the cup-product of their cohomology classes.

2. The theory of Poincaré duality and the Gysin homomorphism can be developed as in [8, §4].

3. A Lefschetz–Verdier fixed point formula for coherent sheaves on a scheme proper over a field, to generalize [21, Theorem 2]. In particular, the determination of the local contribution at a non-simple fixed point presents an interesting topic for future investigation.

4. The still lacking theory of duality for complexes with differential operators as boundary operator, its ties with ordinary singular homology theory and with vector bundles with integrable connections, as suggested in [22].

A remark on references: Theorem 5.1 of Chapter III is referred to as “Theorem 5.1” in Chapter III, and as [III, 5.1] elsewhere. References to the bibliography at the end are given by square brackets with an arabic numeral or some capital letters, e.g., [14, (31.I)] or [EGA III, 2.1.12].

Chapter 2

The Derived Category

2.1 Introduction

Let $\mathcal{A}$ and $\mathcal{B}$ be abelian categories, and let $F: \mathcal{A} \rightarrow \mathcal{B}$ be a functor. Our purpose is to define, functorially for each $X \in \text{Ob } \mathcal{A}$, a complex $\underline{\underline{R}}F(X)$ whose cohomology groups are the right derived functors of F acting on X , namely $R^i F(X)$. In general, we will define $\underline{\underline{R}}F(X^\bullet)$ for any complex $X^\bullet$ of objects of $\mathcal{A}$.

More precisely, $\underline{\underline{R}}F$ will be a functor from the derived category $D(\mathcal{A})$ of $\mathcal{A}$ to the derived category $D(\mathcal{B})$ of $\mathcal{B}$.

The derived category $D(\mathcal{A})$ is obtained as follows: one first considers the category $K(\mathcal{A})$, whose objects are complexes of elements of $\mathcal{A}$, and whose morphisms are homotopy equivalence classes of morphisms of complexes. The category $D(\mathcal{A})$ is obtained by “localizing” $K(\mathcal{A})$ so that every morphism in $K(\mathcal{A})$ which induces an isomorphism on cohomology becomes an isomorphism in $D(\mathcal{A})$. This process of localization will be explained in general below.

Although $\mathcal{A}$ and $\mathcal{B}$ above are assumed to be abelian categories, the categories $K(\mathcal{A})$, $D(\mathcal{A})$, etc., are in general not abelian. However, they can be given a structure which carries enough information for our purposes, namely a structure of triangulated category. Thus we will be led to study triangulated categories.

2.2 Triangulated Categories

2.2.1 Definition

A triangulated category is an additive category $\mathcal{C}$, together with

- a) an automorphism $T: \mathcal{C} \rightarrow \mathcal{C}$ called the *translation functor*, and
- b) a collection of sextuples (X, Y, Z, u, v, w) , called the *triangles* of $\mathcal{C}$, where in each sextuple $X, Y, Z \in \text{Ob } \mathcal{C}$ and

$$u: X \rightarrow Y, \quad v: Y \rightarrow Z, \quad w: Z \rightarrow T(X)$$

are morphisms. A triangle is usually written

$$\begin{array}{ccc} X & \xrightarrow{u} & Y \\ & \swarrow w & \downarrow v \\ & & Z \end{array}$$

A morphism of triangles


$$(X, Y, Z, u, v, w) \rightarrow (X', Y', Z', u', v', w')$$

is a commutative diagram

$$\begin{array}{ccccccc} X & \xrightarrow{u} & Y & \xrightarrow{v} & Z & \xrightarrow{w} & T(X) \\ f \downarrow & & g \downarrow & & h \downarrow & & T(f) \downarrow \\ X' & \xrightarrow{u'} & Y' & \xrightarrow{v'} & Z' & \xrightarrow{w'} & T(X') \end{array}$$

This data is subject to the following axioms:

- (TR1)** (a) Every sextuple isomorphic to a triangle is a triangle.
 (b) Every morphism $u: X \rightarrow Y$ can be embedded in a triangle.
 (c) The sextuple $(X, X, 0, \text{id}_X, 0, 0)$ is a triangle.
- (TR2)** (X, Y, Z, u, v, w) is a triangle if and only if $(Y, Z, T(X), v, w, -T(u))$ is a triangle.
- (TR3)** Given two triangles (X, Y, Z, u, v, w) and (X', Y', Z', u', v', w') , and morphisms $f: X \rightarrow X'$, $g: Y \rightarrow Y'$ such that $u'f = gu$, there exists a morphism $h: Z \rightarrow Z'$ (not necessarily unique) so that (f, g, h) is a morphism of triangles.
- (TR4)** (The octahedral axiom.)



Suppose given triangles

$$\begin{aligned} (X, Y, Z', u, j, \cdot), \\ (Y, Z, X', v, \cdot, i), \\ (X, Z, Y', vu, \cdot, \cdot) \end{aligned}$$

Then there exist morphisms $f: Z' \rightarrow Y'$ and $g: Y' \rightarrow X'$ such that

$$(Z', Y', X', f, g, T(j)i)$$

is a triangle, and the two other faces of the octahedron with f, g as edges, are commutative diagrams.

Definition 1. An additive functor $F: \mathcal{C} \rightarrow \mathcal{C}'$ from one triangulated category to another is called a (covariant) ∂ -functor if it commutes with the translation functor and takes triangles into triangles. A contravariant ∂ -functor takes triangles into triangles with the arrows reversed and sends the translation functor into its inverse.

Definition 2. An additive functor $H: \mathcal{C} \rightarrow \mathcal{A}$ from a triangulated category to an abelian category is called a covariant cohomological functor if whenever (X, Y, Z, u, v, w) is a triangle, the long sequence

$$\cdots \rightarrow H(T^i X) \rightarrow H(T^i Y) \rightarrow H(T^i Z) \rightarrow H(T^{i+1} X) \rightarrow \cdots$$

is exact (the morphisms being $H(T^i u)$, etc.). If H is a cohomological functor, we often write $H^i(X)$ for $H(T^i X)$, $i \in \mathbb{Z}$. One defines a contravariant cohomological functor by reversing the arrows.

Proposition 1. a) The composition of any two consecutive morphisms in a triangle is zero.

b) If $\mathcal{C}$ is a triangulated category and M an object of $\mathcal{C}$, then $\text{Hom}_{\mathcal{C}}(M, \cdot)$ and $\text{Hom}_{\mathcal{C}}(\cdot, M)$ are cohomological functors on $\mathcal{C}$.

c) If in the situation of (TR3) f and g are isomorphisms, then h is also an isomorphism.

Proof. a) Let (X, Y, Z, u, v, w) be a triangle. By (TR2) it is sufficient to show that $vu = 0$. Also by (TR2), $(Y, Z, T(X), v, w, -T(u))$ is a triangle. By (TR1), $(Z, Z, 0, \text{id}_Z, 0, 0)$ is a triangle. We apply (TR3) to the maps $v: Y \rightarrow Z$ and $\text{id}_Z: Z \rightarrow Z$, and conclude that there is a map $h: T(X) \rightarrow 0$ giving a morphism of triangles. It follows that $T(v)(-T(u)) = 0$, or, since T is an automorphism, $vu = 0$.

b) Let $M \in \text{Ob}(\mathcal{C})$, and let (X, Y, Z, u, v, w) be a triangle. To show $\text{Hom}_{\mathcal{C}}(M, \cdot)$ is a cohomological functor, it suffices by (TR2) to show the sequence

$$\text{Hom}_{\mathcal{C}}(M, X) \rightarrow \text{Hom}_{\mathcal{C}}(M, Y) \rightarrow \text{Hom}_{\mathcal{C}}(M, Z)$$

is exact. By a) we know the composition is zero. Now suppose given $g \in \text{Hom}_{\mathcal{C}}(M, Y)$ such that $vg \in \text{Hom}_{\mathcal{C}}(M, Z)$ is zero. We apply (TR3) to the triangles $(M, M, 0, \text{id}_M, 0, 0)$ and (X, Y, Z, u, v, w) and the map $g: M \rightarrow Y$ to conclude there exists $f: M \rightarrow X$ such that $uf = g$.

A similar proof shows that $\text{Hom}_{\mathcal{C}}(\cdot, M)$ is a contravariant cohomological functor.

c) In the situation of (TR3) suppose that f and g are isomorphisms. We apply the cohomological functor $\text{Hom}_{\mathcal{C}}(Z', \cdot)$ to the whole situation, and obtain an exact commutative diagram

$$\begin{array}{ccccccccc} \text{Hom}(Z', X) & \longrightarrow & \text{Hom}(Z', Y) & \longrightarrow & \text{Hom}(Z', Z) & \longrightarrow & \text{Hom}(Z', T(X)) & \longrightarrow & \text{Hom}(Z', T(Y)) \\ f_* \downarrow & & g_* \downarrow & & h_* \downarrow & & T(f)_* \downarrow & & T(g)_* \downarrow \\ \text{Hom}(Z', X') & \longrightarrow & \text{Hom}(Z', Y') & \longrightarrow & \text{Hom}(Z', Z') & \longrightarrow & \text{Hom}(Z', T(X')) & \longrightarrow & \text{Hom}(Z', T(Y')) \end{array}$$

where $f_* = \text{Hom}(Z', f)$ etc. Now since f and g are isomorphisms in $\mathcal{C}$, it follows that f_* , g_* , $T(f)_*$, and $T(g)_*$ are isomorphisms of abelian groups. Hence by the five-lemma, h_* is an isomorphism. We conclude that there exists a $\varphi \in \text{Hom}_{\mathcal{C}}(Z', Z)$ such that $h_*(\varphi) = h\varphi = \text{id}_{Z'} \in \text{Hom}(Z', Z')$.

Similarly using the cohomological functor $\text{Hom}_{\mathcal{C}}(\cdot, Z)$ we find there is a $\psi \in \text{Hom}(Z', Z)$ such that $\psi h = \text{id}_{Z'}$. It follows that $\varphi = \psi$ and h is an isomorphism. $\square$

2.3 $K(\mathcal{A})$ is triangulated

Let $\mathcal{A}$ be an abelian category. A *complex* of objects of $\mathcal{A}$ is a collection $X^\bullet = (X^n)_{n \in \mathbb{Z}}$ of objects of $\mathcal{A}$ together with maps $d^n: X^n \rightarrow X^{n+1}$ such that $d^{n+1}d^n = 0$ for all $n \in \mathbb{Z}$.

A morphism f of complexes $X^\bullet \rightarrow Y^\bullet$ is a collection of maps $f^n: X^n \rightarrow Y^n$ which commute with differentials:

$$f^{n+1}d_X^n = d_Y^n f^n$$

for all n .

Two maps $f, g: X^\bullet \rightarrow Y^\bullet$ are *homotopic* if there exists a collection of maps $k = (k^n)$, $k^n: X^n \rightarrow X^{n-1}$ (not necessarily commuting with d) such that

$$f^n - g^n = d_Y^{n-1}k^n + k^{n+1}d_X^n$$

for all n . Homotopy is an equivalence relation, and compositions of homotopic maps are homotopic.

We define $K(\mathcal{A})$ to be the category whose objects are complexes of objects of $\mathcal{A}$, and whose morphisms are homotopy equivalence classes of morphisms of complexes. A complex $X^\bullet$ is *bounded below* if $X^n = 0$ for $n \ll 0$. We denote by $K^+(\mathcal{A})$ the full subcategory of $K(\mathcal{A})$ of complexes bounded below. Similarly we define $K^-(\mathcal{A})$ (bounded above) and $K^b(\mathcal{A})$ (bounded on both sides).

We now define a triangulated structure on $K(\mathcal{A})$ (resp. $K^+(\mathcal{A})$, etc.). Let T be the left shift functor with sign change:

$$T(X^\bullet)^p = X^{p+1}, \quad d_{T(X)} = -d_X.$$

We often write $X^\bullet[1]$ instead of $T(X^\bullet)$, and $X^\bullet[n]$ instead of $T^n(X^\bullet)$. If $u: X^\bullet \rightarrow Y^\bullet$ is any morphism, consider its *mapping cone* $Z^\bullet$ of u , defined as the complex

$$Z^\bullet = T(X^\bullet) \oplus Y^\bullet$$

with differential

$$d_Z = \begin{pmatrix} T(d_X) & T(u) \\ 0 & d_Y \end{pmatrix}.$$

There are natural morphisms

$$v: Y^\bullet \rightarrow Z^\bullet, \quad w: Z^\bullet \rightarrow T(X^\bullet).$$

We define a triangle in $K(\mathcal{A})$ to be any sextuple isomorphic to one obtained from a morphism u by this construction.

All axioms (TR1)–(TR4) are easily verified, once one notes that the mapping cone of the identity map $\text{id}_{X^\bullet}$ is homotopic to 0. Indeed, the mapping cone is $T(X^\bullet) \oplus X^\bullet$ as described above, matrix

$$k = \begin{pmatrix} 0 & 0 \\ \text{id}_X & 0 \end{pmatrix}$$

is a homotopy operator.

Definition 3. We define H to be the functor from $K(\mathcal{A})$ to $\mathcal{A}$ which sends a complex $X^\bullet$ to its 0^{th} cohomology:

$$H(X^\bullet) = \ker d^0 / \operatorname{im} d^{-1}.$$

This is indeed a functor, since homotopic maps induce the same map on cohomology. We write $H^i = H \circ T^i$ for any $i \in \mathbb{Z}$.

Observe that H is a cohomological functor from $K(\mathcal{A})$ to $\mathcal{A}$. Indeed, it suffices to check the long exact sequence for triangles coming from mapping cones, which is straightforward.

2.4 Localization of categories

Definition 4. Let $\mathcal{C}$ be a category. A collection S of morphisms of $\mathcal{C}$ is called a multiplicative system if it satisfies the following axioms (FR1)–(FR3):

(FR1) If $f, g \in S$ and the composite fg exists, then $fg \in S$. For any $X \in \operatorname{Ob} \mathcal{C}$, $\operatorname{id}_X \in S$.

(FR2) Any diagram

$$X \xrightarrow{u} Y \xleftarrow{s} Z, \quad s \in S$$

can be completed to a commutative diagram

$$\begin{array}{ccc} W & \xrightarrow{v} & Z \\ t \downarrow & & \downarrow s \\ X & \xrightarrow{u} & Y \end{array}$$

with $t \in S$. The analogous statement holds with all arrows reversed.

(FR3) Let $f, g: X \rightarrow Y$ be morphisms in $\mathcal{C}$. The following are equivalent:

- (i) There exists $s: Y \rightarrow Y'$ in S such that $sf = sg$;
- (ii) There exists $t: X' \rightarrow X$ in S such that $ft = gt$.

Definition 5. Let $\mathcal{C}$ be a category and S a collection of morphisms in $\mathcal{C}$. The localization of $\mathcal{C}$ with respect to S is a category $\mathcal{C}_S$ together with a functor $Q: \mathcal{C} \rightarrow \mathcal{C}_S$ such that:

- (a) $Q(s)$ is an isomorphism for every $s \in S$;

(b) Any functor $F: \mathcal{C} \rightarrow \mathcal{D}$ sending each $s \in S$ to an isomorphism factors uniquely through Q .

Remark. One can show that such a localization exists without hypotheses on S , but we will not need this result.

Proposition 2. Let $\mathcal{C}$ be a category and S a multiplicative system in $\mathcal{C}$. The localization $\mathcal{C}_S$ may be constructed as follows: $\text{Ob } \mathcal{C}_S = \text{Ob } \mathcal{C}$, and for all $X, Y \in \text{Ob } \mathcal{C}$,

$$\text{Hom}_{\mathcal{C}_S}(X, Y) = \varinjlim_{I_X} \text{Hom}_{\mathcal{C}}(X', Y),$$

where I_X is the category whose objects are morphisms $s: X' \rightarrow X$ in S , and morphisms are commutative diagrams:

$$\begin{array}{ccc} X'' & \xrightarrow{f} & X' \\ & \searrow s_2 & \downarrow s_1 \\ & & X \end{array}$$

Furthermore, if $\mathcal{C}$ is an additive category, then $\mathcal{C}_S$ is also additive.

Proof. First observe, using (FR1), (FR2), and (FR3) that the category I_X satisfies the axioms L1, L2, L3 of [GT, Ch. I, §1], and hence behaves as well as an inductive system for taking limits. Thus a morphism of X to Y in $\mathcal{C}_S$ is represented by a diagram

$$\begin{array}{ccc} & X' & \\ s \swarrow & & \searrow a \\ X & & Y \end{array}$$

with $s \in S$. This diagram defines the same morphism as another one with $t \in S$

$$\begin{array}{ccc} & X'' & \\ t \swarrow & & \searrow b \\ X & & Y \end{array}$$

if and only if there is a morphism $u: X''' \rightarrow X$ in S and morphisms $f: X''' \rightarrow X'$ and $g: X''' \rightarrow X''$ such that $sf = u = tg$ and $af = bg$.

To compose morphisms

$$\begin{array}{ccc} & X' & \\ s \swarrow & & \searrow a \\ X & & Y \end{array} \quad \text{and} \quad \begin{array}{ccc} & Y' & \\ t \swarrow & & \searrow b \\ Y & & Z \end{array}$$

we use (FR2) to find a commutative diagram

$$\begin{array}{ccccc} & & X'' & & \\ & t' \swarrow & & \searrow c & \\ & X' & & Y' & \\ s \swarrow & & a \searrow & t \swarrow & \searrow b \\ X & & Y & & Z \end{array}$$

with $t' \in S$, and then take X'' , st' , bc to be the composition. One verifies easily that the resulting morphism of X to Z is independent of the representatives of the morphisms of X to Y and Y to Z chosen, and is also independent of the commutative diagram chosen.

One can also verify easily that the functor $Q : C \rightarrow C_S$ has the properties required and that C_S is additive if C is (again using L1, L2, L3 to show that the $\varinjlim$ is a group).

□

Definition 6. Let $\mathcal{C}$ be a triangulated category and S a multiplicative system in $\mathcal{C}$. We say S is compatible with the triangulation if:

(FR4) $s \in S \implies T(s) \in S$, where T is the translation functor;

(FR5) Axiom (TR3) holds with the extra condition: if $f, g \in S$ are the given morphisms of triangles, then $h \in S$.

Proposition 3. If $\mathcal{C}$ is triangulated and S a multiplicative system compatible with triangulation, then C_S has a unique structure of triangulated category such that Q is a ∂ -functor, and Q has the universal property b) above for ∂ -functors into triangulated categories.

Proof. Left to the reader. It helps to observe that one can also calculate $\text{Hom}_{C_S}(X, Y)$ as

$$\varinjlim_{J_Y} \text{Hom}_C(X, Y')$$

where J_Y is the category whose objects are morphisms $s : Y \longrightarrow Y'$ in S , and whose maps are commutative diagrams

$$\begin{array}{ccc} & Y & \\ s_1 \swarrow & & \searrow s_2 \\ Y' & \xrightarrow{f} & Y'' \end{array} .$$

□

Proposition 4. *Let $\mathcal{C}$ be a category, S a multiplicative system in $\mathcal{C}$, and $\mathcal{D} \subseteq \mathcal{C}$ a full subcategory (i.e. $X, Y \in \text{Ob } \mathcal{D}$ implies $\text{Hom}_{\mathcal{C}}(X, Y) = \text{Hom}_{\mathcal{D}}(X, Y)$) such that $S \cap \mathcal{D}$ is a multiplicative system in $\mathcal{D}$. Assume furthermore that one of the following two conditions holds:*

- (i) *For every $s : X' \rightarrow X$ in S with $X \in \text{Ob } \mathcal{D}$, there exists $f : X'' \rightarrow X'$ with $X'' \in \text{Ob } \mathcal{D}$ and $sf \in S$;*
- (ii) *The same with all arrows reversed.*

Then the natural functor

$$\mathcal{D}_{S \cap \mathcal{D}} \rightarrow \mathcal{C}_S$$

is fully faithful, i.e. $\mathcal{D}_{S \cap \mathcal{D}}$ can be identified with a full subcategory of $\mathcal{C}_S$.

Proposition 5. *Let $\mathcal{C}$ be a category, S a multiplicative system, $Q : \mathcal{C} \rightarrow \mathcal{C}_S$ the localization functor. Let $\mathcal{D}$ be any category, and $F, G : \mathcal{C}_S \rightarrow \mathcal{D}$ functors. Then the natural restriction map*

$$\alpha : \text{Hom}(F, G) \rightarrow \text{Hom}(FQ, GQ)$$

between functor morphism sets is bijective.

Proof. To give a morphism of functors $F \longrightarrow G$ means to give for each $X \in \text{Ob } \mathcal{C}_S$, a morphism $F(X) \longrightarrow G(X)$, such that if $X \longrightarrow Y$ is a morphism, then

$$\begin{array}{ccc} F(X) & \longrightarrow & F(Y) \\ \downarrow & & \downarrow \\ G(X) & \longrightarrow & G(Y) \end{array}$$

is a commutative diagram. Thus, since $\text{Ob } \mathcal{C} = \text{Ob } \mathcal{C}_S$, the map α is injective. To show α surjective, suppose given a morphism $FQ \longrightarrow GQ$. Then we have a

morphism $F(X) \rightarrow G(X)$ for each $X \in \text{Ob } C = \text{Ob } C_S$, and commutative diagrams for morphisms $X \rightarrow Y$ in C . A morphism in C_S is represented by a diagram

$$\begin{array}{ccc} & X' & \\ s \swarrow & & \searrow f \\ X & & Y \end{array}$$

of morphisms in C , with $s \in S$. But for $s \in S$, $F(s)$ and $G(s)$ are isomorphisms, so we get the required commutative diagram. $\square$

2.5 Quasi-isomorphisms and the derived category

Let $\mathcal{A}$ be an abelian category, and let $K(\mathcal{A})$ be the triangulated category of complexes up to homotopy introduced in §2. A morphism $f: X^\bullet \rightarrow Y^\bullet$ in $K(\mathcal{A})$ is called a *quasi-isomorphism* if it induces an isomorphism on all cohomology groups. Denote by Qis the collection of all quasi-isomorphisms in $K(\mathcal{A})$.

Proposition 6. *Qis is a multiplicative system in $K(\mathcal{A})$.*

Proof. This is a consequence of the following more general proposition $\square$

Proposition 7. *Let $\mathcal{C}$ be a triangulated category, $\mathcal{A}$ an abelian category, and $H: \mathcal{C} \rightarrow \mathcal{A}$ a cohomological functor. Let S be the class of morphisms $s \in \mathcal{C}$ such that $H(T^i(s))$ is an isomorphism for all $i \in \mathbb{Z}$. Then S is a multiplicative system in $\mathcal{C}$ and is compatible with the triangulation.*

Proof. We verify axioms (FR1)–(FR5). Axioms (FR1) and (FR4) are trivial. Axiom (FR5) follows from the long exact sequence of a cohomological functor and the five-lemma.

To prove (FR2), let a diagram is given:

$$X \xrightarrow{u} Y \xleftarrow{s} Z$$

with $s \in S$. Complete s to a triangle (Z, Y, N, s, f, g) . Complete fu to a triangle (W, X, N, t, fu, h) . Then (u, id_N) is a map of two sides of the second triangle into

the first, so there is a map $v : W \rightarrow Z$ giving a morphism of triangles.

$$\begin{array}{ccccc}
 W & \xrightarrow{v} & Z & & \\
 \downarrow t & \nearrow h & \downarrow & \nwarrow g & \\
 & N & \xrightarrow{\text{id}_N} & N & \\
 \nearrow fu & & \downarrow s & \nearrow f & \\
 X & \xrightarrow{u} & Y & &
 \end{array}$$

Now $sv = ut$, so it remains to prove $t \in S$. Indeed, since $s \in S$, we have $H(T^i(N)) = 0$ for all $i \in \mathbb{Z}$ by the long exact sequence of the first triangle. Applying this to the long exact sequence of the second triangle, we find $H(T^i(t))$ is an isomorphism for all $i \in \mathbb{Z}$.

The opposed statement of (FR2) is proved similarly.

To prove (FR3), we consider the morphism $f - g$, and reduce to showing the following two properties equivalent (where $f : X \rightarrow Y$ is a morphism):

1. There exists an $s : Y \rightarrow Y' \in S$ such that $sf = 0$
2. There exists an $t : X' \rightarrow X \in S$ such that $ft = 0$

Suppose (i) holds.

$$\begin{array}{ccccc}
 X' & \xleftarrow{w} & Z & & \\
 \downarrow t & \nearrow g & \downarrow v & \nwarrow u & \\
 X & \xrightarrow{f} & Y & \xrightarrow{s} & Y'
 \end{array}$$

By (TR1) and (TR2) we can find a triangle (Z, Y, Y', v, s, u) for suitable Z . Now $sf = 0$, so by Proposition 1.1 b), there is a map $g : X \rightarrow Z$ such that $f = vg$. Again by (TR1) and (TR2) we can find a triangle (X', X, Z, t, g, w) for suitable X' . By the same Proposition applied to this second triangle, the existence of v implies that $ft = 0$. We need only show that $t \in S$. Since $s \in S$, $H(T^i(Z)) = 0$ for all $i \in \mathbb{Z}$, by the long exact sequence of cohomology. In turn, this implies that $t \in S$.

The implication (ii) $\implies$ (i) is analogous. □

Definition 7. The derived category of $\mathcal{A}$ is

$$D(\mathcal{A}) := K(\mathcal{A})_{\text{Qis}}.$$

Similarly define

$$D^+(\mathcal{A}) = K^+(\mathcal{A})_{\text{Qis}}, \quad D^-(\mathcal{A}) = K^-(\mathcal{A})_{\text{Qis}}, \quad D^b(\mathcal{A}) = K^b(\mathcal{A})_{\text{Qis}}.$$

By Proposition 1.3.3, these are full subcategories of $D(\mathcal{A})$, and

$$D^+(\mathcal{A}) \cap D^-(\mathcal{A}) = D^b(\mathcal{A}).$$

Definition 8. Let $\mathcal{A}$ be abelian, and $\mathcal{A}' \subset \mathcal{A}$ a thick full abelian subcategory (closed under extensions in $\mathcal{A}$). Define $K_{\mathcal{A}'}(\mathcal{A})$ to be the full subcategory of $K(\mathcal{A})$ consisting of complexes $X^\bullet$ whose cohomology objects $H^i(X^\bullet)$ lie in $\mathcal{A}'$. Since $\mathcal{A}'$ is thick, $K_{\mathcal{A}'}(\mathcal{A})$ is a triangulated subcategory of $K(\mathcal{A})$.

Define

$$D_{\mathcal{A}'}(\mathcal{A}) := K_{\mathcal{A}'}(\mathcal{A})_{\text{Qis}}.$$

By Proposition 1.3.3, $D_{\mathcal{A}'}(\mathcal{A})$ is the full subcategory of $D(\mathcal{A})$ of complexes with all cohomology in $\mathcal{A}'$. One defines similarly $K_{\mathcal{A}'}^+(\mathcal{A})$, $D_{\mathcal{A}'}^+(\mathcal{A})$, etc., for bounded-below complexes.

Example. To help understand the category $D(\mathcal{A})$, let us ask: when does a morphism of complexes $f: X \rightarrow Y$ give the zero map in $D(\mathcal{A})$? The condition is:

(*) There exists an $s: Y \rightarrow Y'$ in Qis such that sf is homotopic to zero (or, equivalently, there exists a $t: X' \rightarrow X$ in Qis such that ft is homotopic to zero).

1. Of course, if $f \sim 0$ (f homotopic to zero), then f satisfies (*). The converse is false. For example, take X to be the complex

$$0 \rightarrow \mathbb{Z} \xrightarrow{2} \mathbb{Z} \rightarrow \mathbb{Z}/2\mathbb{Z} \rightarrow 0,$$

and f to be the identity $\text{id}_X: X \rightarrow X$. Let $g: X \rightarrow 0$ be the zero map. Then $g \in \text{Qis}$, and $gf = 0$, but f is not homotopic to zero, as one sees easily.

2. If f satisfies (*), then f induces the zero map on cohomology, but the converse is false. For example take $f: X \rightarrow Y$ as follows:

$$\begin{array}{ccccccc} \cdots & \longrightarrow & 0 & \longrightarrow & \mathbb{Z} & \xrightarrow{2} & \mathbb{Z} & \longrightarrow & 0 & \longrightarrow & \cdots \\ & & \downarrow & & \downarrow 1 & & \downarrow 2 & & \downarrow & & \\ \cdots & \longrightarrow & 0 & \longrightarrow & \mathbb{Z} & \xrightarrow{1} & \mathbb{Z}/3\mathbb{Z} & \longrightarrow & 0 & \longrightarrow & \cdots \end{array}$$

Now f induces the zero map on cohomology, but there does not exist $t: X' \rightarrow X$ in Qis such that $ft \sim 0$. (The reader can supply the proof as follows: take a cycle $x \in X'$ such that $t(x)$ generates the single cohomology group $\mathbb{Z}/2\mathbb{Z}$ of X . If k is a homotopy operator for ft , show that $2k(x) = 1$, which is impossible.)

So for two maps $f, g: X \rightarrow Y$ of complexes, we see that the following implications are all strict:

$$\begin{aligned} f = g &\implies f \text{ homotopic to } g \\ &\implies f \text{ and } g \text{ give the same morphism in } D(\mathcal{A}) \\ &\implies f \text{ and } g \text{ give the same map on cohomology.} \end{aligned}$$

Proposition 8. *The functor $\mathcal{A} \rightarrow D(\mathcal{A})$ which sends each object X of $\mathcal{A}$ into the complex consisting of X in degree 0, and 0 elsewhere, gives an equivalence of the category $\mathcal{A}$ with the full subcategory of $D(\mathcal{A})$ consisting of those complexes $X^\bullet$ such that $H^i(X^\bullet) = 0$ for $i \neq 0$.*

We now prove three lemmas, which give an alternative description of $D^+(\mathcal{A})$ when $\mathcal{A}$ has enough injectives.

Lemma 1. *Let $\mathcal{A}$ be abelian, $f: Z^\bullet \rightarrow I^\bullet$ a morphism of complexes. Assume:*

- (1) $Z^\bullet$ is acyclic;
- (2) Each I^p is injective in $\mathcal{A}$;
- (3) $I^\bullet$ is bounded below.

Then f is homotopic to zero.

Proof. Well known (and easy). □

Lemma 2. *Let $\mathcal{A}$ be abelian, $s: I^\bullet \rightarrow Y^\bullet$ a morphism of complexes. Assume:*

- (1) s induces an isomorphism on cohomology;
- (2) Each I^p is injective;
- (3) $I^\bullet$ is bounded below.

Then s admits a homotopy inverse.

Proof. Suppose given $s: I^\bullet \rightarrow Y^\bullet$ as above. Let $Z^\bullet = T(I^\bullet) \oplus Y^\bullet$ be the mapping cone of s . Then $Z^\bullet$ is acyclic, and so the triangular morphism $v: Z^\bullet \rightarrow T(I^\bullet)$ satisfies the conditions of Lemma 1.4.4, and so is homotopic to zero. Let us call the homotopy operator

$$(k, t): T(I^\bullet) \oplus Y^\bullet \longrightarrow I^\bullet.$$

Then we have the equation

$$v = (\text{id}_{I^\bullet}, 0) = (k, t)d_Z + d_{I^\bullet}(k, t).$$

Separating the components, we find

$$\text{id}_{T(I^\bullet)} = dk + kd + ts$$

and

$$dt - td = 0.$$

Thus $t: Y^\bullet \rightarrow I^\bullet$ is a morphism of complexes, and $\text{id}_{I^\bullet}$ is homotopic to ts , so t is a homotopy inverse of s . $\square$

Lemma 3. *Let $\mathcal{A}$ be an abelian category.*

1) *Let $P \subseteq \text{Ob } \mathcal{A}$ and assume:*

(i) *Every object of $\mathcal{A}$ admits an injection into an element of P .*

Then every complex $X^\bullet \in K^+(\mathcal{A})$ admits a quasi-isomorphism into a bounded-below complex $I^\bullet$ of objects of P .

2) *Assume further that P satisfies:*

(ii) *If $0 \rightarrow X \rightarrow Y \rightarrow Z \rightarrow 0$ is exact and $X \in P$, then $Y \in P \iff Z \in P$;*

(iii) *There exists a positive integer n such that for any exact sequence*

$$X^0 \rightarrow X^1 \rightarrow \cdots \rightarrow X^{n-1} \rightarrow X^n \rightarrow 0,$$

if $X^0, \dots, X^{n-1} \in P$ then $X^n \in P$.

Then every $X^\bullet \in K(\mathcal{A})$ admits a quasi-isomorphism into a complex $I^\bullet$ of objects of P .

3) *Let $\mathcal{A}'$ be a thick subcategory of $\mathcal{A}$, and suppose $\mathcal{A}'$ has enough $\mathcal{A}$ -injectives. Then every $X^\bullet \in K_{\mathcal{A}'}^+(\mathcal{A})$ admits a quasi-isomorphism into a bounded-below complex $I^\bullet$ of $\mathcal{A}$ -injective objects of $\mathcal{A}'$.*

Proof of 1). We may assume $X^p = 0$ for $p < 0$. Embed $X^0 \hookrightarrow I^0$ with $I^0 \in P$. Having defined $I^0, I^1, \dots, I^p$, choose $I^{p+1} \in P$ containing

$$I^p / \text{im } I^{p-1} \oplus X^{p+1},$$

and define differentials $I^p \rightarrow I^{p+1}$ and morphisms $X^{p+1} \rightarrow I^{p+1}$ canonically. One verifies $X^\bullet \rightarrow I^\bullet$ is a quasi-isomorphism. Note that in this construction all the maps $X^p \rightarrow I^p$ is injective. $\square$

Proof of 2). Let $X^\bullet$ be any complex. For a fixed integer i_0 , by 1), we can find a quasi-isomorphism of the truncated complex

$$0 \rightarrow 0 \rightarrow X^{i_0} \rightarrow X^{i_0+1} \rightarrow \dots$$

into a complex $I^\bullet$ of objects of P , with each $X^i \rightarrow I^i$ injective. Define $X_0^\bullet$ by

$$\dots \rightarrow X^{i_0-2} \rightarrow X^{i_0-1} \rightarrow I^{i_0} \rightarrow I^{i_0+1} \rightarrow \dots$$

Then $X^\bullet \rightarrow X_0^\bullet$ is a quasi-isomorphism, $X_0^i \in P$ for $i \geq i_0$, and each $X^i \rightarrow X_0^i$ is injective.

Suppose given a complex $X_1^\bullet$ with $X_1^i \in P$ for $i \geq i_1$, and let $i_2 < i_1$. Then we will construct a quasi-isomorphism $X_1^\bullet \rightarrow X_2^\bullet$ such that $X_2^i \in P$ for $i \geq i_2$, and $X_1^i = X_2^i$ for $i \geq i_1 + n$. (Here n is the integer of condition (iii) above.) Indeed, by the first step above, we can find a quasi-isomorphism $X_1^\bullet \rightarrow X'^\bullet$ such that $X'^i \in P$ for $i \geq i_2$, and each $X_1^i \rightarrow X'^i$ is injective. Let $Y^i = \text{coker}(X_1^i \rightarrow X'^i)$. Then $Y^\bullet$ is an acyclic complex, and $Y^i \in P$ for $i \geq i_1$, by condition (ii) above. Hence $B^i(Y^\bullet) \in P$ for $i \geq i_1 + n$, by condition (iii). Now define $X_2^\bullet$ by

$$X_2^i = \begin{cases} X'^i & \text{for } i < i_1 + n, \\ B^i(X'^\bullet) \oplus_{X_1^{i-1}} X_1^i & \text{for } i = i_1 + n, \\ X_1^i & \text{for } i > i_1 + n. \end{cases}$$

One sees easily that $X_1^\bullet \rightarrow X_2^\bullet$ is a quasi-isomorphism. It follows from (ii) and the exact sequence

$$0 \rightarrow X_1^i \rightarrow B^i(X'^\bullet) \oplus_{X_1^{i-1}} X_1^i \rightarrow B^i(Y^\bullet) \rightarrow 0$$

that the middle term is in P , for $i \geq i_1 + n$, so $X_2^\bullet$ is as required.

Now, given a complex $X^\bullet \in K(\mathcal{A})$, choose a sequence of integers $i_0 > i_1 > i_2 > \dots$ tending to $-\infty$. Choose $X_0^\bullet$ for i_0 as in the first step, and choose $X_1^\bullet, X_2^\bullet, \dots$ for $i_1, i_2, \dots$ successively as in the second step. Then we have quasi-isomorphisms

$$X^\bullet \rightarrow X_0^\bullet \rightarrow X_1^\bullet \rightarrow X_2^\bullet \rightarrow \dots,$$

and for each i , the sequence $X^i \rightarrow X_0^i \rightarrow X_1^i \rightarrow \dots$ is eventually constant, and eventually in P . Hence $I^\bullet = \varinjlim X_r^\bullet$ is the required complex of objects of P . $\square$

Proof of 3). We may assume $X^i = 0$ for $i < 0$. Embed $H^0(X^\bullet)$ in I^0 , an $\mathcal{A}$ -injective of $\mathcal{A}'$, which is possible since $H^0(X^\bullet) \in \text{Ob } \mathcal{A}'$. Extend this to a map $f^0: X^0 \rightarrow I^0$, which is possible since I^0 is $\mathcal{A}$ -injective.

Having defined $I^0, I^1, \dots, I^p$ and $f^i: X^i \rightarrow I^i$ for $i = 0, \dots, p$, choose I^{p+1} to be an $\mathcal{A}$ -injective of $\mathcal{A}'$ containing

$$I^p / \text{im } I^{p-1} \oplus_{X^p} Z^{p+1}(X^\bullet). \quad (*)$$

We must check that this latter is in $\mathcal{A}'$. Indeed, $\mathcal{A}'$ is a thick subcategory, so it is sufficient to note that $I^p / \text{im } I^{p-1} \in \mathcal{A}'$ (one shows by induction that $B^i(I^\bullet)$ and $Z^i(I^\bullet)$ are in $\mathcal{A}'$ for all i), and that the quotient of $(*)$ by $I^p / \text{im } I^{p-1}$ is $H^{p+1}(X^\bullet)$, which is in $\mathcal{A}'$ by hypothesis. Extend the natural map $Z^{p+1}(X^\bullet) \rightarrow I^{p+1}$ to a map $f^{p+1}: X^{p+1} \rightarrow I^{p+1}$. One checks easily that the resulting map $f: X^\bullet \rightarrow I^\bullet$ is a quasi-isomorphism, as required. $\square$

Proposition 9. *Let $\mathcal{A}$ be abelian, and let $\mathcal{I}$ be the additive subcategory of injective objects of $\mathcal{A}$. The natural functor*

$$\alpha: K^+(\mathcal{I}) \rightarrow D^+(\mathcal{A})$$

is fully faithful. (Note that the results of section 3 carry over to additive subcategories of abelian categories.) Furthermore, if $\mathcal{A}$ has enough injectives (i.e., if every object of $\mathcal{A}$ admits an injection into an injective object), then α is an equivalence of categories.

Proof. We note that $\text{Qis} \cap K^+(\mathcal{I})$ is a multiplicative system in $K^+(\mathcal{I})$, by Proposition 1.4.2, and we observe by Lemma 1.4.5 that condition (ii) of Proposition 1.3.3 is satisfied for $K^+(\mathcal{I}) \subseteq K^+(\mathcal{A})$ and Qis . Hence the natural functor

$$D^+(\mathcal{I}) \longrightarrow D^+(\mathcal{A})$$

is fully faithful. But on the other hand, Lemma 1.4.5 shows also that every quasi-isomorphism in $K^+(\mathcal{I})$ is an isomorphism, hence $K^+(\mathcal{I}) = D^+(\mathcal{I})$.

Now if $\mathcal{A}$ has enough injectives, applying Lemma 1.4.6 in the case $\mathcal{A} = \mathcal{B}$ and $P =$ the injective objects, we see that every object of $D^+(\mathcal{A})$ is isomorphic to an object in $K^+(\mathcal{I})$, so α is an equivalence of categories. $\square$

Proposition 10. *Let $\mathcal{A}$ be abelian, $\mathcal{A}' \subseteq \mathcal{A}$ a thick abelian subcategory. Assume $\mathcal{A}'$ has enough $\mathcal{A}$ -injectives, i.e. every object of $\mathcal{A}'$ can be injected into an $\mathcal{A}$ -injective object of $\mathcal{A}'$. Then the natural functor*

$$D^+(\mathcal{A}') \rightarrow D_{\mathcal{A}'}^+(\mathcal{A})$$

is an equivalence of categories.

Proof. We apply Proposition 1.3.3 to the inclusion $K^+(\mathcal{A}') \hookrightarrow K^+(\mathcal{A})$. Clearly Qis is a multiplicative system in each. If $X^\bullet \rightarrow Y^\bullet$ is a quasi-isomorphism with $X^\bullet \in K^+(\mathcal{A}')$, then $Y^\bullet$ has cohomology in $\mathcal{A}'$, and so by Lemma 1.4.6 admits a quasi-isomorphism $Y^\bullet \rightarrow I^\bullet$ with $I^\bullet \in K^+(\mathcal{A}')$, each I^p injective in $\mathcal{A}$. Hence condition (ii) is satisfied, and so the functor

$$D^+(\mathcal{A}') \longrightarrow D^+(\mathcal{A})$$

is fully faithful. The same Lemma 1.4.6 also shows that the image is $D_{\mathcal{A}'}^+(\mathcal{A})$. $\square$

Exercise. We leave to the reader the analogous statements of the last five results in the case of projective objects of $\mathcal{A}$ and $D^-(\mathcal{A})$.

2.6 Derived functors

We will treat only the question of *right derived covariant functors*, leaving the reader to make the obvious modifications for left derived covariant functors, and right and left derived contravariant functors.

Let $\mathcal{A}, \mathcal{B}$ be abelian categories, and let $F: K(\mathcal{A}) \rightarrow K(\mathcal{B})$ be a ∂ -functor (cf. §1). Such is the case, for example, if we are given an additive functor $F: \mathcal{A} \rightarrow \mathcal{B}$. It extends to $K(\mathcal{A})$.

In general, F will not take quasi-isomorphisms into quasi-isomorphisms – to say that it does is to say that it localizes and gives rise to a functor from $D(\mathcal{A}) \rightarrow D(\mathcal{B})$. That will be the case, for example, if F is an exact functor.

Thus we are led to ask if there is a functor from $D(\mathcal{A})$ to $D(\mathcal{B})$ which is at least close to F , and this gives rise to the notion of derived functor below. Before giving the definition we generalize slightly.

Definition 9. Let $\mathcal{A}$ be abelian, and let $K^*(\mathcal{A}) \subseteq K(\mathcal{A})$ be a triangulated subcategory. By Proposition 1.4.2, $K^*(\mathcal{A}) \cap \text{Qis}$ is a multiplicative system in $K^*(\mathcal{A})$.

We say $K^*(\mathcal{A})$ is a localizing subcategory of $K(\mathcal{A})$ if the natural functor

$$K^*(\mathcal{A})_{K^*(\mathcal{A}) \cap \text{Qis}} \rightarrow K(\mathcal{A})_{\text{Qis}} = D(\mathcal{A})$$

is fully faithful. In this case we write $D^*(\mathcal{A})$ for the $K^*(\mathcal{A})_{K^*(\mathcal{A}) \cap \text{Qis}}$.

Examples.

1. The intersection of any family of localizing subcategories is localizing.
2. $K^+(\mathcal{A}), K^-(\mathcal{A}), K^b(\mathcal{A})$ are localizing subcategories of $K(\mathcal{A})$ (cf. §4).

3. If $\mathcal{A}' \subseteq \mathcal{A}$ is a thick subcategory, then $K_{\mathcal{A}'}(\mathcal{A}), K_{\mathcal{A}'}^+(\mathcal{A}), K_{\mathcal{A}'}^-(\mathcal{A}), K_{\mathcal{A}'}^b(\mathcal{A})$ are localizing subcategories of $K(\mathcal{A})$ (cf. §4).
4. The complexes of finite injective dimension form a localizing subcategory $K^+(\mathcal{A})_{\text{fid}} \subseteq K(\mathcal{A})$ (cf. Corollary 1.7.7).

Definition 10. Let $\mathcal{A}, \mathcal{B}$ be abelian categories, $K^*(\mathcal{A}) \subseteq K(\mathcal{A})$ a localizing subcategory, and

$$F: K^*(\mathcal{A}) \rightarrow K(\mathcal{B})$$

a ∂ -functor.

Denote by Q the localization functor $K^*(\mathcal{A}) \rightarrow D^*(\mathcal{A})$ (resp. $K(\mathcal{B}) \rightarrow D(\mathcal{B})$). The right derived functor of F is a ∂ -functor

$$\underline{\underline{R}}^*F: D^*(\mathcal{A}) \rightarrow D(\mathcal{B})$$

together with a functor morphism

$$\xi: Q \circ F \rightarrow \underline{\underline{R}}^*F \circ Q$$

of functors $K^*(\mathcal{A}) \rightarrow D(\mathcal{B})$, satisfying the following universal property:

If $G: D^*(\mathcal{A}) \rightarrow D(\mathcal{B})$ is a ∂ -functor and any functor morphism

$$\zeta: Q \circ F \rightarrow G \circ Q,$$

there exists a unique functor morphism

$$\eta: \underline{\underline{R}}^*F \rightarrow G$$

such that

$$\zeta = (\eta \circ Q) \circ \xi.$$

Remarks.

1. If $\underline{\underline{R}}^*F$ exists, it is unique up to unique isomorphism of functors.
2. For standard localizing subcategories $K^+(\mathcal{A}), K^-(\mathcal{A}), K_{\mathcal{A}'}(\mathcal{A})$, we write

$$\underline{\underline{R}}^+F, \quad \underline{\underline{R}}^-F, \quad \underline{\underline{R}}_{\mathcal{A}'}F$$

respectively; when no confusion arises, we write simply $\underline{\underline{R}}F$.

3. We will write $R^i F$ for $H^i(\underline{R}F)$, and it will follow from the results below that if F comes from a left-exact functor $F : A \rightarrow B$, and if A has enough injectives, then these are the usual derived functors of F .
4. If $\varphi : F \rightarrow G$ is a morphism of functors from $K^*(A)$ to $K(B)$, and if $\underline{R}F$ and $\underline{R}G$ both exist, then there is a unique morphism of functors

$$\underline{R}\varphi : \underline{R}F \rightarrow \underline{R}G$$

compatible with the ξ 's. This follows immediately from the definition.

5. Let $K^{**}(\mathcal{A}) \subseteq K^*(\mathcal{A})$ be localizing subcategories, if $F : K^*(\mathcal{A}) \rightarrow K(\mathcal{B})$ is a ∂ -functor. and if $\underline{R}^*F$ and $\underline{R}^{**}(F|_{K^{**}(\mathcal{A})})$ exist, there is a natural functor morphism

$$\underline{R}^{**}(F|_{K^{**}(\mathcal{A})}) \rightarrow \underline{R}^*F|_{D^{**}(\mathcal{A})}.$$

We do not know if this is an isomorphism in general, but it holds in all our intended applications (cf. Corollary 1.5.3).

Theorem 1 (Existence of Derived Functors). *Let $\mathcal{A}, \mathcal{B}, K^*(\mathcal{A}), F$ be as above. Suppose there exists a triangulated subcategory $\mathcal{L} \subseteq K^*(\mathcal{A})$ such that:*

- (1) *Every object of $K^*(\mathcal{A})$ admits a quasi-isomorphism into some object of $\mathcal{L}$;*
- (2) *For every acyclic complex $I^\bullet \in \text{Ob } \mathcal{L}$ (i.e. $H^i(I^\bullet) = 0$ for all i), the complex $F(I^\bullet)$ is also acyclic.*

*Then the right derived functor $(\underline{R}^*F, \xi)$ exists. Furthermore, for any $I^\bullet \in \text{Ob } \mathcal{L}$, the map*

$$\xi(I^\bullet) : Q \circ F(I^\bullet) \rightarrow \underline{R}^*F \circ Q(I^\bullet)$$

is an isomorphism in $D(\mathcal{B})$.

Proof. First, the restriction $F|_{\mathcal{L}}$ sends quasi-isomorphisms to quasi-isomorphisms. Indeed, let $s : I_1^\bullet \rightarrow I_2^\bullet$ be a quasi-isomorphism in $\mathcal{L}$; complete s to a triangle in $K^*(\mathcal{A})$, and let $J^\bullet$ be the third side of that triangle. Then $J^\bullet$ is acyclic, so $F(J^\bullet)$ is acyclic. Hence $F(s)$ is a quasi-isomorphism.

Thus F passes to the quotient to give a functor

$$\overline{F} : \mathcal{L}_{Q\text{is}} \rightarrow D(\mathcal{B})$$

satisfying $\overline{F} \circ Q = Q \circ F$ on $\mathcal{L}$ (we denote as usual by Q the morphism from a category to its localization).

Second, by the condition of Proposition 1.3.3, (ii) are satisfied for $\mathcal{L}, K^*(\mathcal{A}), \text{Qis}$, and so the natural functor

$$T: \mathcal{L}_{\text{Qis}} \rightarrow D^*(\mathcal{A})$$

is an equivalence of categories, using 1) above. Let U be a quasi-inverse of T , i.e. a functor $U: D^*(\mathcal{A}) \rightarrow \mathcal{L}_{\text{Qis}}$ with functorial isomorphisms

$$\alpha: \text{id}_{\mathcal{L}_{\text{Qis}}} \rightarrow U \circ T, \quad \beta: \text{id}_{D^*(\mathcal{A})} \rightarrow T \circ U.$$

Define

$$\underline{\underline{R}}^*F := \overline{F} \circ U.$$

We now define the functor morphism

$$\xi: Q \circ F \rightarrow \underline{\underline{R}}^*F \circ Q = \overline{F} \circ U \circ Q$$

as follows. Given $X^\bullet \in \text{Ob } K^*(\mathcal{A})$, choose $I^\bullet \in \text{Ob } \mathcal{L}$ such that $Q(I^\bullet) \cong U \circ Q(X^\bullet)$ in $D^*(\mathcal{A})$.

We have an isomorphism in $D^*(\mathcal{A})$,

$$\beta(Q(X^\bullet)): Q(X^\bullet) \xrightarrow{\sim} T \circ U(Q(X^\bullet)) = T(Q(I^\bullet)).$$

This isomorphism may be represented by a diagram in $K^*(\mathcal{A})$

$$X^\bullet \xrightarrow{t} Y^\bullet \xleftarrow{s} I^\bullet \quad (*)$$

with $s, t \in \text{Qis}$ and $Y^\bullet \in \text{Ob } K^*(\mathcal{A})$. Furthermore by hypothesis (1), we may assume $Y^\bullet \in \text{Ob } \mathcal{L}$. Applying the functor F , we obtain a diagram in $K(\mathcal{B})$:

$$F(X^\bullet) \xrightarrow{F(t)} F(Y^\bullet) \xleftarrow{F(s)} F(I^\bullet).$$

As observed previously, $F(s)$ is also a quasi-isomorphism.

This induces a morphism in $D(\mathcal{B})$:

$$\xi(X^\bullet): Q \circ F(X^\bullet) \rightarrow Q \circ F(I^\bullet) = \overline{F} \circ Q(I^\bullet) = \overline{F} \circ U \circ Q(X^\bullet) = \underline{\underline{R}}^*F \circ Q(X^\bullet).$$

One can now check without difficulty that $\xi(X^\bullet)$ does not depend on the choice of the diagram (*) above, that ξ defines a functor morphism $Q \circ F \rightarrow \underline{\underline{R}}^*F \circ Q$, and that the pair $(\underline{\underline{R}}^*F, \xi)$ satisfies the universal property of the right derived functor.

If $X^\bullet \in \text{Ob } \mathcal{L}$, then $F(t)$ is also a quasi-isomorphism in the construction above, so $\xi(X^\bullet)$ is an isomorphism in $D(\mathcal{B})$, as required. $\square$

Proposition 11. *Let $\mathcal{A}, \mathcal{B}, K^*(\mathcal{A}), F$ be as above, and let $K^{**}(\mathcal{A}) \subseteq K^*(\mathcal{A})$ be another localizing subcategory of $K(\mathcal{A})$. Suppose there exists a triangulated subcategory $\mathcal{L} \subseteq K^*(\mathcal{A})$ satisfying hypotheses (1),(2) of Theorem 1.5.1, and such that $\mathcal{L} \cap K^{**}(\mathcal{A})$ satisfies hypothesis (1) for $K^{**}(\mathcal{A})$. Then $\underline{\underline{R}}^*F$ and $\underline{\underline{R}}^{**}(F|_{K^{**}(\mathcal{A})})$ both exist, and the natural morphism*

$$\underline{\underline{R}}^{**}(F|_{K^{**}(\mathcal{A})}) \rightarrow \underline{\underline{R}}^*F|_{D^{**}(\mathcal{A})}$$

is an isomorphism.

Proof. Existence follows immediately from Theorem 1.5.1. For the isomorphism, any object of $D^{**}(\mathcal{A})$ is isomorphic to one coming from an object of $\mathcal{L}$, we may assume that $X'' = Q(I^\bullet)$ for some $I^\bullet \in \text{Ob } \mathcal{L} \cap K^{**}(\mathcal{A})$. The claim then follows from the final assertion of Theorem 1.5.1. $\square$

Corollary 1. $\alpha)$ *Let $\mathcal{A}, \mathcal{B}$ be abelian categories, $F: K^+(\mathcal{A}) \rightarrow K(\mathcal{B})$ a ∂ -functor (e.g., induced by an additive functor $F_0: \mathcal{A} \rightarrow \mathcal{B}$). If $\mathcal{A}$ has enough injectives, then $\underline{\underline{R}}^+F$ exists.*

$\beta)$ *Let $\mathcal{A}, \mathcal{B}$ be abelian categories, $F: \mathcal{A} \rightarrow \mathcal{B}$ additive. Suppose there exists an subset $P \subseteq \text{Ob } \mathcal{A}$ satisfying (i),(ii) of Lemma 1.4.6, and:*

(iv) F sends short exact sequences in $\mathcal{A}$ to short exact sequences in $\mathcal{B}$.

Then $\underline{\underline{R}}^+F$ exists (we also denote by F its extension to a ∂ -functor $K^+(\mathcal{A}) \rightarrow K^+(\mathcal{B})$).

$\gamma)$ *Let $\mathcal{A}, \mathcal{B}$ be abelian categories, $F: \mathcal{A} \rightarrow \mathcal{B}$ additive. Assume:*

(a) The hypotheses of β hold;

(b) F has finite cohomological dimension: there exists $n > 0$ such that $R^i F(Y) = 0$ for all $Y \in \text{Ob } \mathcal{A}$ and all $i > n$.

Then $\underline{\underline{R}}F$ exists on all of $D(\mathcal{A})$, and its restriction to $D^+(\mathcal{A})$ coincides with $\underline{\underline{R}}^+F$.

Remark. Case α is a special case of β : if $\mathcal{A}$ has enough injectives, the class of injective objects satisfies (i),(ii),(iv) for any additive F .

Proof. $\alpha)$ Let $\mathcal{L} \subseteq K^+(\mathcal{A})$ be the triangulated subcategory of bounded-below injective complexes. By Lemma 1.4.6(1), every object of $K^+(\mathcal{A})$ admits a quasi-isomorphism into an object of $\mathcal{L}$. By Lemma 1.4.5, every quasi-isomorphism in $\mathcal{L}$ is an isomorphism, so acyclic objects of $\mathcal{L}$ are zero in $K(\mathcal{A})$. Thus condition (2) of Theorem 1.5.1 holds for any ∂ -functor F , so $\underline{\underline{R}}^+F$ exists.

$\beta)$ Take $\mathcal{L} \subseteq K^+(\mathcal{A})$ to be the triangulated subcategory of complexes of objects of P (Note $\mathcal{L}$ is a triangulated subcategory because P is stable under direct sums by

(ii). Condition (1) holds of the theorem is satisfied as above. For (2), let $Z^\bullet \in \mathcal{L}$ be acyclic; using (ii) repeatedly, all kernels $\ker d_Z^p$ lie in P . By (iv), F preserves exactness, so $F(Z^\bullet)$ is acyclic.

γ) Let P' be the collection of F -acyclic objects of $\mathcal{A}$ ($R^i F(X) = 0, \forall i > 0$). Then P' satisfies (i),(ii),(iii) of Lemma 1.4.6. Let $\mathcal{L} \subseteq K(\mathcal{A})$ be complexes of objects of P' . By Lemma 1.4.6 and the argument for β , the hypotheses of Theorem 1.5.1 are satisfied, so $\underline{\underline{R}}F$ exists.

By Proposition 1.5.2, the restriction of $\underline{\underline{R}}F$ to $D^+(\mathcal{A})$ is isomorphic to $\underline{\underline{R}}^+F$. $\square$

Proposition 12. *Let $\mathcal{A}, \mathcal{B}, \mathcal{C}$ be abelian categories, $K^*(\mathcal{A}) \subseteq K(\mathcal{A})$, $K^\dagger(\mathcal{B}) \subseteq K(\mathcal{B})$ localizing subcategories, and let*

$$F: K^*(\mathcal{A}) \rightarrow K(\mathcal{B}), \quad G: K^\dagger(\mathcal{B}) \rightarrow K(\mathcal{C})$$

be ∂ -functors.

a) *Assume $F(K^*(\mathcal{A})) \subseteq K^\dagger(\mathcal{B})$, that $\underline{\underline{R}}^*F, \underline{\underline{R}}^\dagger G, \underline{\underline{R}}^*(G \circ F)$ exist, and $\underline{\underline{R}}^*F(D^*(\mathcal{A})) \subseteq D^\dagger(\mathcal{B})$. Then there exists a unique functor morphism*

$$\zeta: \underline{\underline{R}}^*(G \circ F) \rightarrow \underline{\underline{R}}^\dagger G \circ \underline{\underline{R}}^*F$$

so the the diagram is commutative.

$$\begin{array}{ccc} Q \cdot G \cdot F & \xrightarrow{\xi_G} & \underline{\underline{R}}^+G \cdot Q \cdot F \\ \xi_{G \cdot F} \downarrow & & \downarrow \xi_F \\ \underline{\underline{R}}^*(G \cdot F) \cdot Q & \xrightarrow{\zeta \cdot Q} & \underline{\underline{R}}^+G \cdot \underline{\underline{R}}^*F \cdot Q \end{array}$$

b) *Suppose there exist triangulated subcategories $\mathcal{L} \subseteq K^*(\mathcal{A})$, $M \subseteq K^\dagger(\mathcal{B})$ satisfying Theorem 1.5.1's hypotheses for F, G respectively, and $F(\mathcal{L}) \subseteq M$. Then the hypotheses of (a) hold, and the canonical morphism ζ is an isomorphism.*

Proof. Straightforward $\square$

Remarks.

1. If F, G, H are three consecutive functors, then there is a commutative diagram of ζ s (provided they all exist):

$$\begin{array}{ccc}
 \underline{\underline{R}}(H \cdot G \cdot F) & \xrightarrow{\xi_{G \cdot F, H}} & \underline{\underline{R}}H \cdot \underline{\underline{R}}(G \cdot F) \\
 \downarrow \zeta_{F, H \cdot G} & & \downarrow \zeta_{F, G} \\
 \underline{\underline{R}}(H \cdot G) \cdot \underline{\underline{R}}F & \longrightarrow & \underline{\underline{R}}H \cdot \underline{\underline{R}}G \cdot \underline{\underline{R}}F
 \end{array}$$

2. This proposition shows the convenience of derived functors in the context of derived categories. What used to be a spectral sequence becomes now simply a composition of functors. (And of course one can recover the old spectral sequence from this proposition by taking cohomology and using the spectral sequence of a double complex.)

Corollary 2. *Left to the reader: Illustrate the Proposition in the style of Corollary 5.3.*

Proposition 13. *Let $\mathcal{A}$ be an abelian category, let $\mathcal{A}'$ be a thick abelian subcategory, let $\mathcal{B}$ be another abelian category, and let $F: K^+(\mathcal{A}) \rightarrow K^+(\mathcal{B})$ be a ∂ -functor. Suppose that $\underline{\underline{R}}^+F$ and $\underline{\underline{R}}^+(F|_{\mathcal{A}'})$ both exist. Then there is a natural morphism*

$$\zeta: \underline{\underline{R}}^+(F|_{\mathcal{A}'}) \longrightarrow \varphi \circ \underline{\underline{R}}^+F$$

of functors from $D^+(\mathcal{A}')$ to $D(\mathcal{B})$, where

$$\varphi: D^+(\mathcal{A}') \longrightarrow D^+(\mathcal{A})$$

is the natural functor. If furthermore $\mathcal{A}'$ has enough $\mathcal{A}$ -injectives, and $\mathcal{A}$ has enough injectives, then ζ is an isomorphism.

Proof. The existence of ζ follows from the definition of the derived functor. If $\mathcal{A}'$ has enough $\mathcal{A}$ -injectives, and $\mathcal{A}$ has enough injectives, then we can use $\mathcal{A}$ -injectives to calculate both functors above, by Corollary 1.5.3 α , and so ζ is an isomorphism. (Recall by Proposition 1.4.8 that φ is an equivalence of categories in that case.) $\square$

2.7 Examples. Ext and $\underline{\underline{R}}\text{Hom}$

Definition 11. *Let $\mathcal{A}$ be an abelian category, and let $X^\bullet, Y^\bullet \in D(\mathcal{A})$. Define the i th hyperext groups by*

$$\text{Ext}^i(X^\bullet, Y^\bullet) = \text{Hom}_{D(\mathcal{A})}(X^\bullet, T^i(Y^\bullet)).$$

Remarks.

1. If $X^\bullet, Y^\bullet \in D^+(\mathcal{A})$, then Ext^i computed in $D^+(\mathcal{A})$ agrees with the above, since $D^+(\mathcal{A})$ is a full subcategory of $D(\mathcal{A})$.
2. This gives in particular a definition of $\text{Ext}^i(X, Y)$ for any $X, Y \in \mathcal{A}$. We shall see that if $\mathcal{A}$ has enough injectives, this coincides with the classical definition of Ext groups.

Proposition 14. *Let*

$$0 \rightarrow X^\bullet \rightarrow Y^\bullet \rightarrow Z^\bullet \rightarrow 0$$

be a short exact sequence of complexes over $\mathcal{A}$, and let $V^\bullet$ be another complex. Then we have long exact sequences

$$\cdots \rightarrow \text{Ext}^i(V^\bullet, X^\bullet) \rightarrow \text{Ext}^i(V^\bullet, Y^\bullet) \rightarrow \text{Ext}^i(V^\bullet, Z^\bullet) \rightarrow \text{Ext}^{i+1}(V^\bullet, X^\bullet) \rightarrow \cdots$$

and

$$\cdots \rightarrow \text{Ext}^i(Z^\bullet, V^\bullet) \rightarrow \text{Ext}^i(Y^\bullet, V^\bullet) \rightarrow \text{Ext}^i(X^\bullet, V^\bullet) \rightarrow \text{Ext}^{i+1}(Z^\bullet, V^\bullet) \rightarrow \cdots$$

Proof. Let $W^\bullet$ be the third side of a triangle on $X^\bullet \rightarrow Y^\bullet$.

$$\begin{array}{ccccccc} & & & W^\bullet & & & \\ & & \swarrow & \uparrow & \searrow & g & \\ 0 & \longrightarrow & X^\bullet & \longrightarrow & Y^\bullet & \longrightarrow & Z^\bullet \longrightarrow 0 \end{array}$$

Then by Proposition 1.1(b), there is a morphism of complexes $g: W^\bullet \rightarrow Z^\bullet$. Using the long exact sequence of cohomology of the short exact sequence, and of the triangle, and using the five-lemma, we see that g is a quasi-isomorphism, i.e., an isomorphism in $D(\mathcal{A})$. Hence we may replace $Z^\bullet$ by $W^\bullet$ in the conclusion, which then follows from the same proposition. $\square$

Remark. It follows from the proof that any short exact sequence of complexes

$$0 \rightarrow X^\bullet \rightarrow Y^\bullet \rightarrow Z^\bullet \rightarrow 0$$

determines a morphism $Z^\bullet \rightarrow T(X^\bullet)$ in $D(\mathcal{A})$, making $X^\bullet, Y^\bullet, Z^\bullet$ into a triangle.

Now we will define a functor whose cohomology gives the Ext groups.

Definition 12. For complexes $X^\bullet, Y^\bullet$ over $\mathcal{A}$, define the complex $\text{Hom}^\bullet(X^\bullet, Y^\bullet)$ by

$$\text{Hom}^n(X^\bullet, Y^\bullet) = \prod_{p \in \mathbb{Z}} \text{Hom}_{\mathcal{A}}(X^p, Y^{p+n}),$$

with differential

$$d^n = \prod_{p \in \mathbb{Z}} (d_X^{p-1} + (-1)^{n+1} d_Y^{p+n}).$$

Notice under this definition the n -cycles of $\text{Hom}^\bullet(X^\bullet, Y^\bullet)$ correspond bijectively to morphisms $X^\bullet \rightarrow T^n(Y^\bullet)$ of complexes. The n -boundaries correspond to morphisms homotopic to zero. In other words

$$H^n(\text{Hom}^\bullet(X^\bullet, Y^\bullet)) \cong \text{Hom}_{K(\mathcal{A})}(X^\bullet, T^n(Y^\bullet)).$$

Clearly $\text{Hom}^\bullet$ is a bi- ∂ -functor

$$\text{Hom}^\bullet: K(\mathcal{A})^\circ \times K(\mathcal{A}) \rightarrow K(\mathbf{Ab}).$$

If $\mathcal{A}$ has enough injectives, we can calculate its derived functors.

Lemma 4. Let $X^\bullet \in K(\mathcal{A})$ be a complex, $Y^\bullet \in K^+(\mathcal{A})$ be a complex of injectives. Assume either

- (a) $Y^\bullet$ is acyclic, or
- (b) $X^\bullet$ is acyclic.

Then $\text{Hom}^\bullet(X^\bullet, Y^\bullet)$ is acyclic.

Proof. By the remark above, one has only to check that any morphism of $X^\bullet$ to $T^n(Y^\bullet)$ is homotopic to zero, or, since $T^n(Y^\bullet)$ also satisfies the hypotheses of the lemma, it is enough to show that any morphism of $X^\bullet$ to $Y^\bullet$ is homotopic to zero. In case (a), $Y^\bullet$ is split exact, and it is easy to construct the homotopy (left to reader). In case (b) the result follows from Lemma 1. $\square$

Suppose $\mathcal{A}$ has enough injectives, and let $\mathcal{L} \subseteq K^+(\mathcal{A})$ be the triangulated subcategory of bounded-below injective complexes. By Lemma 1.6.2(a), for fixed $X^\bullet \in K(\mathcal{A})$, the functor

$$\text{Hom}^\bullet(X^\bullet, \cdot): K^+(\mathcal{A}) \rightarrow K(\mathbf{Ab})$$

satisfies the hypotheses of Theorem 1.5.1, hence admits a right derived functor. This is functorial in $X^\bullet$, giving a bi- ∂ -functor

$$\underline{\underline{R}}_{\text{II}} \text{Hom}^\bullet: K(\mathcal{A})^\circ \times D^+(\mathcal{A}) \rightarrow D(\mathbf{Ab}).$$

Now using the lemma, part b), we see that this functor is "exact in the first variable, i.e., takes acyclic complexes into acyclic complexes, and hence passes to the quotient, giving a trivial derived functor

$$\underline{R}_I \underline{R}_{II} \text{Hom}^\bullet : D(A)^\circ \times D^+(A) \longrightarrow D(\mathbf{Ab}).$$

(We will denote this functor by $\underline{R} \text{Hom}^\bullet$ when no confusion can

Suppose on the other hand that A has enough projectives by the usual process of "reversing the arrows" we see that the also a functor

$$\underline{R}_{II} \underline{R}_I \text{Hom}^\bullet : D^-(A)^\circ \times D(A) \longrightarrow D(A).$$

Now if A has enough injectives and enough projectives, then functors $\underline{R}_I \underline{R}_{II} \text{Hom}^\bullet$ and $\underline{R}_{II} \underline{R}_I \text{Hom}^\bullet$ are defined on $D^-(A)^\circ \times D^+$

and are canonically isomorphic, as we see by the lemma below. Thus we are justified in using the ambiguous notation $\underline{R} \text{Hom}^\bullet$.

Lemma 5. *Let $\mathcal{A}, \mathcal{B}, \mathcal{C}$ be abelian categories,*

$$T : K^*(\mathcal{A}) \times K^+(\mathcal{B}) \rightarrow K(\mathcal{C})$$

a bi- ∂ -functor. Suppose that the derived functors $\underline{R}_I \underline{R}_{II} T$ and $\underline{R}_{II} \underline{R}_I T$ both exist, (where the subscripts I, II denote the derived functor in the first or second variable, respectively). Then there is a unique isomorphism between them compatible with the morphisms $\xi_1 : T \rightarrow \underline{R}_I \underline{R}_{II} T$ and $\xi_2 : T \rightarrow \underline{R}_{II} \underline{R}_I T$

Proof. Immediate from the universal property of derived functors. $\square$

Theorem 2 (Yoneda). *Let $\mathcal{A}$ have enough injectives. For all $X^\bullet \in D(\mathcal{A})$, $Y^\bullet \in D^+(\mathcal{A})$,*

$$H^i(\underline{R} \text{Hom}^\bullet(X^\bullet, Y^\bullet)) = \text{Ext}^i(X^\bullet, Y^\bullet).$$

Corollary 3. *If $\mathcal{A}$ has enough injectives, the derived-category definition of $\text{Ext}^i(X, Y)$ coincides with the classical Ext groups for all $X, Y \in \mathcal{A}$.*

Proof of Corollary 1.6.5. Take an injective resolution $I^\bullet$ of Y . Then

$$\text{Ext}^i(X, Y) = H^i(\text{Hom}^\bullet(X, I^\bullet)),$$

which is the standard definition. $\square$

Proof of Theorem 1.6.4. Choose a quasi-isomorphism $s: Y^\bullet \rightarrow I^\bullet$ with $I^\bullet$ a bounded-below injective complex. Then

$$\mathrm{Ext}^i(X^\bullet, Y^\bullet) = \mathrm{Ext}^i(X^\bullet, I^\bullet) = \mathrm{Hom}_{D(\mathcal{A})}(X^\bullet, T^i(I^\bullet)).$$

By Lemma 1.4.5, one sees that every morphism in $D(\mathcal{A})$ of a complex $X^\bullet$ to a complex of injectives bounded below, say $T^i(I^\bullet)$, is represented by an actual morphism of complexes. Hence the above is equal to

$$\begin{aligned} \mathrm{Hom}_{D(\mathcal{A})}(X^\bullet, T^i(I^\bullet)) &= \mathrm{Hom}_{K(\mathcal{A})}(X^\bullet, T^i(I^\bullet)) \\ &= H^i(\mathrm{Hom}^\bullet(X^\bullet, I^\bullet)) \\ &= H^i(\underline{\mathrm{R}}\mathrm{Hom}^\bullet(X^\bullet, Y^\bullet)). \end{aligned}$$

□

2.8 Way-out functors and Isomorphisms

Definition 13. Let $\mathcal{A}, \mathcal{B}$ be abelian categories, and let $F: D(\mathcal{A}) \rightarrow D(\mathcal{B})$ be a (co-)variant ∂ -functor. We say F is *way-out (right)* if for every $n_1 \in \mathbb{Z}$, there exists $n_2 \in \mathbb{Z}$ such that for any complex $X^\bullet \in D(\mathcal{A})$ satisfying $H^i(X^\bullet) = 0$ for all $i < n_2$, we have $H^i(F(X^\bullet)) = 0$ for all $i < n_1$.

One defines *way-out left* and *way-out in both directions* similarly. For a contravariant ∂ -functor $F: D(\mathcal{A}) \rightarrow D(\mathcal{B})$, the definition of *way-out right* is the same, except reversing the inequality to $i > n_2$.

Examples.

1. If $F_0: \mathcal{A} \rightarrow \mathcal{B}$ is additive and satisfies the hypotheses of Corollary 1.5.3(α) or (β), then $\underline{\mathrm{R}}^+F$ is way-out right. If F_0 satisfies (γ), then $\underline{\mathrm{R}}F$ is way-out in both directions.
2. For unbounded $X^\bullet \in D(\mathcal{A})$, the functor $\underline{\mathrm{R}}\mathrm{Hom}(X^\bullet, Y^\bullet)$ for $Y^\bullet \in \mathrm{Ob} D^+(\mathcal{A})$ is generally not way-out in $Y^\bullet$.

Proposition 15 (Lemma on Way-out Functors). Let $\mathcal{A}, \mathcal{B}$ be abelian categories, $\mathcal{A}' \subseteq \mathcal{A}$ a thick subcategory. Let F, G be ∂ -functors from $D_{\mathcal{A}'}^+(\mathcal{A})$ or $D_{\mathcal{A}'}(\mathcal{A})$ to $D(\mathcal{B})$, and $\eta: F \rightarrow G$ a morphism of functors.

- (i) If $\eta(X)$ is an isomorphism for all $X \in \mathrm{Ob} \mathcal{A}'$, then $\eta(X^\bullet)$ is an isomorphism for all $X^\bullet \in D_{\mathcal{A}'}^b(\mathcal{A})$.

- (ii) If $\eta(X)$ is an isomorphism for all $X \in \text{Ob } \mathcal{A}'$, and F, G are way-out right, then $\eta(X^\bullet)$ is an isomorphism for all $X^\bullet \in D_{\mathcal{A}'}^+(\mathcal{A})$.
- (iii) If $\eta(X)$ is an isomorphism for all $X \in \text{Ob } \mathcal{A}'$, and F, G are way-out in both directions, then $\eta(X^\bullet)$ is an isomorphism for all $X^\bullet \in D_{\mathcal{A}'}(\mathcal{A})$.
- (iv) Let $P \subseteq \text{Ob } \mathcal{A}'$ be such that every object of $\mathcal{A}'$ embeds into an object of P . If $\eta(X)$ is an isomorphism for all $X \in P$, and F, G are way-out right, then $\eta(X)$ is an isomorphism for all $X \in \text{Ob } \mathcal{A}'$.

Definition 14. Let $X^\bullet$ be a complex over abelian category $\mathcal{A}$, $n \in \mathbb{Z}$. Define truncation functors:

$$\begin{aligned} \tau_{>n}(X^\bullet) &: \cdots \rightarrow 0 \rightarrow X^{n+1} \rightarrow X^{n+2} \rightarrow \cdots, \\ \tau_{\leq n}(X^\bullet) &: \cdots \rightarrow X^{n-1} \rightarrow X^n \rightarrow 0 \rightarrow \cdots, \\ \sigma_{>n}(X^\bullet) &: \cdots \rightarrow 0 \rightarrow \text{im } d^n \rightarrow X^{n+1} \rightarrow X^{n+2} \rightarrow \cdots, \\ \sigma_{\leq n}(X^\bullet) &: \cdots \rightarrow X^{n-1} \rightarrow \ker d^n \rightarrow 0 \rightarrow \cdots. \end{aligned}$$

There are natural morphisms inducing short exact sequences of complexes:

$$0 \rightarrow \tau_{>n}(X^\bullet) \rightarrow X^\bullet \rightarrow \tau_{\leq n}(X^\bullet) \rightarrow 0, \quad (2.1)$$

$$0 \rightarrow \sigma_{\leq n}(X^\bullet) \rightarrow X^\bullet \rightarrow \sigma_{>n}(X^\bullet) \rightarrow 0. \quad (2.2)$$

The σ -truncations satisfy:

$$H^i(\sigma_{>n}(X^\bullet)) = \begin{cases} H^i(X^\bullet) & i > n, \\ 0 & i \leq n; \end{cases} \quad H^i(\sigma_{\leq n}(X^\bullet)) = \begin{cases} H^i(X^\bullet) & i \leq n, \\ 0 & i > n. \end{cases}$$

Lemma 6. For any complex $X^\bullet$ over $\mathcal{A}$, there exist triangles in $D(\mathcal{A})$:

(3)

$$\begin{array}{ccc} & \tau_{>n}(X^\bullet) & \\ \swarrow & & \searrow \\ \tau_{\leq n}(X^\bullet) & \xrightarrow{\quad} & X^n \end{array}$$

(4)

$$\begin{array}{ccc} & \sigma_{>n}(X^\bullet) & \\ \swarrow & & \searrow \\ H^n(X^\bullet) & \xrightarrow{\quad} & \sigma_{\geq n}(X^\bullet) \end{array}$$

Proof. Triangle (3) comes from the short exact sequence

$$0 \rightarrow \tau_{>n}(X^\bullet) \rightarrow \tau_{\geq n}(X^\bullet) \rightarrow X^n \rightarrow 0.$$

For the second, define $\sigma'_{\geq n}(X^\bullet)$ by

$$\dots \rightarrow 0 \rightarrow X^n / \text{im } d^{n-1} \rightarrow X^{n+1} \rightarrow \dots.$$

There is a quasi-isomorphism $\sigma_{\geq n}(X^\bullet) \rightarrow \sigma'_{\geq n}(X^\bullet)$, and an exact sequence

$$0 \rightarrow H^n(X^\bullet) \rightarrow \sigma'_{\geq n}(X^\bullet) \rightarrow \sigma_{>n}(X^\bullet) \rightarrow 0.$$

This induces the required triangle in $D(\mathcal{A})$. $\square$

Proof of Proposition 1.7.1. (i) Let $X^\bullet \in \text{Ob } D_{\mathcal{A}'}^b(\mathcal{A})$. We prove, by descending induction on n , that

$$\eta(\sigma_{>n}(X^\bullet)) : F(\sigma_{>n}(X^\bullet)) \rightarrow G(\sigma_{>n}(X^\bullet))$$

is an isomorphism for all n . If n is large enough, then $\sigma_{>n}(X^\bullet)$ has zero cohomology, since $X^\bullet$ has bounded cohomology. Hence it is the zero object of $D(\mathcal{A})$, and η of it is an isomorphism. The induction step follows from the hypotheses and Proposition 1.1(c), applied to the triangle (4) above. Now for n small enough, the natural map $X^\bullet \rightarrow \sigma_{>n}(X^\bullet)$ is a quasi-isomorphism, hence an isomorphism in $D(\mathcal{A})$, and we are done.

(ii) Let $X^\bullet \in \text{Ob } D_{\mathcal{A}'}^+(\mathcal{A})$. To show $\eta(X^\bullet)$ is an isomorphism, it is sufficient to show that

$$H^j(\eta(X^\bullet)) : H^j(F(X^\bullet)) \rightarrow H^j(G(X^\bullet))$$

is an isomorphism for all $j \in \mathbb{Z}$. Given j , let $n_1 \geq j + 2$, and choose n_2 as in the definition of way-out functors, to work for F and G both. From the exact sequence (2) above, we have a triangle in $D(\mathcal{A})$.

$$\begin{array}{ccc} & \tau_{>n_2}(X^\bullet) & \\ \swarrow & & \searrow \\ \tau_{\geq n_2}(X^\bullet) & \xrightarrow{\quad} & X^\bullet \end{array}$$

Since

$$H^i(\sigma_{>n_2}(X^\bullet)) = 0 \quad \text{for } i \leq n_2,$$

we have

$$H^i(F(\sigma_{>n_2}(X^\bullet))) = 0 = H^i(G(\sigma_{>n_2}(X^\bullet)))$$

for $i < n_1$, in particular for $i = j, j + 1$. Therefore from the long exact sequence of cohomology we get isomorphisms

$$\begin{aligned} H^j(F(\sigma_{\leq n_2}(X^\bullet))) &\xrightarrow{\sim} H^j(F(X^\bullet)), \\ H^j(G(\sigma_{\leq n_2}(X^\bullet))) &\xrightarrow{\sim} H^j(G(X^\bullet)). \end{aligned}$$

But $\sigma_{\leq n_2}(X^\bullet) \in D_{\mathcal{A}'}^b(\mathcal{A})$, so η is an isomorphism on it by what we have just proved. Hence $H^j(\eta(X^\bullet))$ is an isomorphism, as required.

(iii) Given $X^\bullet \in D_{\mathcal{A}'}(\mathcal{A})$, we treat first $\sigma_{\leq 0}(X^\bullet)$ and $\sigma_{> 0}(X^\bullet)$ by case (ii), then glue the results by a triangle deduced from the exact sequence (2).

(iv) Given $X \in \text{Ob } \mathcal{A}'$, we can find by Lemma 1.4.6 a resolution $I^\bullet \in D_{\mathcal{A}'}^+(\mathcal{A})$ of X , with each $I^p \in P$. Thus it is sufficient to show that $\eta(I^\bullet)$ is an isomorphism. This follows by applying the technique of (i) and (ii), but using the functors τ instead of σ . $\square$

Proposition 16. *Let $\mathcal{A}, \mathcal{B}$ be abelian categories, and let $\mathcal{A}' \subseteq \mathcal{A}, \mathcal{B}' \subseteq \mathcal{B}$ be thick abelian subcategories, F be a ∂ -functor from $D_{\mathcal{A}'}^+(\mathcal{A})$ or $D_{\mathcal{A}'}(\mathcal{A})$ to $D(\mathcal{B})$,.*

- (i) *If $F(X) \in D_{\mathcal{B}'}(\mathcal{B})$ for all $X \in \text{Ob } \mathcal{A}'$, then $F(X^\bullet) \in D_{\mathcal{B}'}(\mathcal{B})$ for all $X^\bullet \in D_{\mathcal{A}'}^b(\mathcal{A})$.*
- (ii) *If $F(X) \in D_{\mathcal{B}'}(\mathcal{B})$ for all $X \in \text{Ob } \mathcal{A}'$ and F is way-out right, then $F(X^\bullet) \in D_{\mathcal{B}'}(\mathcal{B})$ for all $X^\bullet \in D_{\mathcal{A}'}^+(\mathcal{A})$.*
- (iii) *As in (ii) if F is way-out in both directions, then $F(X^\bullet) \in D_{\mathcal{B}'}(\mathcal{B})$ for all $X^\bullet \in D_{\mathcal{A}'}(\mathcal{A})$.*
- (iv) *Let $P \subseteq \text{Ob } \mathcal{A}'$ such that every object of $\mathcal{A}'$ embeds into P . If $F(X) \in D_{\mathcal{B}'}(\mathcal{B})$ for all $X \in P$ and F is way-out right, then $F(X) \in D_{\mathcal{B}'}(\mathcal{B})$ for all $X \in \text{Ob } \mathcal{A}'$.*

Proof. The proof is analogous to that of the previous proposition, and will be left to the reader. $\square$

Proposition 17. *Let $\mathcal{A}$ have enough injectives, $F: \mathcal{A} \rightarrow \mathcal{B}$ additive of cohomological dimension $\leq n$. Let $P \subseteq \text{Ob } \mathcal{A}$ be all X with $R^i F(X) = 0$ for $i \neq n$. Assume every object of $\mathcal{A}$ is a quotient of an object in P . Set $G = R^n F$. Then $\underline{R}F, \underline{L}G$ exist, and there is a functorial isomorphism*

$$\psi: \underline{R}F \xrightarrow{\sim} \underline{L}G[-n],$$

where $[-n]$ denotes shifting right by n degrees.

Proof. $\underline{\underline{R}}F$ exists by Corollary 1. To see that $\underline{\underline{L}}G$ exists, we apply Theorem 1 with the arrows reversed. By hypothesis P satisfies condition (i)* of Lemma 3 (where $*$ denotes reversal of arrows), and one sees easily from the definition of P that it satisfies also (ii)*, (iii)*, and (iv)* of Corollary 1. Therefore, if we let $\mathcal{L} \subseteq K(\mathcal{A})$ be the triangulated subcategory consisting of complexes of elements of P , we see as in the proof of Corollary 1(γ) that the hypotheses 1)* and 2)* = 2) of the theorem are satisfied, and so $\underline{\underline{L}}G$ exists.

To define the morphism ψ , we observe, as in the proof of Theorem 1, that $\mathcal{L}_{\text{Qis}} \rightarrow D(\mathcal{A})$ is an equivalence of categories, so it will be sufficient to define ψ on $\mathcal{L}_{\text{Qis}}$. Or, using Proposition 5, it will be sufficient to define ψ on $\mathcal{L}$ itself. In other words, for every complex $X^\bullet$ of objects of P we must give a morphism in $D(\mathcal{B})$,

$$\psi(X^\bullet): \underline{\underline{R}}F(X^\bullet) \rightarrow \underline{\underline{L}}G(X^\bullet)[-n],$$

such that whenever $f: X^\bullet \rightarrow Y^\bullet$ is a morphism of such complexes, there is the usual commutative diagram $\underline{\underline{L}}G(f) \cdot \psi(X^\bullet) = \psi(Y^\bullet) \cdot \underline{\underline{R}}F(f)$.

Definition 15. Let $\mathcal{A}$ be an abelian category, $X^\bullet$ a complex of objects of $\mathcal{A}$. We refer to [M, Ch. IV, §4] for the definition of double complexes, maps and homotopies of double complexes, and injective resolutions of complexes. We recall the following facts. A (right) Cartan–Eilenberg resolution of $X^\bullet$ is a double complex $C^\bullet$ of injective objects of $\mathcal{A}$, with $C^{p,q} = 0$ if $p < 0$, together with an augmentation morphism $\epsilon: X^\bullet \rightarrow C^{\bullet,0}$ such that for every $p \in \mathbb{Z}$, the induced maps

$$B^p(\epsilon): B^p(X^\bullet) \rightarrow B_1^p(C^\bullet), \quad H^p(\epsilon): H^p(X^\bullet) \rightarrow H_1^p(C^\bullet)$$

are injective resolutions in the usual sense (cf. [M, Ch. XVII, §1] where this is called an injective resolution of $X^\bullet$).

Lemma 7.

- a) If $\mathcal{A}$ has enough injectives, every complex $X^\bullet$ of objects of $\mathcal{A}$ has a Cartan–Eilenberg resolution.
- b) If $f: X^\bullet \rightarrow Y^\bullet$ is a morphism of complexes, and $C^\bullet, D^\bullet$ are Cartan–Eilenberg resolutions of $X^\bullet, Y^\bullet$, then there exists a map $F: C^\bullet \rightarrow D^\bullet$ of double complexes lying over f .
- c) If $f, g: X^\bullet \rightarrow Y^\bullet$ are homotopic maps, and $F, G: C^\bullet \rightarrow D^\bullet$ are maps of Cartan–Eilenberg resolutions lying over f, g , then F is homotopic to G .

- d) If $F, G: C^\bullet \rightarrow D^\bullet$ are homotopic maps of double complexes, then $s(F), s(G): s(C^\bullet) \rightarrow s(D^\bullet)$ are homotopic maps of the associated simple complexes.
- e) If $F, G: C^\bullet \rightarrow D^\bullet$ are homotopic maps of double complexes, and $\sigma_{\leq n}^\Pi$ denotes truncation with respect to the second differential d_2 , then the restrictions of F and G to maps $\sigma_{\leq n}^\Pi(C^\bullet) \rightarrow \sigma_{\leq n}^\Pi(D^\bullet)$ are homotopic.
- f) If $\epsilon: X^\bullet \rightarrow C^\bullet$ augments $X^\bullet$ to a double complex $C^\bullet$ with $C^{p,q} = 0$ for $q < 0$ and $q > n_0$ (for suitable n_0), and such that for each p the map $X^p \rightarrow C^{p,\bullet}$ is a resolution, then the natural map $X^\bullet \rightarrow s(C^\bullet)$ into the associated simple complex is a quasi-isomorphism.

Proof of Lemma 1.7.5. Parts a), b), and c) are in [M, Ch. XVII, Prop. 1.2]. Part d) is in [M, Ch. IV, §4]. Part e) is easy. Part f) follows either from the spectral sequence of a double complex [EGA 0_{III} 11.3.3(ii)] or by a direct calculation. $\square$

Proof of Proposition 1.7.4, continued. Given $X^\bullet \in \text{Ob } \mathcal{L}$, choose a Cartan–Eilenberg resolution $C^\bullet$ of $X^\bullet$ (exists by Lemma 1.7.5 a)). Define $C'^\bullet = \sigma_{\leq n}^\Pi(C^\bullet)$; explicitly

$$C'^{p,q} = \begin{cases} C^{p,q} & q < n, \\ \ker d_2^{p,n} & q = n, \\ 0 & q > n. \end{cases}$$

For each $p \in \mathbb{Z}$, we have an exact sequence

$$0 \rightarrow X^p \rightarrow C'^{p,0} \rightarrow C'^{p,1} \rightarrow \dots \rightarrow C'^{p,n-1} \rightarrow C'^{p,n} \rightarrow 0.$$

The terms $C'^{p,0}, \dots, C'^{p,n-1}$ are injective and $X^p \in P$ implies $C'^{p,n}$ is F -acyclic. Hence this resolution may be used to calculate $\underline{\underline{R}}F(X^p)$ (see end of Theorem 1.5.1), giving an isomorphism

$$\varphi(t^p): G(X^p) = R^n F(X^p) \xrightarrow{\sim} F(C'^{p,n}) / \text{im } F(C'^{p,n-1}),$$

where t^p is the quasi-isomorphism $X^p \rightarrow C^{p,\bullet}$. This may be used to construct a map

$$\alpha^p: F(C^{p,\bullet}) \rightarrow G(X^p),$$

whence a map of double complexes

$$\alpha: F(C'^\bullet) \rightarrow G(X^\bullet),$$

where $G(X^\bullet)$ is concentrated in the n -th row.

Passing to the associated simple complexes we obtain

$$s(\alpha): F(s(C'^{\bullet})) \rightarrow G(X^{\bullet})[-n].$$

Now $u: X^{\bullet} \rightarrow s(C'^{\bullet})$ is a quasi-isomorphism into a complex of F -acyclic objects, by Lemma 1.7.5(f), so there is an isomorphism

$$\varphi(u): \underline{\underline{R}}F(X^{\bullet}) \xrightarrow{\sim} F(s(C'^{\bullet})).$$

Since $X^{\bullet}$ consists of G -acyclic objects, there is an isomorphism

$$\varphi(\text{id}_{X^{\bullet}}): G(X^{\bullet}) \xrightarrow{\sim} \underline{\underline{L}}G(X^{\bullet}).$$

Composing, we define

$$\psi(X^{\bullet}): \underline{\underline{R}}F(X^{\bullet}) \longrightarrow \underline{\underline{L}}G(X^{\bullet})[-n].$$

I do not care whether $\psi(X^{\bullet})$ is dependent of the choice of the Cartan–Eilenberg resolution $C^{\bullet}$. It will be sufficient to verify that whenever $f: X^{\bullet} \rightarrow Y^{\bullet}$ is a morphism of complexes in $\mathcal{L}$, then $\psi(X^{\bullet})$ and $\psi(Y^{\bullet})$ defined as above, fit into a commutative diagram. This is not hard to show using the results of the lemma above, and can safely be left to the reader.

Thus we have a morphism of functors $\psi: \underline{\underline{R}}F \rightarrow \underline{\underline{L}}G[-n]$.

To show that ψ is an isomorphism, we note that $\underline{\underline{R}}F$ and $\underline{\underline{L}}G$ are way-out in both directions, and so, by the lemma on way-out functors (Proposition 15), we reduce to showing $\psi(X)$ is an isomorphism for any $X \in P$. But that is clear from the construction. $\square$

Proposition 18. *Let $\mathcal{A}$ have enough injectives, $X^{\bullet} \in K^+(\mathcal{A})$. The following are equivalent:*

- (i) $X^{\bullet}$ admits a quasi-isomorphism $X^{\bullet} \rightarrow I^{\bullet}$ into a bounded complex of injective objects of $\mathcal{A}$.
- (ii) The functor $\underline{\underline{R}}\text{Hom}(\cdot, X^{\bullet}): D(\mathcal{A})^{\circ} \rightarrow D(\mathbf{Ab})$ is way-out left (hence way-out both ways).
- (iii) There exists $n_0 \in \mathbb{Z}$ such that $\text{Ext}^i(Y, X^{\bullet}) = 0$ for all $Y \in \text{Ob } \mathcal{A}$ and all $i > n_0$.

Proof. (i) $\Rightarrow$ (ii): We may assume that $X^{\bullet}$ is a bounded complex of injectives, and calculate $F(Y^{\bullet})$ as $\text{Hom}(Y^{\bullet}, X^{\bullet})$. Then it is clear that F is way-out in both directions.

(ii) $\Rightarrow$ (iii): Since F is way-out left, choose n_0 such that whenever $Y^\bullet \in D(\mathcal{A})$ and $H^i(Y^\bullet) = 0$ for $i < n_0$, then $H^i(F(Y^\bullet)) = 0$ for $i > 0$. Then this n_0 will do. Indeed, let $Y \in \text{Ob } \mathcal{A}$, and let $Y'^\bullet$ be the complex $T^{-n_0}(Y)$. Then $H^i(Y'^\bullet) = 0$ for $i < n_0$, so $H^i(F(Y'^\bullet)) = 0$ for $i > 0$. But $H^i(F(Y'^\bullet)) = H^i(F(T^{-n_0}Y)) = H^{i+n_0}(F(Y))$. This says $\text{Ext}^j(Y, X^\bullet) = 0$ for $j > n_0$, as required.

(iii) $\Rightarrow$ (i): Let $X^\bullet$ satisfy condition (iii) for n_0 . Since $\mathcal{A}$ has enough injectives, we can find an injective resolution of $X^\bullet$ (Lemma 3), i.e., a quasi-isomorphism $s: X^\bullet \rightarrow X'^\bullet$ where $X'^\bullet$ is a bounded-below complex of injective objects of $\mathcal{A}$.

I claim that $H^i(X'^\bullet) = 0$ for all $i > n_0$. Suppose to the contrary that $H^m(X'^\bullet) \neq 0$ for some $m > n_0$. Then from the exact sequence

$$0 \rightarrow B^m(X'^\bullet) \rightarrow Z^m(X'^\bullet) \rightarrow H^m(X'^\bullet) \rightarrow 0$$

we see that the first inclusion is strict. Hence there exists a $Y \in \text{Ob } \mathcal{A}$ (e.g., $Y = Z^m(X'^\bullet)$) such that the inclusion

$$\text{Hom}(Y, B^m(X'^\bullet)) \rightarrow \text{Hom}(Y, Z^m(X'^\bullet))$$

is strict. However, there is a commutative diagram

$$\begin{array}{ccc} \text{Hom}(Y, B^m(X'^\bullet)) & \longrightarrow & \text{Hom}(Y, Z^m(X'^\bullet)) \\ \downarrow & & \downarrow \cong \\ B^m(\text{Hom}^\bullet(Y, X'^\bullet)) & \longrightarrow & Z^m(\text{Hom}^\bullet(Y, X'^\bullet)) \end{array}$$

with an isomorphism on the right since Hom is left exact. But by hypothesis $\text{Ext}^m(Y, X^\bullet) = H^m(\text{Hom}^\bullet(Y, X'^\bullet)) = 0$, so the arrow on top is an isomorphism, which gives a contradiction. Hence $H^i(X'^\bullet) = 0$ for $i > n_0$, as claimed.

It follows that the map

$$i: \sigma_{\leq n}(X'^\bullet) \rightarrow X'^\bullet$$

is a quasi-isomorphism for any $n \geq n_0$ (using the notation introduced earlier in this section). Now I claim that $Z^{n+1}(X'^\bullet) = B^{n+1}(X'^\bullet)$ is injective for any $n \geq n_0$. This will show that $\sigma_{\leq n+1}(X'^\bullet)$ is a bounded complex of injectives.

Consider the exact sequence of complexes

$$0 \rightarrow \sigma_{\leq n}(X'^\bullet) \rightarrow \tau_{\leq n}(X'^\bullet) \rightarrow B^{n+1}(X'^\bullet)[-n] \rightarrow 0$$

for $n \geq n_0$. We will show that $B^{n+1} = B^{n+1}(X'^\bullet)$ is injective by showing that $\text{Ext}^1(Y, B^{n+1}) = 0$ for all $Y \in \text{Ob } \mathcal{A}$. For any Y , we have

$$\text{Ext}^1(Y, B^{n+1}) = \text{Ext}^{n+1}(Y, B^{n+1}[-n])$$

and there is an exact sequence

$$\cdots \rightarrow \operatorname{Ext}^{n+1}(Y, \tau_{\leq n}(X'^{\bullet})) \rightarrow \operatorname{Ext}^{n+1}(Y, B^{n+1}[-n]) \rightarrow \operatorname{Ext}^{n+2}(Y, \sigma_{\leq n}(X'^{\bullet})) \rightarrow \cdots$$

The group on the left is zero since $\tau_{\leq n}(X'^{\bullet})$ is an injective complex which stops in degree n . The group on the right is zero by our hypothesis and the fact that $\sigma_{\leq n}(X'^{\bullet})$ is isomorphic to $X'^{\bullet}$ in $D^+(\mathcal{A})$. Hence the middle group is zero.

Now for $n \geq n_0$ we have seen that $\sigma_{\leq n}(X'^{\bullet})$ is a bounded complex of injectives, and that the map

$$i: \sigma_{\leq n}(X'^{\bullet}) \rightarrow X'^{\bullet}$$

is a quasi-isomorphism. By Lemma 2, i has a homotopy inverse, say f . Therefore the composition

$$fs: X^{\bullet} \rightarrow \sigma_{\leq n}(X'^{\bullet})$$

gives the required quasi-isomorphism of $X^{\bullet}$ into a bounded complex of injectives. $\square$

Definition 16. *If $X^{\bullet} \in K^+(\mathcal{A})$ satisfies the equivalent conditions of Proposition 1.7.6, we say $X^{\bullet}$ has finite injective dimension.*

Corollary 4. *The complexes of finite injective dimension form a triangulated localizing subcategory of $K(\mathcal{A})$.*

Proof. The fact that they form a triangulated subcategory follows e.g. from condition (iii) of Proposition 18 and the long exact sequence of Ext . We see that the subcategory is localizing by applying Proposition 4(i), since anything in $K^+(\mathcal{A})$ quasi-isomorphic to an element of the subcategory is already in the subcategory, and $K^+(\mathcal{A})$ is a localizing subcategory of $K(\mathcal{A})$. $\square$

Definition 17. *Denote by $K^+(\mathcal{A})_{\text{fid}}$ the full subcategory of $K^+(\mathcal{A})$ consisting of complexes of finite injective dimension. The corresponding derived subcategory is $D^+(\mathcal{A})_{\text{fid}}$.*

Remark. These subcategories are essentially different from the subcategories of the form $K_{\mathcal{A}'}(\mathcal{A})$ and $D_{\mathcal{A}'}(\mathcal{A})$ we have considered before, because they cannot be characterized by putting conditions on the cohomology objects $H^i(X^{\bullet})$.

Chapter 3

Application to Preschemes

3.1 Categories of sheaves

3.2 The derived functors of f_* and Γ

3.3 The derived functor of $\mathcal{H}om$

3.4 The derived functors of $\otimes$ and f^*

3.5 Relations among the derived functors

3.6 Compatibilities among the relations of §3.5

3.7 Injective sheaves on a locally noetherian prescheme

Chapter 4

Duality for Projective Morphisms

4.1 Differentials

4.2 $f^\sharp$ for a smooth morphism f

4.3 Recall of the explicit calculations

4.4 The trace map for projective space

4.5 The duality theorem for projective space

4.6 Duality for a finite morphism

4.7 The fundamental local isomorphism

4.8 $f^!$ for embeddable morphisms

4.9 The residue symbol

4.10 Trace for projective morphisms

4.11 Duality for projective morphisms

Chapter 5

Local Cohomology

5.1 Local cohomology groups, sheaves, and complexes

5.2 Depth and the Cousin complex

5.3 Generalization to complexes

Chapter 6

Dualizing Complexes and Local Duality

6.1 Introduction

6.2 Example: duality for abelian groups

6.3 Dualizing complexes

6.4 Uniqueness of the dualizing complex

6.5 Local cohomology on a prescheme

6.6 Dualizing functors on a local noetherian ring

6.7 Local duality

6.8 Application to dualizing complexes

6.9 Pointwise dualizing complexes and $f^\sharp$

6.10 Gorenstein preschemes

6.11 Existence of dualizing complexes

Chapter 7

Residual Complexes

7.1 Introduction

7.2 Functorial properties

7.3 $f^!$ for residual complexes

7.4 Trace for residual complexes

7.5 Behavior with respect to certain base changes

Chapter 8

The Duality Theorem

8.1 Curves over an Artin ring

8.2 The residue theorem

8.3 The duality theorem for proper morphisms

8.4 Smooth morphisms

Index of Definitions

acyclic complex	1.5
bounded complex	1.2
Cartan-Eilenberg resolution	1.7
catenary	V.7
Cech complex	III.3
codimension	IV.1
codimension function	V.7
co-finite type module	V.5
Cohen-Macaulay complex	IV.3
Cohen-Macaulay morphism	V.9
Cohen-Macaulay-1 ring	V.9
Cohen-Macaulay sheaf	IV.2
cohomological functor	I.1
complex	1.2
Cousin complex	IV.3
Cousin complex of $\mathcal{F}$	IV.2
derived category	I.1
∂ -functor	I.4
depth	IV.2
derived functor	1.5
duality morphism	III.5
dualizing complex	III.8
dualizing functor	V.2
embeddable morphism	V.5
$\mathcal{F}$ -acyclic	1.5
family of supports	IV.1
filtration associated to a dualizing complex	V.7
finite cohomological dimension	1.5

finite injective dimension	IV.3
finite Tor-dimension	1.7
Gorenstein complex	II.4
Gorenstein prescheme	III.1, V.9
Gorenstein morphism	V.9
Gorenstein ring	V.9
homotopy	1.6
hyperext	III.7
Koszul complex	1.2
Krull dimension	V.2
local complete intersection	III.1
local hyperext	II.3
local hyper Tor	II.3
localization of a category	1.3
localizing subcategory	II.7
locally noetherian category	1.5
local parameter	VII.1
mapping cone	1.2
multiplicative system	1.3
noetherian object	II.7
normalized dualizing complex	V.6
pointwise bounded below	V.8
pointwise dualizing complex	V.8
pointwise finite injective dimension	V.8
projectively embeddable	III.10
rational point	VII.1
reflexive complex	III.1
regular prescheme	V.2
relative differentials	V.2
residual complex	VI.1
residually stable morphism	VI.5
residue of a differential	VII.1
residue symbol	III.9
sheaf of families of supports	IV.1
smooth morphism	III.1
translation functor	I.1
triangle	IV.2
triangulated category	I.1

way-out functor	I.1
Z/Z' -skeleton	1.7

Index of Notations

Of course there is the usual collection of variable notations: A, B for abelian categories or rings, functors or sheaves, X, Y for preschemes, x, y for points of them, f, g for morphisms of preschemes, ϕ for morphisms of functors on sheaves, etc.

In general no underline denotes a group, as $\mathrm{Hom}(\mathcal{F}, \mathcal{G})$, one underline denotes a sheaf, as $\underline{\mathrm{Hom}}(\mathcal{F}, \mathcal{G})$ or $\underline{\mathrm{Tor}}(\mathcal{F}, \mathcal{G})$, two underlines denotes an object in a derived category, as $\underline{\underline{\mathrm{R}}}\mathrm{Hom}^\bullet(\mathcal{F}^\bullet, \mathcal{G}^\bullet)$, and a dot denotes a complex.

In the list of stable notations below, we have distinguished five categories: Latin alphabet, other alphabets, superscripts, subscripts, and arbitrary symbols.

Latin alphabet

Symbol	Meaning	Reference
Ab	the category of abelian groups	
acc	ascending chain condition	V.2
Ass	associated primes	
B	boundaries of a complex	
Coh	category of coherent sheaves	II.1
Coz	category of Cousin complexes	
D	dualizing functor	V.0
$\mathcal{D}(\mathcal{A})$	derived category	I.4
$\mathcal{D}(X)$	$= \mathcal{D}(\mathrm{Mod}(X))$	II.1
dcc	descending chain condition	V.2
dim	dimension	
Dual	category of dualizing complexes	VI.1
E	associated Cousin complex	IV.2, IV.3
Ext	Ext groups	I.6
$\underline{\mathrm{Ext}}$	sheaf Ext	II.3

Symbol	Meaning	Reference
fid	finite injective dimension	I.7.6, II.7.20
FR1–FR5	axioms of multiplicative systems	I.3
H	cohomology	I.1, IV.1
Hom	Hom groups	I.6
<u>Hom</u>	sheaf Hom	II.3
i_x	closed immersion	II.7
Icz	category of injective Cousin complexes	
id	identity map	
im	image	
$J(x), J(x, x')$	standard injectives	II.7
$\mathcal{K}(A)$		II.7
$k(x)$	residue field of a point x	
$K_\bullet(\underline{f}), K^\bullet(\underline{f}; F)$	Koszul complexes	III.7
ker	kernel	
L1–L3	axioms of direct systems	I.3
$Der L$	left derived functor	I.5
lfr	locally free of finite rank	II.5
Lno	category of locally noetherian preschemes	III.8
$\text{Mod}(X)$	category of $\mathcal{O}_X$ -modules	II.1
Ob	objects of a category	
pfid	pointwise finite injective dimension	V.8
Ptwdual	category of pointwise dualizing complexes	VI.1
Q	functor into derived category	I.3
Qco	category of quasi-coherent sheaves	II.1
Qis	quasi-isomorphisms	I.4
<u>R</u>	right derived functor	I.5
Res	category of residual complexes	VI.1
	residue symbol	
	residue of a differential	III.9
Spec	spectrum of a ring	VII.1
Supp	support of a sheaf or module	II.7
T	translation functor	I.1
<u>Tor</u>		
Tr	trace map	III.10, VI.4, VII.1
TR1–TR4	axioms of a triangulated category	I.1
TRA	axioms of trace	III.10.5, VI.4.2

Symbol	Meaning	Reference
tr. d	transcendence degree	
$\mathrm{Tr} f$	trace for a finite morphism	III.6
$\mathrm{Tr} p$	trace for a projective morphism	III.11
VAR	axioms for $f^!$ or f^Δ	III.8.7, VI.3.1
Z	cycles of a complex	

Greek and other alphabets

Symbol	Meaning	Reference
γ	trace for smooth morphism	III.3.4, III.11.2, VII.4.1
Γ	global sections of a sheaf	II.1.5
ζ	change of differentials	III.7.3
η	fundamental local isomorphism	III.7.3
θ	duality morphism	III.7.3
ξ	map of a functor to its derived functor	III.7.3
$\sigma_{\geq n}$	truncated complex	I.7
$\tau_{\geq n}$	truncated complex	I.7
ψ	residue isomorphism	III.8
ω	sheaf of differentials	III.1
Ω	sheaf of differentials	III.1
$\mathcal{O}_X$	structure sheaf of a prescheme	
$\mathfrak{m}$	maximal ideal of a ring	
$\mathfrak{p}$	prime ideal of a ring	
$\mathbb{A}$	affine space	
$\mathbb{P}$	projective space	
$\mathbb{Q}$	rational numbers	
$\mathbb{Z}$	integers	

Subscripts

Symbol	Meaning	Reference
--------	---------	-----------

$\mathcal{A}'$	as in $\mathcal{K}_{\mathcal{A}'}(\mathcal{A}), \mathcal{D}_{\mathcal{A}'}(\mathcal{A})$	I.4
c	as in $\mathcal{D}_c(X)$	II.1
fid	as in $\mathcal{D}(X)_{\text{fid}}$	I.7
fTd	as in $\mathcal{D}(X)_{\text{fTd}}$	II.4
$\text{Gor}(Z^\bullet)$	as in $\mathcal{D}(X)_{\text{Gor}(Z^\bullet)}$	IV.3
$\mathfrak{p}$	as in $A_{\mathfrak{p}}, M_{\mathfrak{p}}$: localization at a prime ideal	
qc	as in $\mathcal{D}_{\text{qc}}(X)$	II.1
x	as in $\mathcal{O}_x$: local ring at a point x	
	as in $\mathcal{F}_x$: stalk of a sheaf at x	
	as in H_x^i : local cohomology	
Z	as in Γ_Z, H_Z^i	IV.1
I, II	as in $\underline{\underline{R}}_I, \underline{\underline{R}}_{II}$	IV.1
$\geq n$	as in $\sigma_{\geq n}, \tau_{\geq n}$	IV.1
$*$	as in f_* : direct image	IV.1

Superscripts

Symbol	Meaning	Reference
b	as in $\mathcal{K}^b, \mathcal{D}^b$: bounded	I.2, I.4
i	as in X^i : i -th term of a complex	
	as in H^i : i cohomology group	
	as in $R^i F$: i -th derived functor	
y	as in f^y	VI.2
z	as in f^z	VI.2
$\bullet$	as in $\mathcal{F}^\bullet, X^\bullet$: denotes a complex	
$\bullet\bullet$	as in $\mathcal{C}^{\bullet\bullet}$: double complex	I.7
$\bullet$	as in $\mathcal{K}(A)^\bullet$: opposed category	
$*$	as in f^* : inverse image	I.7, II.4
$+$	as in $\mathcal{K}^+, \mathcal{D}^+, \underline{\underline{R}}^+$: bounded below	I.2, I.4, I.5
$-$	as in $\mathcal{K}^-, \mathcal{D}^-, \underline{\underline{R}}^-$: bounded above	I.2, I.4, I.5
$!$	as in $f^!$	Intr., III.8, VII.3
$\flat$	$f^\flat$	IX.6
$\sharp$	$f^\sharp$	VI.3
Δ	f^Δ	VI.3
$\vee$	as in $L^\vee$: dual sheaf	II.5.16, III.1
$-$	as in $\{x\}^-$: closure	

$\sim$	as in $\tilde{M}$: sheaf associated to a module
$\hat{\wedge}$	as in $\hat{A}$: completion of local ring

Arbitrary Symbols

Symbol	Meaning	Reference
$ $	restriction	
$\gg$	enough larger than	
$\times$	product	
$\otimes$	tensor product	
$\oplus$	direct sum	
$\coprod$	disjoint union / sum	
$[]$	shift operator	I.1
	closed interval	I.1
	reference to Bibliography or other chapter	
$\cup$	union	
$\{ \}$	the set of	
$\rightarrow$	morphism	
$\mapsto$	effect of a map on elements	
$\cdot \rightarrow$	distinguished side of triangle	
$--\rightarrow$	morphism being constructed	
$\circ$	composition of functors ($F \circ G$)	I.2

Bibliography

- [AB66] Michael Francis Atiyah and Raoul Bott. “A Lefschetz fixed point formula for elliptic differential operators”. In: *Bulletin of the American Mathematical Society* 72 (1966), pp. 245–250.
- [Art62] Michael Artin. *Grothendieck Topologies*. Mimeographed Seminar Notes, Harvard University. Cambridge, MA, 1962.
- [Bas62] Hyman Bass. “Injective dimension in noetherian rings”. In: *Transactions of the American Mathematical Society* 102 (1962), pp. 18–29.
- [Bas63] Hyman Bass. “On the ubiquity of Gorenstein rings”. In: *Mathematische Zeitschrift* 82 (1963), pp. 8–28.
- [CE56] Henri Cartan and Samuel Eilenberg. *Homological Algebra*. Princeton, NJ: Princeton University Press, 1956.
- [Che51] Claude Chevalley. *Introduction to the theory of algebraic functions of one variable*. American Mathematical Society Surveys. Providence, RI: American Mathematical Society, 1951.
- [ES53] Beno Eckmann and Arnold Schopf. “Über injektive Moduln”. In: *Archiv der Mathematik* 4 (1953), pp. 75–78.
- [Gab62] Pierre Gabriel. “Des Catégories Abéliennes”. In: *Bulletin de la Société Mathématique de France* 90 (1962), pp. 323–448 (cit. on p. 7).
- [Gir60] Jean Giraud. “Unpublished Thesis”. To appear. PhD thesis. Institut des Hautes Études Scientifiques, 1960 (cit. on p. 8).
- [God58] Roger Godement. *Topologie algébrique et théorie des faisceaux*. Paris: Hermann, 1958.
- [Gor52] Daniel Gorenstein. “An arithmetic theory of adjoint plane curves”. In: *Transactions of the American Mathematical Society* 72 (1952), pp. 414–436.

- [Gro57a] Alexander Grothendieck. “Sur quelques points d’algèbre homologique”. In: *Tohoku Mathematical Journal* 9 (1957), pp. 119–221.
- [Gro57b] Alexander Grothendieck. *Théorèmes de dualité pour les faisceaux algébriques cohérents*. Séminaire Bourbaki, No. 149, I.H.P. Paris. 1957 (cit. on pp. i, 2).
- [Gro60a] Alexander Grothendieck. *Éléments de Géométrie Algébrique*. 4, 8, 11, 17, 20, 24, 28 vols. Publications Mathématiques de l’IHÉS. 1960 ff. Paris: Institut des Hautes Études Scientifiques, 1960.
- [Gro60b] Alexander Grothendieck. “The cohomology theory of abstract algebraic varieties”. In: *Proceedings of the International Congress of Mathematicians, Edinburgh 1958*. Cambridge: Cambridge University Press, 1960, pp. 103–118 (cit. on pp. i, 2).
- [Gro61] Alexander Grothendieck. *Local Cohomology*. Mimeographed Seminar Notes (Notes by R. Hartshorne), Harvard University. Cambridge, MA, 1961.
- [Gro62] Alexander Grothendieck. *Séminaire de Géométrie Algébrique*. Polycopied Notes, I.H.E.S. Paris. 1960–1961; 1962. 1962.
- [Gro63] Alexander Grothendieck. *Residus et dualité*. Pré-notes pour un Séminaire Hartshorne, Manuscript. Paris, 1963 (cit. on p. i).
- [Gro66] Alexander Grothendieck. “De Rham cohomology of algebraic varieties”. In: *Publications Mathématiques de l’IHÉS* (1966). To appear.
- [Har66] Robin Hartshorne. “A property of A-sequences”. In: *Bulletin de la Société Mathématique de France* 94 (1966), pp. 61–66.
- [Mac63] Saunders Mac Lane. *Categorical Algebra*. Colloquium Lectures, 68th AMS Summer Meeting, Boulder. Aug. 1963.
- [Mat58] Ephraim Matlis. “Injective modules over noetherian rings”. In: *Pacific Journal of Mathematics* 8 (1958), pp. 511–528 (cit. on p. 7).
- [Nag62] Masayoshi Nagata. *Local Rings*. Vol. 13. Interscience Tracts in Pure and Applied Mathematics. New York: J. Wiley and Sons, 1962.
- [Ser55a] Jean-Pierre Serre. “Faisceaux algébriques cohérents”. In: *Annals of Mathematics* 61 (1955), pp. 197–278 (cit. on pp. 1, 2).
- [Ser55b] Jean-Pierre Serre. “Un théorème de dualité”. In: *Commentarii Mathematici Helvetici* 29 (1955), pp. 9–26 (cit. on p. 2).
- [Ser59] Jean-Pierre Serre. *Groupes algébriques et corps de classes*. Paris: Hermann, 1959 (cit. on p. 1).

- [Ver63a] Jean-Louis Verdier. “Théorème de dualité de Poincaré”. In: *Comptes Rendus de l’Académie des Sciences* 256 (1963), pp. 2084–2086.
- [Ver63b] Jean-Louis Verdier. “Unpublished Thesis”. To appear. PhD thesis. Institut des Hautes Études Scientifiques, 1963 (cit. on p. 7).
- [Zar52] Oscar Zariski. “Complete linear systems on normal varieties and a generalization of a lemma of Enriques-Severi”. In: *Annals of Mathematics* 55 (1952), pp. 552–592 (cit. on p. 1).
- [Zar56] Oscar Zariski. “Algebraic sheaf theory”. In: *Bulletin of the American Mathematical Society* 62 (1956), pp. 117–141 (cit. on p. 1).
- [ZS58] Oscar Zariski and Pierre Samuel. *Commutative Algebra*. Vol. 1, 2. Vol.1:1958, Vol.2:1960. Princeton, NJ: Van Nostrand, 1958.

Cohomologie à Support Propre, et Construction du Foncteur $f^!$

by P. Deligne

Errata