Zeta_r Drivers

Petra Guy

10 May 2018

Modeling the response of zeta_r to the abiotica/heterogeneity varaibles






```
##   Northing        PHI    meandbh     meanph     Buffer    meanSOM
##   1.730083   1.215926   2.031896   1.377587   1.819210   1.162576
##    meanLBA     meanTD area_ratio     no_NVC     no_MSG
##   1.706256   1.950731   1.230848   1.299008   1.276061
```

```
##   Northing        PHI     Buffer     no_MSG     no_NVC      sd_pH
##   1.713097   1.275714   1.860168   1.241747   1.277452   1.328220
##     sd_SOM     sd_LBA sd_meandbh      sd_TD area_ratio
##   1.412783   1.188175   1.521227   1.501602   1.257226
```

The variance inflation factors in the sd dataset are also low

The models which had a delta <2 were selected from the MuMin dredge funtion as the top model set.



The graph shows the averaged effect sizes of the model with delta < 2. Number of NVC codes, mean tree density and mean soil organic matter effect the plot level Z



In the sd dataset model the standard deviaiton of mean dbh has an effect. ##Using the model for prediction


Observed versus predicted data, mean dataset
R2 = 0.39
empirical_Z
predicted_Z
0.8
0.7
0.6
0.5
0.4
0.3
0.45
0.50
0.55
0.60
0.65
0.70


Observed versus predicted data, sd dataset
R2 = 0.31
empirical_Z
predicted_Z
0.8
0.7
0.6
0.5
0.4
0.3
0.50
0.55
0.60
0.65
0.70