

Colon & Colorectal Cancer — Clinical Trials Intelligence

Alert Window: Last 30 Days · Generated: 2026-05-02 10:17:00

Situation Summary

Phase I/II: 68 trials
Recruiting: 372 trials
Biomarker: 309 biomarker-directed trials
Breakthrough: 151 trials scored ≥ 4

Active Trial Timeline

Showing 151 high-significance trials (score ≥ 4). 464 additional lower-scored trials are collapsed below. Phase III trials appear first. Click any title to view full trial details on ClinicalTrials.gov.

MEDIUM **PHASE3** **Stage IV** **MSI/MMR** **TARGETED THERAPY** • **RECRUITING**

⚡ 11

Botensilimab + Balstilimab vs Best Supportive Care as Therapy in Chemo-refractory, Unresectable, Colorectal Adenocarcinoma

This clinical trial is testing the combination of botensilimab and balstilimab in patients with chemo-refractory, unresectable colorectal adenocarcinoma. The intervention involves two immunotherapy drugs aimed at potentially improving overall survival and quality of life for participants. Currently in Phase 3 and actively recruiting, this trial may provide insights into the effectiveness and safety of these treatments for eligible patients. Participants may need to meet specific biomarker criteria, including MSI, MSI-H, MMR, dMMR, and microsatellite status.

NCT: NCT07152821 · Enrollment: 834 · Sites: Greenfield Park, Canada, Kingston, Canada, Oshawa, Canada, Ottawa, Canada, Prince George, Canada

MEDIUM **PHASE3** **Stage III** **MSI/MMR** **TARGETED THERAPY** • **NOT_YET_RECRUITING**

⚡ 11

Neoadjuvant/Adjuvant AK104 in Microsatellite Instability-high or Mismatch Repair-deficient, Resectable Colon Cancer

This clinical trial is testing the efficacy and safety of neoadjuvant/adjuvant treatment with AK104 (Cadonilimab) in patients with resectable microsatellite instability-high (MSI-H) or mismatch repair-deficient (dMMR) colon cancer. The intervention involves AK104 in combination with standard chemotherapy agents, including Oxaliplatin, Capecitabine, 5-Fluorouracil, and Calcium Folate, which is significant for potentially improving treatment outcomes in this specific patient population. This is a Phase 3 trial that is currently not yet recruiting, meaning patients will need to wait for enrollment to begin before participating. Eligibility criteria include having resectable colon cancer with specific biomarkers such as MSI-H or dMMR.

NCT: NCT07412613 · Enrollment: 386 · Sites: Guangzhou, China

MEDIUM **PHASE3** **Stage IV** **BRAF** **TARGETED THERAPY** • **RECRUITING**

⚡ 11

Symbiotic-GI-03: A Study to Learn About the Study Medicine Called PF-08634404 in Combination With Chemotherapy in Adult Participants With Metastatic Colorectal Cancer

The Symbiotic-GI-03 trial is investigating the efficacy of the study medicine PF-08634404 in combination with chemotherapy for adult participants with metastatic colorectal cancer. This trial aims to determine how well PF-08634404 works alongside standard chemotherapy and Bevacizumab, a biological therapy, in patients whose cancer has spread or

returned after previous treatments. Currently in Phase 3 and actively recruiting, this trial offers eligible patients the opportunity to receive potentially beneficial treatment options while contributing to research on colorectal cancer therapies. To participate, individuals must be 18 years or older, have metastatic colorectal cancer, and meet specific health criteria, including not being pregnant prior to treatment.

NCT: NCT07222800 · Enrollment: 800 · Sites: Aberdeen, Abilene, Albany, Amarillo, Arlington

MEDIUM

PHASE3

Stage IV

MSI/MMR

TARGETED THERAPY

• NOT_YET_RECRUITING

11

Irinotecan Liposomes in Total Neoadjuvant Therapy in Locally Advanced Rectal Cancer

This clinical trial is testing the efficacy and safety of irinotecan liposomes in combination with standard total neoadjuvant therapy for patients with locally advanced rectal cancer. The intervention involves the use of NALIRI (irinotecan liposomes), along with LCRT (radiation) and FOLFOX (chemotherapy), to evaluate its impact on treatment outcomes. This is a Phase 3 trial that is not yet recruiting, meaning that patients will need to wait until enrollment begins to participate. Eligibility criteria include specific biomarkers such as MSI, MSI-H, MMR, DMMR, microsatellite status, and mismatch repair status.

NCT: NCT07074353 · Enrollment: 360 · Sites: Hangzhou, China

MEDIUM

PHASE2, PHASE3

Stage IV

MSI/MMR

TARGETED THERAPY

• RECRUITING

11

Node-Sparing Short-Course Radiotherapy Plus Chemotherapy, Bevacizumab and PD-1 Inhibitor in Metastatic pMMR/MSS Colorectal Cancer (MODIFI-CRC)

This clinical trial, known as MODIFI-CRC, is testing the effectiveness of node-sparing short-course radiotherapy combined with chemotherapy, bevacizumab, and a PD-1 inhibitor in patients with metastatic pMMR/MSS colorectal cancer. The intervention aims to enhance tumor response by utilizing radiotherapy to improve the efficacy of immunotherapy and targeted treatments, addressing the limitations of current standard therapies. Currently in Phase 2/3 and actively recruiting participants, this trial may offer patients an opportunity to receive a novel treatment approach that could potentially improve their outcomes. Eligibility criteria include specific biomarker assessments such as MSI, MMR, and NGS.

NCT: NCT06958419 · Enrollment: 286 · Sites: Guangzhou, China

MEDIUM

PHASE3

All Stages

KRAS

TARGETED THERAPY

• RECRUITING

11

A Study of Amivantamab and FOLFIRI Versus Cetuximab/Bevacizumab and FOLFIRI in Participants With KRAS/NRAS and BRAF Wild-type Colorectal Cancer Who Have Previously Received Chemotherapy

This clinical trial is testing the efficacy of amivantamab combined with FOLFIRI chemotherapy in patients with KRAS/NRAS and BRAF wild-type colorectal cancer who have previously undergone chemotherapy. The intervention involves the biological agent amivantamab alongside standard chemotherapy agents, which is important for evaluating its potential benefits compared to existing treatment options. This is a Phase 3 trial currently recruiting 700 participants, which means that eligible patients may have the opportunity to receive a treatment that is being compared to standard therapies in a larger population. Eligibility criteria include specific biomarker requirements related to KRAS, NRAS, and BRAF status, as well as other factors such as microsatellite instability (MSI) and mismatch repair (MMR) status.

NCT: NCT06750094 · Enrollment: 700 · Sites: Abilene, Ann Arbor, Arlington, Athens, Atlanta

MEDIUM

PHASE3

Stage III

MSI/MMR

TARGETED THERAPY

• RECRUITING

11

A Study of Neoadjuvant QL1706 in Participants With Untreated dMMR/MSI-H Resectable Colon Cancer

This clinical trial is testing the efficacy of neoadjuvant QL1706 in participants with untreated dMMR/MSI-H resectable colon cancer. The intervention involves administering QL1706 alongside CAPEOX/Capecitabine to assess its impact on tumor response prior to surgery. Currently in Phase 3 and actively recruiting, this trial aims to determine the proportion of patients achieving a pathological complete response (pCR), which indicates no residual tumor after treatment. Eligibility criteria include having T4N0 or Stage III colon cancer with specific biomarkers such as MSI-H or dMMR.

NCT: NCT06686576 · Enrollment: 363 · Sites: Guangzhou, China

SIGNAL

PHASE3

Stage III

ctDNA

LIQUID BIOPSY

• RECRUITING

11

ctDNA-guided Selection of Adjuvant Chemotherapy Regimens for Elderly Colon Cancer Patients After Surgery: A Single-center, Randomized, Controlled Study

This clinical trial is testing the effectiveness of ctDNA-guided selection of adjuvant chemotherapy regimens in elderly patients with high-risk stage II and stage III colon cancer following surgery. Participants will receive either 6 months of 5-FU monotherapy or an intensive XELOX treatment regimen, with the aim of improving disease-free survival outcomes. Currently in Phase 3 and actively recruiting, this trial seeks to determine how ctDNA detection can inform treatment decisions and patient prognosis. Eligibility criteria include being an elderly patient with high-risk stage II or stage III colon cancer and undergoing ctDNA testing to assess disease status.

NCT: NCT06609551 · Enrollment: 312 · Sites: Hangzhou, China

MEDIUM

PHASE2, PHASE3

Stage IV

MSI/MMR

TARGETED THERAPY

• RECRUITING

11

Sacituzumab Govitecan in Metastatic Colorectal Cancer

This clinical trial is testing the efficacy of Sacituzumab Govitecan (SG) in patients with metastatic colorectal cancer who have become refractory to at least two lines of standard chemotherapy and are not candidates for local therapy. The intervention involves administering SG compared to the physician's choice of treatment, which is significant as it may provide an alternative option for patients with limited treatment choices. This is a Phase II/III trial currently recruiting participants, indicating that patients may have the opportunity to access this investigational treatment while contributing to the research. Eligibility criteria include documented progression or intolerability to specific chemotherapy regimens and a minimum 6-month interval since the last use of Irinotecan.

NCT: NCT06243393 · Enrollment: 80 · Sites: Heidelberg, Germany

MEDIUM

PHASE3

Stage IV

MSI/MMR

TARGETED THERAPY

• ACTIVE_NOT_RECRUITING

11

Injection of SHR-A1811 Versus Physician Chocied Treatment in Patients With Advanced Colorectal Cancer Who Had Failed to Respond to Oxaliplatin, 5-fu, and Irinotecan

This clinical trial is testing the efficacy and safety of SHR-A1811 in patients with advanced colorectal cancer who have not responded to previous treatments with oxaliplatin, 5-FU, and irinotecan. The intervention involves the drug SHR-A1811, which is being compared to physician-chosen treatments such as TAS-102, regorafenib, and fruquintinib. This is a Phase 3 trial that is currently active but not recruiting, meaning that while the study is ongoing, no new participants are being enrolled at this time. Eligibility criteria include specific biomarkers such as MSI, MSI-H, MMR, DMMR, and RAS.

NCT: NCT06199973 · Enrollment: 130 · Sites: Shanghai, China

MEDIUM

PHASE3

Stage IV

MSI/MMR

TARGETED THERAPY

• RECRUITING

11

Preoperative Short-course Radiation Followed by Envafoimab Plus CAPEOX for MSS Locally Advanced Rectal Adenocarcinoma

This clinical trial is testing the combination of preoperative short-course radiation followed by Envafoimab and CAPEOX chemotherapy in patients with microsatellite stable (MSS) locally advanced rectal adenocarcinoma. The intervention involves the use of Envafoimab, a PD-L1 monoclonal antibody, which may enhance the effectiveness of short-course radiotherapy and chemotherapy. This is a Phase 3 trial currently recruiting participants, which means eligible patients may have the opportunity to receive this investigational treatment while contributing to important research. Eligibility criteria include specific biomarker requirements such as MSI, MSI-H, MMR, DMMR, and RAS status.

NCT: NCT05752136 · Enrollment: 108 · Sites: Hangzhou, China

MEDIUM

PHASE3

Stage IV

MSI/MMR

TARGETED THERAPY

• RECRUITING

11

Total Neoadjuvant Treatment ±Immunotherapy for High Risk Locally Advanced Rectal Cancer (TNTi)

The TNTi clinical trial is testing the effectiveness of total neoadjuvant treatment with and without the immunotherapy drug camrelizumab for patients with high-risk locally advanced rectal cancer. This intervention is significant as it aims to determine the pathologic complete response (PCR) rate, which is a critical measure of treatment effectiveness. Currently in Phase 3 and actively recruiting, this trial offers patients the opportunity to participate in research that may influence future treatment protocols. Eligibility criteria include specific biomarkers such as MSI, MMR, DMMR, RAS, and microsatellite status.

NCT: NCT06229041 · Enrollment: 472 · Sites: Beijing, China

MEDIUM

PHASE3

Stage IV

MSI/MMR

TARGETED THERAPY

• ACTIVE_NOT_RECRUITING

11

Study of XL092 + Atezolizumab vs Regorafenib in Participants With Metastatic Colorectal Cancer

This clinical trial is testing the combination of XL092 and atezolizumab against regorafenib in participants with metastatic colorectal cancer who have progressed during, after, or are intolerant to standard-of-care therapy. The intervention involves the use of XL092 and atezolizumab, which may provide a different therapeutic approach compared to regorafenib. This is a Phase 3 trial that is currently active but not recruiting, indicating that it is in the final stages of testing and may soon provide important data on treatment efficacy. Eligibility criteria include specific biomarkers such as MSI, MSI-H, MMR, DMMR, RAS, and microsatellite status.

NCT: NCT05425940 · Enrollment: 901 · Sites: Albuquerque, Billings, Charlotte, Chattanooga, Cincinnati

MEDIUM

PHASE3

Stage III

MSI/MMR

TARGETED THERAPY

• RECRUITING

11

PACE: PD-1 Antibody For dMMR/MSI-H Stage III Colorectal Cancer

The PACE trial is investigating the efficacy of the PD-1 antibody Sintilimab in patients with dMMR/MSI-H Stage III colorectal cancer. Participants will receive either Sintilimab alone or a combination of oxaliplatin and capecitabine, which are standard chemotherapy agents, to assess their impact on disease-free survival. This is a Phase 3 study currently recruiting 323 participants, indicating that patients may have the opportunity to access this treatment while contributing to important clinical research. Eligibility criteria include having local advanced colon cancer with specific biomarkers such as dMMR or MSI-H.

NCT: NCT05236972 · Enrollment: 323 · Sites: Guangzhou, China

MEDIUM

PHASE3

Stage IV

BRAF

TARGETED THERAPY

• ACTIVE_NOT_RECRUITING

11

A Study of Encorafenib Plus Cetuximab With or Without Chemotherapy in People With Previously Untreated Metastatic Colorectal Cancer

This clinical trial is evaluating the combination of encorafenib and cetuximab, with or without standard chemotherapy, in patients with previously untreated metastatic colorectal cancer that has a BRAF mutation. The intervention involves administering encorafenib orally and cetuximab via intravenous infusion, either alone or alongside chemotherapy agents such as oxaliplatin and irinotecan. This is a Phase 3 trial that is currently active but not recruiting, indicating that it is in the final stages of testing before potential approval, and patients may not be able to enroll at this time. Eligibility criteria include the presence of specific biomarkers such as BRAF mutations, MSI, MSI-H, MMR, or DMMR.

NCT: NCT04607421 · Enrollment: 831 · Sites: Aventura, Basking Ridge, Berkeley Heights, Chicago, City of Saint Peters

SIGNAL

PHASE2, PHASE3

All Stages

RESEARCH

• RECRUITING

10

Antegrade Intestinal Fluid Reinfusion for Prevention of Low Anterior Resection Syndrome After Low Anterior Resection: a Single-Center, Prospective Randomized Controlled Trial.

This clinical trial is testing the effectiveness of antegrade intestinal fluid reinfusion in preventing low anterior resection syndrome (LARS) in patients with a prophylactic ileal stoma. The intervention involves administering either intestinal fluid or potable water through the ileal stoma prior to ileostomy reversal surgery, which is important for understanding optimal management strategies for LARS. Currently in Phase 2 and Phase 3, the trial is actively recruiting 160 participants, which means eligible patients can join to contribute to this research. Eligibility criteria include having undergone anterior rectal resection and being scheduled for ileostomy reversal.

NCT: NCT07537998 · Enrollment: 160 · Sites: Guangzhou, China

MEDIUM

PHASE3

Stage IV

HER2

TARGETED THERAPY

• RECRUITING

10

A Study of Tucatinib With Trastuzumab and mFOLFOX6 Versus Standard of Care Treatment in First-line HER2+ Metastatic Colorectal Cancer

This clinical trial is testing the efficacy of tucatinib in combination with trastuzumab and mFOLFOX6 compared to standard of care treatments for patients with HER2 positive metastatic colorectal cancer. The intervention involves the use of tucatinib, a targeted therapy, alongside other established cancer drugs, to evaluate its effectiveness and side effects in this patient population. Currently in Phase 3 and actively recruiting, this trial aims to provide insights into progression-free survival outcomes for participants. Eligibility criteria include having HER2 positive colorectal cancer that is metastatic or unresectable, with specific biomarker assessments for HER2 and RAS.

MEDIUM

PHASE2

Stage IV

MSI/MMR

LIQUID BIOPSY

• ACTIVE_NOT_RECRUITING

⚡ 10

Fecal Microbiota Transplant and Re-introduction of Anti-PD-1 Therapy (Pembrolizumab or Nivolumab) for the Treatment of Metastatic Colorectal Cancer in Anti-PD-1 Non-responders

This phase II clinical trial is investigating the combination of fecal microbiota transplantation and the re-introduction of anti-PD-1 therapy (pembrolizumab or nivolumab) for patients with metastatic colorectal cancer who have not responded to previous anti-PD-1 treatments. The intervention includes procedures such as fecal microbiota transplantation and the use of specific drugs, which aim to enhance the immune response against cancer by restoring the gut microbiome. Currently, the trial is active but not recruiting new participants, indicating that it is ongoing with enrolled patients receiving treatment. Eligibility criteria include specific biomarkers such as MSI, MSI-H, MMR, DMMR, CTDNA, microsatellite status, and mismatch repair status.

NCT: NCT04729322 · Enrollment: 15 · Sites: Houston

SIGNAL

PHASE3

Stage IV

MSI/MMR

RESEARCH

• RECRUITING

⚡ 9

Neoadjuvant CAPOX With or Without Pucotenlimab Plus Selective Radiotherapy for Locally Advanced Rectal Cancer

This clinical trial is testing the effectiveness of neoadjuvant CAPOX with or without pucotenlimab in patients with locally advanced rectal cancer. The intervention involves administering four cycles of CAPOX, a chemotherapy regimen, combined with the investigational drug pucotenlimab, which may enhance treatment outcomes. This is a phase III trial currently recruiting 556 participants, which means patients may have the opportunity to access this treatment approach while contributing to the evaluation of its efficacy and safety. Eligible patients must have pMMR/MSS locally advanced rectal cancer and will be randomized based on mesorectal fascia status.

NCT: NCT07528209 · Enrollment: 556 · Sites: Guangzhou, China

MEDIUM

PHASE2

Stage IV

MSI/MMR

TARGETED THERAPY

• NOT_YET_RECRUITING

⚡ 9

Neoadjuvant Immunotherapy ± Radiotherapy in MSI-H/dMMR Locally Advanced Colorectal Cancer

This phase II clinical trial is testing the efficacy and safety of three neoadjuvant regimens for patients with locally advanced microsatellite instability-high/mismatch repair-deficient (MSI-H/dMMR) colorectal cancer. The interventions include dual immune checkpoint blockade with nivolumab and ipilimumab, nivolumab plus radiotherapy, and nivolumab monotherapy, which are important for evaluating different approaches to enhance treatment outcomes. The trial is not yet recruiting participants, meaning patients will need to wait until enrollment begins to participate. Eligibility criteria include the presence of MSI, MSI-H, MMR, or dMMR biomarkers.

NCT: NCT07403877 · Enrollment: 114 · Sites: Shanghai, China

MEDIUM

PHASE3

All Stages

MSI/MMR

TARGETED THERAPY

• NOT_YET_RECRUITING

⚡ 9

Prophylactic PIPAC in Patients With High-risk Colorectal Cancer

This clinical trial is testing the efficacy of oxaliplatin-based Pressurized Intraperitoneal Aerosol Chemotherapy (PIPAC) in patients with high-risk colorectal cancer. The intervention involves administering PIPAC in conjunction with standard surgery and adjuvant chemotherapy, which may provide a targeted approach to prevent peritoneal metastasis. This is a Phase 3 trial that is not yet recruiting, meaning patients will need to wait until enrollment begins to participate. Eligibility criteria include specific biomarkers such as MSI, MMR, DMMR, and NGS, which will help identify suitable candidates for the study.

NCT: NCT06839092 · Enrollment: 160 ·

MEDIUM

PHASE3

All Stages

MSI/MMR

TARGETED THERAPY

• NOT_YET_RECRUITING

⚡ 9

Study Evaluating Two Treatment Strategies for Rectal Cancer in Patients ≥75 Years Old

The NACRE-2 trial is evaluating two treatment strategies for patients aged 75 years and older with locally advanced rectal cancer. The study compares the effectiveness of adding chemotherapy (FOLFOX4s) after short-course radiotherapy against short-course radiotherapy alone. This is a Phase 3 trial that is not yet recruiting, meaning patients will need to wait

for enrollment to begin before participating. Eligibility criteria include specific biomarkers such as MSI, MMR, DMMR, and microsatellite status.

NCT: NCT07118800 · Enrollment: 160 ·

SIGNAL **PHASE3** **Stage IV** **RESEARCH** • **RECRUITING**

🔥 9

AK112 and Chemotherapy in First-line Metastatic Colorectal Cancer

This Phase III clinical trial is evaluating the efficacy and safety of AK112 in combination with chemotherapy for patients with first-line metastatic colorectal cancer. The intervention includes AK112 along with standard chemotherapy agents such as oxaliplatin, irinotecan, leucovorin, and 5-FU, compared to bevacizumab with chemotherapy. Currently, the trial is in the recruiting phase, meaning eligible patients can still enroll to participate. Enrollment is set for 560 participants, and the primary outcome being assessed is progression-free survival as determined by blinded independent central review.

NCT: NCT06951503 · Enrollment: 560 · Sites: Guanzhou, China

SIGNAL **PHASE3** **All Stages** **ctDNA** **LIQUID BIOPSY** • **RECRUITING**

🔥 9

tislelizUMaB in canceR Patients With molEcuLar residual Disease

This clinical trial is testing the efficacy of tislelizumab in patients with colorectal cancer who have molecular residual disease (MRD+) after surgery and perioperative treatments. The intervention involves administering tislelizumab, an immunotherapy drug, compared to a placebo, with the aim of improving disease-free survival (DFS) in this patient population. This is a Phase 3 trial currently recruiting 717 participants, which indicates that it is in the later stages of testing and may provide critical information on the effectiveness of tislelizumab for patients with MRD+. Eligibility criteria include the presence of circulating tumor DNA (ctDNA) and specific RAS biomarker requirements.

NCT: NCT06332274 · Enrollment: 717 · Sites: Villejuif, France

HIGH **PHASE3** **All Stages** **PHASE III TRIAL** • **ACTIVE_NOT_RECRUITING**

🔥 9

A Phase III Study to Evaluate the Efficacy, Immunogenicity and Safety of the 9-valent HPV Vaccine in Chinese Males

This Phase III clinical trial is evaluating the efficacy, immunogenicity, and safety of the 9-valent Human Papillomavirus (HPV) vaccine in Chinese males aged 18-45 years. The intervention involves administering the 9-valent HPV recombinant vaccine, which targets multiple HPV types, to determine its effectiveness in reducing the incidence of genital warts compared to a placebo. The trial is currently active but not recruiting participants, indicating that it is in the process of collecting data from enrolled subjects. Eligibility criteria include being a male aged 18-45 years, with no specific biomarker requirements mentioned.

NCT: NCT06465914 · Enrollment: 9000 · Sites: Changsha, China, Chengdu, China, Kunming, China, Nanning, China, Taiyuan, China

SIGNAL **PHASE3** **Stage IV** **MSI/MMR** **RESEARCH** • **RECRUITING**

🔥 9

DOSAGE Study: Upfront Dose-Reduced Chemotherapy in Older Patients with Metastatic Colorectal Cancer

The DOSAGE Study is a phase III clinical trial designed to evaluate the effectiveness of upfront dose-reduced chemotherapy compared to standard dose chemotherapy in older patients (≥70 years) with metastatic colorectal cancer. The trial investigates various chemotherapy regimens, including doublet chemotherapy and monotherapy, both at standard and reduced doses, to assess their impact on progression-free survival (PFS). Currently, the study is in the recruiting phase, meaning eligible patients can still enroll to participate in this research. Eligibility criteria include age and risk classification for chemotherapy toxicity, with specific biomarkers such as MSI and microsatellite status being relevant for patient selection.

NCT: NCT06275958 · Enrollment: 587 · Sites: 's-Hertogenbosch, Netherlands, Alkmaar, Netherlands, Amstelveen, Netherlands, Amsterdam, Netherlands, Arnhem, Netherlands

SIGNAL **PHASE3** **Stage IV** **BRAF** **RESEARCH** • **RECRUITING**

🔥 9

Intermittent or Continuous Panitumumab Plus FOLFIRI for Left Sided RAS/B-RAF Wild-type Metastatic Colorectal Cancer

This clinical trial is testing the effectiveness of intermittent versus continuous administration of Panitumumab combined with FOLFIRI in patients with left-sided RAS/B-RAF wild-type metastatic colorectal cancer. The intervention involves the use of Panitumumab, Irinotecan, 5-fluorouracil, and L-folinic acid, which are important for evaluating treatment strategies that may optimize patient outcomes. Currently in Phase 3 and actively recruiting, this trial aims to determine if the intermittent regimen can achieve a Time to Treatment Failure comparable to the standard continuous regimen, which is relevant for patients considering their treatment options. Eligibility criteria include specific biomarker requirements related to BRAF, MMR, and RAS status.

NCT: NCT06509126 · Enrollment: 500 · Sites: Naples, Italy

MEDIUM **PHASE2** **Stage IV** **BRAF** **TARGETED THERAPY** • **RECRUITING**

🔥 9

A Study To Evaluate The Efficacy And Safety Of Ifinatamab Deruxtecan (I-DXd) In Subjects With Recurrent Or Metastatic Solid Tumors (IDeate-PanTumor02)

This clinical trial is evaluating the efficacy and safety of ifinatamab deruxtecan (I-DXd) in patients with recurrent or metastatic solid tumors, including colorectal cancer. The intervention, ifinatamab deruxtecan, is a drug that targets specific cancer biomarkers, which may influence treatment outcomes. This is a Phase 2 trial currently recruiting 520 participants, indicating that patients may have the opportunity to access this treatment while contributing to the understanding of its effectiveness. Eligibility criteria include specific biomarkers such as BRAF, MSI, MSI-H, MMR, DMMR, HER2, and RAS.

NCT: NCT06330064 · Enrollment: 520 · Sites: Amarillo, Chattanooga, Dallas, Fairfax, Germantown

SIGNAL **PHASE3** **Stage III** **RESEARCH** • **RECRUITING**

🔥 9

Thymosin-alpha 1 for Adjuvant Treatment After Radical Resection of High-risk Stage II and III Colorectal Cancer

This clinical trial is testing the efficacy and safety of Thymosin-alpha 1 as an adjuvant treatment for patients with high-risk stage II and III colorectal cancer following radical resection. The intervention, Thymosin-alpha 1, is a drug believed to enhance immune response, potentially reducing the risk of cancer recurrence and metastasis. This is a Phase 3 trial currently recruiting 2,500 participants, which indicates that it is in the final stages of testing before potential approval and may provide valuable information for patients regarding treatment options. Eligibility criteria may include specific high-risk classifications of colorectal cancer, although detailed biomarker requirements are not specified.

NCT: NCT05086614 · Enrollment: 2500 · Sites: Shanghai, China

SIGNAL **PHASE3** **Stage IV** **RAS** **RESEARCH** • **ACTIVE_NOT_RECRUITING**

🔥 9

Neoadjuvant FOLFOX Chemotherapy for Patients With Locally Advanced Colon Cancer

This clinical trial is testing the effectiveness of neoadjuvant FOLFOX chemotherapy in patients with locally advanced colon cancer. The intervention involves administering FOLFOX chemotherapy before surgery, which may enhance treatment outcomes by targeting microscopic metastatic cancer cells earlier and reducing surgical complications. This is a Phase 3 trial that is currently active but not recruiting, indicating that it is in the final stages of testing and results may soon be available for clinical application. Eligibility criteria include the presence of RAS biomarkers, which may be relevant for patient selection.

NCT: NCT03426904 · Enrollment: 708 · Sites: Daegu, South Korea, Hwasun, South Korea, Seoul, South Korea, Suwon, South Korea

SIGNAL **PHASE3** **Stage IV** **RESEARCH** • **RECRUITING**

🔥 9

Bevacizumab Plus mFOLFOXIRI as First-line Treatment for Patients With Unresectable Metastatic Colorectal Cancer

This clinical trial is testing the efficacy of mFOLFOXIRI combined with Bevacizumab versus mFOLFOX6 combined with Bevacizumab as first-line treatments for patients with unresectable metastatic colorectal cancer. The intervention involves two chemotherapy regimens, which are significant as they aim to improve treatment outcomes while considering the unique responses of different populations, particularly in China. This is a Phase 3 trial currently recruiting 528 participants, which means eligible patients may have the opportunity to receive one of the investigational treatments while contributing to important research data. Eligibility criteria may include specific biomarkers, although details on these requirements are not provided.

NCT: NCT04230187 · Enrollment: 528 · Sites: Guangzhou, China

SIGNAL **PHASE3** **Stage III** **RESEARCH** • **RECRUITING**

⚡ 9

Randomised Study Evaluating Adjuvant Chemotherapy After Resection of Stage III Colonic Adenocarcinoma in Patients of 70 and Over

This clinical trial is evaluating the effectiveness of adjuvant chemotherapy in patients aged 70 and over who have undergone resection for stage III colonic adenocarcinoma. The interventions being tested include LV5FU2, capecitabine, FOLFOX4, XELOX, and observation, which are important for understanding the potential benefits of chemotherapy in this age group. Currently in Phase 3 and actively recruiting, this trial aims to provide insights into disease-free survival outcomes for elderly patients, which may influence treatment decisions. Eligibility criteria include being 70 years or older and having a diagnosis of stage III colonic adenocarcinoma.

NCT: NCT02355379 · Enrollment: 774 · Sites: Abbeville, France, Agen, France, Aix-en-Provence, France, Albi, France, Alès, France

SIGNAL **PHASE2, PHASE3** **All Stages** **RAS** **RESEARCH** • **NOT_YET_RECRUITING**

⚡ 8

Injection of IP-001 Into Thermally Ablated Hepatic Tumors in Patients With Colorectal Liver Metastases

This clinical trial is testing the efficacy of the injectable drug IP-001 in patients with colorectal cancer that has metastasized to the liver, specifically following standard microwave tumor ablation. The intervention involves administering either a high-dose or low-dose solution of IP-001 after the ablation procedure to assess its potential in prolonging progression-free survival compared to ablation alone. This study is in Phase 2 and Phase 3 and is currently not yet recruiting, indicating that patients will need to wait for enrollment to begin before participating. Eligibility criteria include being at least 18 years old and having RAS biomarker status relevant to the study.

NCT: NCT07321847 · Enrollment: 717 ·

SIGNAL **PHASE2, PHASE3** **Stage III** **MSI/MMR** **RESEARCH** • **RECRUITING**

⚡ 8

Efficacy and Safety Comparison of Short-course Radiotherapy Followed by CapeOX Chemotherapy Plus Toripalimab With or Without Concurrent Surufatinib in Neoadjuvant Therapy for Mid-to-low Localized Rectal Cancer of High-risk Criteria

This clinical trial is testing the efficacy and safety of short-course radiotherapy followed by CapeOX chemotherapy combined with Toripalimab, with or without concurrent Surufatinib, in patients with mid-to-low localized rectal cancer classified as high-risk. The intervention involves a combination of radiotherapy, chemotherapy, and immunotherapy, which aims to enhance the pathological complete response rate in this patient population. Currently in Phase 2 and Phase 3, the trial is recruiting participants, indicating that eligible patients may have the opportunity to access this investigational treatment. Eligibility criteria include specific biomarkers such as MMR and RAS, which may be relevant for patient selection.

NCT: NCT07505472 · Enrollment: 212 · Sites: Changchun, China

SIGNAL **PHASE3** **Stage IV** **RAS** **PHASE I TRIAL** • **RECRUITING**

⚡ 8

A Phase Ib/III Study of Suvemcitug Plus FTD/TPI in Participants With Refractory Metastatic Colorectal Cancer

This clinical trial is testing the combination of Suvemcitug and trifluridine/tipiracil tablets in participants with refractory metastatic colorectal cancer. The intervention involves administering Suvemcitug injections alongside trifluridine/tipiracil tablets to assess its safety and efficacy compared to a placebo. This is a Phase III study currently recruiting 464 participants, which means eligible patients may have the opportunity to receive a potentially effective treatment option while contributing to research. Eligibility criteria include the presence of RAS biomarkers.

NCT: NCT07361003 · Enrollment: 464 · Sites: Fuzhou, China, Hangzhou, China, Harbin, China, Jinan, China, Nanjing, China

MEDIUM **PHASE2** **Stage IV** **MSI/MMR** **TARGETED THERAPY** • **NOT_YET_RECRUITING**

⚡ 8

A Phase II Clinical Study Evaluating SSGJ-706 in Combination Therapy for Advanced Gastrointestinal Cancers

This phase II clinical trial is evaluating the combination of SSGJ-706 with standard therapies for patients with advanced gastrointestinal cancers. The interventions include SSGJ-706 combined with XELOX, Bevacizumab, AG, and TP, which are important for assessing the safety and effectiveness of this treatment approach. The trial is currently not yet recruiting

participants, meaning that patients will need to wait for enrollment to begin before they can participate. Eligibility criteria include specific biomarkers such as MSI, MSI-H, MMR, DMMR, and HER2.

NCT: NCT07233850 · Enrollment: 300 · Sites: Wuhan, China

SIGNAL **PHASE3** **Stage III** **MSI/MMR** **RESEARCH** • NOT_YET_RECRUITING

⚡ 8

Neoadjuvant Chemoradiotherapy Followed by Chemotherapy With or Without Tislelizumab for Resectable Ultra-low Rectal Cancer: The RELIEVE-02 Study

The RELIEVE-02 study is a Phase 3 clinical trial designed for patients with pathologically confirmed, previously untreated, resectable ultra-low rectal adenocarcinoma, specifically those with MSI-L or MSS/pMMR tumors. The trial investigates the efficacy of long-course chemoradiotherapy followed by chemotherapy with or without the addition of Tislelizumab, an immunotherapy drug, which may enhance treatment outcomes. Currently, the trial is not yet recruiting participants, meaning patients will need to wait for enrollment to begin before they can participate. Eligibility criteria include specific biomarker requirements such as MSI, MMR, and NGS status.

NCT: NCT07132463 · Enrollment: 154 ·

SIGNAL **PHASE2, PHASE3** **Stage III** **RESEARCH** • RECRUITING

⚡ 8

Total Neoadjuvant Therapy and Organ Preservation Versus Surgery for Rectal Cancer.

This clinical trial is testing the effectiveness of total neoadjuvant therapy (TNT) and a watch-and-wait strategy for patients with rectal cancer. The intervention includes radiation therapy, chemoradiotherapy, and consolidation chemotherapy, with the aim of potentially allowing patients to preserve their rectum if they achieve a complete or near-complete clinical response. Currently in Phase 2 and Phase 3, the trial is recruiting 400 participants, which means eligible patients may have the opportunity to contribute to research that could impact treatment approaches. Eligibility criteria include patients with cT1N1, T2-T3 N0-1 rectal cancer, and specific markers such as MRF- and EMVI-.

NCT: NCT06758830 · Enrollment: 400 · Sites: Vilnius, Lithuania

MEDIUM **PHASE3** **All Stages** **MSI/MMR** **IMMUNOTHERAPY** • RECRUITING

⚡ 8

Personalized Long-course Radiotherapy Plus Chemotherapy With or Without Immunotherapy for LARC: PALACE Study

The PALACE study is a phase III clinical trial designed for patients with locally advanced rectal cancer (LARC) to evaluate the complete response rate to personalized long-course radiotherapy combined with chemotherapy, with or without the anti-PD-1 antibody drug Serplulimab. This intervention aims to determine the effectiveness of adding immunotherapy to standard treatment in achieving a complete response, defined as a pathological complete response plus a clinical complete response sustained for over one year. Currently, the trial is recruiting participants, which means eligible patients have the opportunity to join and contribute to this research. Eligibility criteria include specific biomarkers such as MMR and RAS.

NCT: NCT07020247 · Enrollment: 184 · Sites: Chengdu, China

SIGNAL **PHASE3** **Stage III** **MSI/MMR** **RESEARCH** • RECRUITING

⚡ 8

Planned Organ Preservation for Low Rectal Cancer After Neoadjuvant Chemotherapy (OPLAR)

The OPLAR trial is testing the effectiveness of total neoadjuvant therapy (TNT) versus immunotherapy combined with TNT (iTNT) in patients with low and intermediate-risk rectal cancer. The intervention involves administering two cycles of XELOX chemotherapy followed by either long-course chemoradiotherapy with consolidation chemotherapy or the same regimen with added immunotherapy, aiming to improve organ preservation rates. This is a Phase 3 trial currently recruiting participants, which means eligible patients may have the opportunity to receive either treatment while contributing to research that could inform future care. Eligibility criteria include having measurable disease with specific biomarkers such as MMR and RAS.

NCT: NCT07070622 · Enrollment: 186 · Sites: Chengdu, China

MEDIUM **PHASE2** **Stage IV** **MSI/MMR** **TARGETED THERAPY** • RECRUITING

⚡ 8

Dostarlimab for Locally Advanced or Metastatic Cancer (Non-colorectal/Non-endometrial) With Tumor dMMR/MSI

This clinical trial is testing the efficacy of dostarlimab, an immunotherapy drug, for adult patients with locally advanced or metastatic cancers that are not colorectal or endometrial and have deficient mismatch repair (dMMR) or microsatellite instability (MSI). The intervention involves comparing dostarlimab to standard chemotherapy as a first-line treatment, which is significant for understanding alternative therapeutic options for these specific cancer types. This is a Phase 2 trial currently recruiting 120 participants, which means eligible patients may have the opportunity to receive either treatment while contributing to research on their effectiveness. Eligibility criteria include adults aged 18 and older with specific cancer types, including duodenum and small bowel adenocarcinoma, gastric adenocarcinoma, and others with documented dMMR/MSI status.

NCT: NCT06333314 · Enrollment: 120 · Sites: Angers, France, Avignon, France, Besançon, France, Brest, France, Caen, France

MEDIUM

PHASE2

Stage IV

KRAS

LIQUID BIOPSY

• ACTIVE_NOT_RECRUITING

8

Phase II Study of ctDNA-guided Encorafenib Plus Cetuximab Retreatment in Patients BRAF V600E Mutated mCRC

This clinical trial is testing the effectiveness of ctDNA-guided retreatment with encorafenib and cetuximab in patients with BRAF V600E mutated metastatic colorectal cancer (mCRC). The intervention involves the combination of encorafenib and cetuximab, which is significant for patients who have previously benefited from these treatments. This is a Phase II study that is currently active but not recruiting, indicating that while the trial is ongoing, no new participants are being enrolled at this time. Eligible participants must have specific biomarker profiles, including BRAF V600E mutation and wild-type KRAS, NRAS, and MAP2K1, as well as non-amplified MET status on ctDNA at the time of entry.

NCT: NCT06578559 · Enrollment: 16 · Sites: Meldola, Italy, Milan, Italy, Monserrato, Italy, Naples, Italy, Padova, Italy

SIGNAL

PHASE3

Stage IV

BRAF

RESEARCH

• NOT_YET_RECRUITING

8

Nimotuzumab Combined With Trifluridine/Tipiracil in the Treatment of Refractory Metastatic Colorectal Cancer

This clinical trial is testing the efficacy and safety of nimotuzumab combined with trifluridine/tipiracil in patients with refractory metastatic colorectal cancer (mCRC). The intervention involves the administration of nimotuzumab, a monoclonal antibody, alongside trifluridine/tipiracil, a chemotherapy regimen, to assess its impact on overall survival. This is a Phase 3 trial that is not yet recruiting, meaning patients will need to wait until enrollment begins to participate. Eligibility criteria include specific biomarkers such as BRAF, MMR, and RAS.

NCT: NCT06343116 · Enrollment: 420

HIGH

PHASE3

Stage IV

PHASE III TRIAL

• RECRUITING

8

Phase III Study Evaluating Induction Chemotherapy Followed by Chemoradiotherapy Compared to Standard Chemoradiotherapy for Locally Advanced SCCA

This Phase III clinical trial is evaluating the effectiveness of induction chemotherapy using the modified DCF protocol (mDCF) followed by chemoradiotherapy compared to standard chemoradiotherapy for patients with locally advanced squamous cell carcinoma of the anus (SCCA). The intervention involves administering mDCF, followed by concomitant chemotherapy with Capecitabine and Mitomycin-C, along with radiotherapy, aiming to improve disease-related event-free survival (DREFS) at two years. The trial is currently recruiting 310 participants, which means eligible patients can enroll to potentially access this treatment approach. Eligibility criteria may include specific tumor stages and the presence of biomarkers related to HPV.

NCT: NCT06207981 · Enrollment: 310 · Sites: Agen, France, Aix-en-Provence, France, Amiens, France, Antony, France, Argenteuil, France

MEDIUM

N/A

Stage IV

MSI/MMR

TARGETED THERAPY

• RECRUITING

8

The Efficacy of Watch and Wait Strategy or Surgery After Neoadjuvant Immunotherapy for Locally Advanced Colorectal Cancer With dMMR/MSI-H Guided by MRD Dynamic Monitoring: A Single-center, Open-label, Prospective, Phase II Clinical Trial.

This clinical trial is testing the efficacy of a watch and wait strategy compared to surgery after neoadjuvant immunotherapy in patients with locally advanced colorectal cancer characterized by deficient mismatch repair or microsatellite instability-high (dMMR/MSI-H). The intervention involves the use of Tislelizumab, a PD-1 inhibitor, which is significant for its potential to enhance the immune response against cancer cells. Currently, the trial is in the recruiting phase, indicating that eligible patients can still enroll to participate in this study. Eligibility criteria include specific biomarkers such as MSI, MSI-H, MMR, dMMR, RAS, and microsatellite status, as well as the use of next-generation sequencing (NGS) for patient selection.

MEDIUM

PHASE2, PHASE3

All Stages

MSI/MMR

TARGETED THERAPY

• RECRUITING

⚡ 8

DETERMINE Trial Treatment Arm 02: Atezolizumab in Adult, Paediatric and Teenage/Young Adult Patients With Cancers With High Tumour Mutational Burden (TMB) or Microsatellite Instability-high (MSI-high) or Proven Constitutional Mismatch Repair Deficiency (CMMRD) Disposition

The DETERMINE Trial Treatment Arm 02 is investigating the efficacy of atezolizumab in adult, pediatric, and teenage/young adult patients with cancers characterized by high tumor mutational burden (TMB), microsatellite instability-high (MSI-high), or proven constitutional mismatch repair deficiency (CMMRD). Atezolizumab is a drug that has been approved for several cancer types and is being evaluated for its potential benefits in additional cancers with specific genetic markers. This trial is currently in Phase 2 and Phase 3 and is actively recruiting participants, which means eligible patients may have the opportunity to receive this treatment while contributing to important research. Eligibility criteria include the presence of specific biomarkers such as MSI, TMB, and MMR, among others.

NCT: NCT05770102 · Enrollment: 30 · Sites: Belfast, United Kingdom, Birmingham, United Kingdom, Bristol, United Kingdom, Cambridge, United Kingdom, Cardiff, United Kingdom

MEDIUM

PHASE2

All Stages

KRAS

TARGETED THERAPY

• RECRUITING

⚡ 8

Pre-operative Targeted Treatments in Molecularly Selected Resectable Colorectal Cancer (UNICORN)

The UNICORN trial is testing pre-operative targeted treatments in patients with non-metastatic resectable colorectal cancer who have specific targetable molecular alterations. The intervention includes a combination of drugs such as Trastuzumab deruxtecan, Durvalumab, Panitumumab, Botensilimab, and Balstilimab, which are administered as a short-course strategy prior to standard surgical care. This is a Phase 2 trial currently recruiting participants, which means eligible patients may have the opportunity to receive these targeted therapies before surgery. Eligibility criteria include the presence of biomarkers such as KRAS, NRAS, BRAF, MSI, MSI-H, MMR, DMMR, and HER2.

NCT: NCT05845450 · Enrollment: 197 · Sites: Milan, Italy

SIGNAL

PHASE2, PHASE3

Stage IV

BRAF

RESEARCH

• RECRUITING

⚡ 8

Regorafenib With Low-dose Chemotherapies and Aspirin Followed by Standard Chemotherapies in Metastatic Colorectal Cancer

This clinical trial is evaluating the effectiveness of regorafenib combined with metronomic chemotherapies and low-dose aspirin as an induction therapy for patients with second-line metastatic colorectal cancer. The intervention involves administering regorafenib along with other drugs and assessing quality of life through questionnaires and blood samples. Currently in Phase 2 and Phase 3, the trial is actively recruiting participants, which means eligible patients may have the opportunity to receive this treatment approach. Eligibility criteria include specific biomarker requirements related to BRAF, RAS, and microsatellite status.

NCT: NCT05462613 · Enrollment: 446 · Sites: Besançon, France, Clermont-Ferrand, France, Créteil, France, Dijon, France, Lyon, France

MEDIUM

PHASE2, PHASE3

All Stages

HER2

TARGETED THERAPY

• RECRUITING

⚡ 8

DETERMINE Trial Treatment Arm 04: Trastuzumab in Combination With Pertuzumab in Adult, Paediatric and Teenage/Young Adult Patients With Cancers With HER2 Amplification or Activating Mutations

The DETERMINE Trial Treatment Arm 04 is investigating the efficacy of trastuzumab in combination with pertuzumab in adult, pediatric, and teenage/young adult patients with cancers that exhibit HER2 amplification or activating mutations. This combination of drugs is already approved as a standard treatment for adult patients with breast cancer, highlighting its established role in targeting HER2-related malignancies. Currently in Phase 2 and Phase 3, the trial is actively recruiting participants, which means eligible patients may have the opportunity to access this treatment while contributing to important research. Eligibility criteria include the presence of HER2 amplification or activating mutations, as determined by next-generation sequencing or other biomarker assessments.

NCT: NCT05786716 · Enrollment: 30 · Sites: Belfast, United Kingdom, Birmingham, United Kingdom, Bristol, United Kingdom, Cambridge, United Kingdom, Cardiff, United Kingdom

SIGNAL **PHASE3** **Stage IV** **BRAF** **RESEARCH** • ACTIVE_NOT_RECRUITING

8

Combination With Sintilimab and XELOX+Bevacizumab as 1st Line Therapy in RAS-mutant Metastatic Colorectal Cancer

This clinical trial is testing the efficacy of a combination therapy involving sintilimab, XELOX (oxaliplatin and capecitabine), and bevacizumab in patients with RAS-mutant metastatic colorectal cancer who have not received prior treatment. The intervention includes sintilimab, a PD-1 monoclonal antibody, which is being evaluated for its potential to enhance anti-tumor immunity when used alongside chemotherapy. This is a Phase 3 trial that is currently active but not recruiting, indicating that it is in the advanced stages of testing and may provide insights into treatment options for eligible patients. Eligibility criteria include the presence of specific biomarkers such as BRAF, RAS, and microsatellite status.

NCT: NCT05171660 · Enrollment: 446 · Sites: Hangzhou, China

SIGNAL **PHASE2, PHASE3** **Stage IV** **RAS** **RESEARCH** • RECRUITING

8

Study Evaluating the Tailored Management of Locally-advanced Rectal Carcinoma

This clinical trial is evaluating the tailored management of locally advanced rectal carcinoma for patients who may benefit from personalized treatment approaches. The intervention includes a modified FOLFIRINOX regimen for induction chemotherapy, early tumor response evaluation by MRI volumetry, radiochemotherapy, and radical proctectomy with total mesorectal excision, which are designed to improve surgical outcomes and reduce metastatic risks. Currently in Phase 2 and Phase 3, the trial is recruiting participants, which means eligible patients can join to potentially receive these interventions. Eligibility criteria include the presence of RAS biomarkers, which may help in assessing treatment response.

NCT: NCT04749108 · Enrollment: 1075 · Sites: Amiens, France, Annecy, France, Besançon, France, Bordeaux, France, Clermont-Ferrand, France

SIGNAL **PHASE2, PHASE3** **Stage III** **RAS** **RESEARCH** • RECRUITING

8

Adjuvant mFOLFIRINOX for High-risk Stage III Colon Cancer

This clinical trial is testing the effectiveness of mFOLFIRINOX compared to mFOLFOX6 as adjuvant treatment for patients with high-risk stage III colon cancer. The intervention involves two chemotherapy regimens, mFOLFIRINOX and mFOLFOX6, which are important for potentially improving disease-free survival outcomes. Currently in Phase 2 and Phase 3, the trial is actively recruiting participants, which means eligible patients may have the opportunity to receive one of the treatments while contributing to the research. Eligibility criteria include specific tumor characteristics, such as RAS biomarker status.

NCT: NCT05179889 · Enrollment: 308 · Sites: Daejeon, South Korea

SIGNAL **PHASE3** **Stage IV** **BRAF** **RESEARCH** • RECRUITING

8

Conversion Therapy of RAS/BRAF Wild-Type Colorectal Cancer Patients With Initially Unresectable Liver Metastases

This clinical trial is testing the effectiveness of two treatment combinations for patients with RAS/BRAF wild-type colorectal cancer who have initially unresectable liver metastases. The interventions being compared are modified FOLFOXIRI (mFOLFOXIRI) combined with cetuximab and mFOLFOXIRI combined with bevacizumab, as both combinations may improve response rates and the potential for surgical resection of liver metastases. This is a Phase 3 trial currently recruiting 508 participants, which means eligible patients may have the opportunity to receive one of the treatment regimens while contributing to important research outcomes. Eligibility criteria include confirmation of RAS/BRAF wild-type status and the presence of liver-only metastases, as determined by a local multidisciplinary team.

NCT: NCT04687631 · Enrollment: 508 · Sites: Shanghai, China

SIGNAL **PHASE3** **Stage IV** **KRAS** **RESEARCH** • ACTIVE_NOT_RECRUITING

8

Study to Evaluate the Efficacy of FOLFOX + Panitumumab Followed by FOLFIRI + Bevacizumab (Sequence 1) Versus FOLFOX + Bevacizumab Followed by FOLFIRI + Panitumumab (Sequence 2) in Untreated Patients With Wild-type RAS Metastatic, Primary Left-sided, Unresectable Colorectal Cancer

This clinical trial is evaluating the efficacy of two treatment sequences for untreated patients with wild-type RAS metastatic, primary left-sided, unresectable colorectal cancer. The interventions being tested are the FOLFOX regimen combined with either panitumumab or bevacizumab, followed by the FOLFIRI regimen with the alternate drug, to

determine which sequence is more effective. This is a Phase 3 trial that is currently active but not recruiting, indicating that it is in the final stages of testing before potential approval, and patients may not be able to enroll at this time. Eligibility criteria include having wild-type RAS status, which is essential for determining the appropriateness of the targeted therapies being studied.

NCT: NCT03635021 · Enrollment: 419 · Sites: Madrid, Spain

SIGNAL **PHASE3** **Stage IV** **RESEARCH** • ACTIVE_NOT_RECRUITING

⚡ 8

Consolidation Chemotherapy for Locally Advanced Mid or Low Rectal Cancer After Neoadjuvant Concurrent Chemoradiotherapy

This clinical trial is testing the efficacy and feasibility of consolidation chemotherapy for patients with locally advanced mid or low rectal cancer following neoadjuvant concurrent chemoradiotherapy. The intervention involves administering chemotherapy to evaluate its impact on achieving a pathologic complete response. Currently, the trial is in Phase 3 and is active but not recruiting, which means that while the study is ongoing, no new participants are being enrolled at this time. Eligibility criteria may include specific characteristics related to the cancer stage and treatment history.

NCT: NCT02843191 · Enrollment: 358 · Sites: Busan, South Korea, Cheonan, South Korea, Chuncheon, South Korea, Daegu, South Korea, Daejeon, South Korea

SIGNAL **PHASE3** **Stage IV** **KRAS** **RESEARCH** • RECRUITING

⚡ 8

Systemic Oxaliplatin or Intra-arterial Chemotherapy Combined With LV5FU2 +/- Irinotecan and an Target Therapy in First Line Treatment of Metastatic Colorectal Cancer Restricted to the Liver

This clinical trial is testing the effectiveness of systemic oxaliplatin versus intra-arterial chemotherapy combined with LV5FU2, with or without irinotecan and panitumumab, in patients with metastatic colorectal cancer restricted to the liver. The intervention involves administering oxaliplatin both intravenously and intra-arterially, along with other chemotherapy agents and targeted therapy, to potentially enhance treatment efficacy for patients with unresectable liver metastases. This is a Phase 3 trial currently recruiting 348 participants, which indicates that it is in the later stages of testing and may provide valuable information on treatment options for patients. Eligibility criteria include specific biomarker assessments for KRAS, NRAS, and RAS.

NCT: NCT02885753 · Enrollment: 348 · Sites: Angers, France, Antony, France, Avignon, France, Bayonne, France, Beauvais, France

SIGNAL **PHASE3** **All Stages** **MSI/MMR** **RESEARCH** • NOT_YET_RECRUITING

⚡ 8

Finding the Best Dose of Aspirin to Prevent Lynch Syndrome Cancers

This clinical trial is testing the effectiveness of different doses of enteric coated aspirin in preventing cancers associated with Lynch Syndrome, specifically for individuals who carry a germline pathological mismatch repair gene defect. The intervention involves comparing daily doses of 600mg, 300mg, and 100mg of aspirin to determine the optimal dose for cancer prevention. This is a Phase 3 trial that is currently not yet recruiting participants, meaning that patients will need to wait until recruitment begins to consider participation. Eligibility criteria include being a carrier of a germline pathological mismatch repair gene defect associated with Lynch Syndrome.

NCT: NCT02497820 · Enrollment: 1800 · Sites: Tel Aviv, Israel

MEDIUM **PHASE2** **Stage IV** **MSI/MMR** **TARGETED THERAPY** • ACTIVE_NOT_RECRUITING

⚡ 8

Study of Pembrolizumab (MK-3475) in Participants With Advanced Solid Tumors (MK-3475-158/KEYNOTE-158)

This clinical trial is testing the efficacy of pembrolizumab (MK-3475) in participants with advanced solid tumors who have progressed on standard of care therapy. The intervention, pembrolizumab, is a biological agent that targets the immune system to help fight cancer, which is significant for patients with limited treatment options. This is a Phase 2 trial that is currently active but not recruiting, indicating that while the study is ongoing, no new participants are being enrolled at this time. Eligibility criteria include specific biomarkers such as MSI, MSI-H, MMR, DMMR, TMB, and microsatellite instability, which may be relevant for patient selection.

NCT: NCT02628067 · Enrollment: 1609 ·

SIGNAL **PHASE3** **Stage III** **RESEARCH** • ACTIVE_NOT_RECRUITING

⚡ 8

Safety of a Boost (CXB or EBRT) in Combination With Neoadjuvant Chemoradiotherapy for Early Rectal Adenocarcinoma

This clinical trial is testing the safety and effectiveness of a boost using either Contact X-ray brachytherapy (CXB) or 3D conformal external beam radiation therapy (EBRT) in combination with neoadjuvant chemoradiotherapy for patients with early rectal adenocarcinoma. The intervention involves administering a radiation boost after standard treatment to potentially improve the rate of rectum preservation without the need for major surgery. This is a Phase 3 trial that is currently active but not recruiting, indicating that it is in the later stages of testing and may provide valuable insights for patients in the near future. Eligibility criteria include patients with cT2 or cT3a-b tumors less than 5 cm.

NCT: NCT02505750 · Enrollment: 148 · Sites: Besançon, France, Geneva, Switzerland, Guildford, United Kingdom, Hull, United Kingdom, Liverpool, United Kingdom

SIGNAL **PHASE3** **All Stages** **RESEARCH** • ACTIVE_NOT_RECRUITING

8

A Trial of Aspirin on Recurrence and Survival in Colon Cancer Patients

This clinical trial is testing the effectiveness of acetylsalicylic acid in reducing recurrence and improving survival in patients with colon cancer. The intervention involves administering acetylsalicylic acid compared to a placebo, which is significant for understanding its potential role in cancer management. This is a Phase 3 trial that is currently active but not recruiting, indicating that it is in the final stages of testing before potential approval. Eligibility criteria have not been specified, but the study involves a total enrollment of 770 participants.

NCT: NCT02301286 · Enrollment: 770 · Sites: Almelo, Netherlands, Amersfoort, Netherlands, Assen, Netherlands, Breda, Netherlands, Capelle aan den IJssel, Netherlands

SIGNAL **PHASE3** **All Stages** **RESEARCH** • ACTIVE_NOT_RECRUITING

8

The Northern-European Initiative on Colorectal Cancer

The Northern-European Initiative on Colorectal Cancer (NordICC) is a Phase 3 clinical trial designed to evaluate the effectiveness of colonoscopy screening in reducing colorectal cancer (CRC) incidence and mortality among individuals aged 55-64 years. The intervention involves a colonoscopy procedure, which is significant as it can detect and remove precursor lesions that may lead to CRC, potentially preventing the disease. Currently, the trial is active but not recruiting, indicating that it is ongoing with enrolled participants, and results will be assessed after 15 years of follow-up. Eligibility criteria include being part of the specified age group and drawn randomly from population registries in the participating Nordic countries, the Netherlands, and Poland.

NCT: NCT00883792 · Enrollment: 95000 · Sites: Boston, New York

SIGNAL **PHASE3** **Stage IV** **BRAF** **RESEARCH** • NOT_YET_RECRUITING

7

Pressurized Intraperitoneal Aerosolized Chemotherapy With Mitomycin for the Treatment of Unresectable Appendix or Colorectal Cancer With Peritoneal Metastases, The IMPACT Trial

This phase III clinical trial, known as the IMPACT Trial, is testing the effectiveness of pressurized intraperitoneal aerosolized chemotherapy (PIPAC) with mitomycin in patients with unresectable appendix or colorectal cancer that has metastasized to the peritoneal cavity. The intervention involves delivering chemotherapy as a pressurized mist directly into the abdominal cavity during a minimally invasive procedure, which may enhance drug absorption and distribution compared to standard chemotherapy regimens. The trial is currently not yet recruiting participants, which means patients will need to wait for enrollment to begin before they can consider participation. Eligibility criteria include specific biomarker assessments such as BRAF, RAS, mismatch repair status, and next-generation sequencing (NGS).

NCT: NCT07271355 · Enrollment: 129 · Sites: Duarte, Goodyear, Irvine, Newnan, Zion

MEDIUM **PHASE1, PHASE2** **Stage IV** **MSI/MMR** **TARGETED THERAPY** • RECRUITING

7

A Clinical Research Study Evaluating EIK1005, a Werner Helicase Inhibitor, as Monotherapy and in Combination With Pembrolizumab in Participants With Advanced Solid Tumors Including Microsatellite Instability High (MSI-H) Tumors

The EIK1005-002 clinical trial is evaluating the efficacy and safety of EIK1005, a Werner Helicase inhibitor, both as a monotherapy and in combination with pembrolizumab in participants with advanced solid tumors, including those with microsatellite instability high (MSI-H) tumors. The intervention involves administering EIK1005 to determine the most effective and safe dosage, as well as assessing its tolerability when combined with pembrolizumab. This trial is currently in Phase 1 and Phase 2 and is actively recruiting participants, which means eligible patients may have the opportunity to access these investigational treatments. Eligibility criteria include specific biomarkers such as MSI, MSI-H, MMR, DMMR, RAS, and microsatellite status.

MEDIUM

PHASE1

Stage IV

BRAF

TARGETED THERAPY

• RECRUITING

7

A Study to Investigate the Safety, Tolerability, Pharmacokinetics, and Anti-tumor Activity of CBI-1214 T Cell Engager in Participants With Advanced or Metastatic MSS/MSI-L Colorectal Cancer

This clinical trial is investigating the safety, tolerability, pharmacokinetics, and anti-tumor activity of CBI-1214 T Cell Engager in participants with advanced or metastatic Microsatellite Stable (MSS) or Microsatellite Instability Low (MSI-L) colorectal cancer. The intervention, CBI-1214, is a biological agent designed to engage T cells to target cancer cells, which is important for potentially improving treatment outcomes in this patient population. Currently in Phase 1 and actively recruiting, this trial aims to assess the safety and tolerability of CBI-1214 at various dose levels, which is crucial for determining the appropriate dosing for future studies. Eligibility criteria include specific biomarkers such as BRAF, MSI, MSI-H, MMR, DMMR, and HER2.

NCT: NCT07321106 · Enrollment: 80 · Sites: Fairfax, Grand Rapids, Los Angeles, San Antonio

MEDIUM

PHASE1

All Stages

MSI/MMR

TARGETED THERAPY

• RECRUITING

7

Phase I Study of [177Lu]Lu-DFC413 in Patients With Solid Tumors

This clinical trial is testing the safety and efficacy of the drugs 177Lu-DFC413 and 68Ga-NNS309 in patients aged 18 years and older with solid tumors. The intervention involves administering these drugs to evaluate their safety, tolerability, dosimetry, and preliminary efficacy, which is important for understanding their potential therapeutic benefits. Currently in Phase I and actively recruiting participants, this trial aims to assess the incidence and severity of dose-limiting toxicities associated with 177Lu-DFC413. Eligibility criteria include specific biomarkers such as MSI, MSI-H, MMR, DMMR, HER2, and microsatellite instability.

NCT: NCT07261631 · Enrollment: 180 · Sites: Essen, Germany, Haifa, Israel, Montreal, Canada, Odense C, Denmark, Singapore, Singapore

SIGNAL

PHASE3

Stage IV

RESEARCH

• RECRUITING

7

Modulated Electro-Hyperthermia in Combination With Multimodal Therapy for Locally Advanced Rectal Cancer

This clinical trial is testing the effectiveness of modulated electro-hyperthermia (mEHT) in combination with Total Neoadjuvant Therapy (TNT) for adult patients with locally advanced rectal adenocarcinoma. The intervention involves the use of the Oncotherm EHY-2030 device to apply hyperthermia, which may enhance tumor down-staging and improve pathological responses. Currently in Phase 3 and actively recruiting, this trial aims to determine if the addition of mEHT can significantly improve outcomes compared to TNT alone. Eligible participants include adults aged 20 years and older with specific stages of rectal cancer at high risk of recurrence.

NCT: NCT07372300 · Enrollment: 126 · Sites: Chiayi City, Taiwan

HIGH

PHASE3

All Stages

PHASE III TRIAL

• RECRUITING

7

Phase 3 Trial of eRapa in Patients With Familial Adenomatous Polyposis

This Phase 3 clinical trial is testing the efficacy of eRapa (encapsulated rapamycin) in patients diagnosed with Familial Adenomatous Polyposis (FAP). The intervention involves administering eRapa compared to a placebo to determine its impact on slowing disease progression and its safety profile. Currently, the trial is recruiting 168 participants, which means eligible patients can still join to contribute to the research. Eligibility criteria include a diagnosis of FAP, and participants will undergo regular check-ups and endoscopies to monitor their condition throughout the study.

NCT: NCT06950385 · Enrollment: 168 · Sites: Ann Arbor, Arcadia, Baltimore, Boston, Chicago

MEDIUM

NA

Stage IV

MSI/MMR

TARGETED THERAPY

• ACTIVE_NOT_RECRUITING

7

Reducing the Resection Range After Neoadjuvant Therapy for Locally Advanced Upper Rectal and Rectosigmoid Junction Cancers

This clinical trial is evaluating the safety and efficacy of reduced surgical resection margins in patients with locally advanced upper rectal and rectosigmoid junction cancers who have shown regression following neoadjuvant therapy. The intervention involves laparoscopic resection with reduced margins of 3 cm distally and 5 cm proximally, which is significant for potentially minimizing surgical impact while maintaining effective cancer treatment. The trial is currently active but not

recruiting, indicating that it is ongoing and may provide insights into surgical practices for eligible patients in the future. Eligibility criteria include specific biomarkers such as MSI, MSI-H, MMR, DMMR, RAS, and microsatellite status.

NCT: NCT07038122 · Enrollment: 874 · Sites: Guangzhou, China

SIGNAL **PHASE3** **All Stages** **KRAS** **RESEARCH** • **RECRUITING**

7

A Study of Tucidinostat in Combination With Sintilimab and Bevacizumab in MSS/pMMR Colorectal Cancer Patients

This clinical trial is testing the combination of tucidinostat, sintilimab, and bevacizumab in patients with MSS/pMMR colorectal cancer. The intervention aims to evaluate the efficacy and safety of this combination compared to fruquintinib monotherapy, which is significant for understanding potential treatment options. Currently in Phase 3 and actively recruiting, this trial may provide eligible patients with access to a novel treatment approach while contributing to the overall understanding of colorectal cancer therapies. Eligibility criteria include specific biomarkers such as KRAS, MSI, MMR, and RAS.

NCT: NCT06497985 · Enrollment: 430 · Sites: Guangzhou, China

SIGNAL **PHASE3** **All Stages** **RESEARCH** • **NOT_YET_RECRUITING**

7

Preoperative Chemoradiotherapy and Transanal Endoscopic Microsurgery Versus Ttransanal Endoscopic Microsurgery in T1 N0, M0 Rectal Cancer (TAUTEM-T1 Study)

the T1 N0, M0 rectal cancer stage. This trial is testing the combination of preoperative chemoradiotherapy using Capecitabine and 50.4 Gy of radiation followed by Transanal Endoscopic Microsurgery (TEM) against TEM alone. As a Phase 3 trial that is not yet recruiting, it aims to evaluate rectal preservation rates in patients with early-stage rectal cancer, which may influence treatment decisions and outcomes. Eligibility criteria include patients diagnosed with T1, N0, M0 rectal adenocarcinoma without poor prognostic factors.

NCT: NCT06450574 · Enrollment: 106 ·

SIGNAL **PHASE3** **All Stages** **RESEARCH** • **RECRUITING**

7

CompariSon Between the EuroPeAn and Japanese pathologiCal InvEstigation for Colon Cancer (SPACE)

The SPACE trial is testing the diagnostic accuracy of Japanese versus European pathological investigation methods for stage III colon cancer in patients. The intervention involves comparing two procedures: the Japanese pathological investigation, which includes surgeon involvement in lymph node extraction and specimen assessment, and the European method, which primarily relies on pathologists. This is a Phase 3 trial currently recruiting 430 participants, which means patients may have the opportunity to contribute to important findings regarding diagnostic practices. Eligibility criteria include patients diagnosed with stage III colon cancer, although specific biomarker requirements are not mentioned.

NCT: NCT06119867 · Enrollment: 430 · Sites: Moscow, Russia

SIGNAL **PHASE3** **All Stages** **RAS** **RESEARCH** • **RECRUITING**

7

Regorafenib Combined With Irinotecan Drug-Eluting Beads for Colorectal Cancer Liver Metastases

This clinical trial is testing the efficacy and safety of regorafenib combined with Irinotecan Drug-Eluting Beads in patients with colorectal cancer liver metastases who have failed first- and second-line standard chemotherapy. The intervention involves the use of regorafenib and DIBIRI (drug-eluting beads), which is significant as it explores a combination treatment approach for this patient population. This is a Phase 3 trial currently recruiting 126 participants, which means that eligible patients may have the opportunity to access this investigational treatment while contributing to important research. Eligibility criteria include the presence of RAS biomarkers.

NCT: NCT05794971 · Enrollment: 126 · Sites: Guangdong, China

SIGNAL **PHASE3** **All Stages** **NGS** **RESEARCH** • **RECRUITING**

7

Induction Chemotherapy for Locally Recurrent Rectal Cancer

This clinical trial is testing the effectiveness of induction chemotherapy followed by neoadjuvant chemoradiotherapy and surgery in patients with locally recurrent rectal cancer. The intervention involves a combination of drugs, radiation, and

surgical procedures, which may improve surgical outcomes by increasing the likelihood of achieving clear resection margins. Currently in phase III and actively recruiting participants, this trial aims to enroll 364 patients, which may provide valuable insights into treatment options for this patient population. Eligibility criteria include the requirement for next-generation sequencing (NGS) biomarkers.

NCT: NCT04389086 · Enrollment: 364 · Sites: Amsterdam, Netherlands, Eindhoven, Netherlands, Ghent, Belgium, Gothenburg, Sweden, Groningen, Netherlands

MEDIUM

NA

All Stages

TARGETED THERAPY

• ACTIVE_NOT_RECRUITING

7

Targeted Agent Evaluation in Digestive Cancers in China Based on Molecular Characteristics

This clinical trial is evaluating the effectiveness of various targeted agents in patients with previously treated and refractory gastrointestinal tumors, including colorectal cancer. The intervention involves multiple inhibitors and therapies, such as FGFR, IDH1, HER2, PARP, BRAF, MEK, immune checkpoint inhibitors, and others, aimed at providing individualized treatment based on molecular characteristics. The trial is currently active but not recruiting, indicating that while it is ongoing, no new participants are being enrolled at this time. Eligibility criteria may include specific molecular biomarkers identified through tumor DNA sequencing, which will guide the selection of targeted therapies for each patient.

NCT: NCT04584008 · Enrollment: 600 · Sites: Beijing, China

SIGNAL

PHASE2, PHASE3

Stage II

RAS

RESEARCH

• ACTIVE_NOT_RECRUITING

7

Duloxetine to Prevent Oxaliplatin-Induced Peripheral Neuropathy in Patients With Stage II-III Colorectal Cancer

This clinical trial is testing the effectiveness of duloxetine in preventing oxaliplatin-induced peripheral neuropathy in patients with stage II-III colorectal cancer. The intervention involves administering duloxetine, which may help alleviate pain, tingling, and numbness associated with oxaliplatin treatment by increasing certain brain chemicals. Currently in phase II/III and marked as active but not recruiting, this trial is significant for patients as it explores a potential method to improve their quality of life during chemotherapy. Eligibility criteria include the presence of RAS biomarkers.

NCT: NCT04137107 · Enrollment: 220 · Sites: Aberdeen, Akron, Albany, Albuquerque, Allentown

SIGNAL

PHASE3

All Stages

RAS

RESEARCH

• RECRUITING

7

Surgery Plus Chemo Versus Chemoradiotherapy Followed by Surgery Plus Chemo for Locally Recurrent Rectal Cancer

The JCOG1801 trial is testing the effectiveness of preoperative chemoradiotherapy followed by surgery and adjuvant chemotherapy in patients with locally recurrent rectal cancer. The intervention involves a combination of chemotherapy, preoperative radiotherapy, and surgical procedures, aiming to improve local relapse-free survival compared to standard treatment. This is a Phase 3 trial currently recruiting 110 participants, which means eligible patients may have the opportunity to receive a potentially more effective treatment approach. Eligibility criteria include the presence of RAS biomarkers.

NCT: NCT04288999 · Enrollment: 110 · Sites: Chiba, Japan, Gifu, Japan, Hidaka, Japan, Hirakata, Japan, Hiroshima, Japan

SIGNAL

PHASE3

All Stages

RAS

RESEARCH

• RECRUITING

7

Performance of SGM-101 for the Delineation of Primary and Recurrent Tumor and Metastases in Patients Undergoing Surgery for Colorectal Cancer

This clinical trial is testing the performance of SGM-101, an intraoperative imaging agent, in patients undergoing surgery for colorectal cancer. The intervention involves using SGM-101 to enhance the visualization of primary and recurrent tumors and metastases compared to standard "white light" visualization, which is important for improving surgical outcomes. This is a Phase 3 trial currently recruiting 300 participants, indicating that it is in the later stages of testing and may provide valuable information on the effectiveness of SGM-101 for surgical procedures. Eligibility criteria include the presence of RAS biomarkers.

NCT: NCT03659448 · Enrollment: 300 · Sites: Boston, Duarte, La Jolla, Philadelphia, Weston

SIGNAL

PHASE3

All Stages

RESEARCH

• ACTIVE_NOT_RECRUITING

7

Effect of EPA-FFA on Polypectomy in Familial Adenomatous Polyposis

This clinical trial is testing the effect of eicosapentaenoic acid free fatty acid (EPA-FFA) on polypectomy outcomes in patients with familial adenomatous polyposis (FAP). The intervention involves administering EPA-FFA gastro-resistant capsules compared to a placebo, which is important for assessing its potential impact on the number of polypectomies required. This is a Phase 3 trial that is currently active but not recruiting, indicating that it is in the later stages of testing and may provide significant insights into treatment efficacy for patients. Eligibility criteria may include a diagnosis of familial adenomatous polyposis, although specific biomarker requirements are not detailed.

NCT: NCT03806426 · Enrollment: 204 · Sites: Bologna, Italy

MEDIUM

PHASE1, PHASE2

Stage IV

KRAS

LIQUID BIOPSY

• RECRUITING

7

A Study of Bispecific Antibody MCLA-158 in Patients With Advanced Solid Tumors

This clinical trial is testing the bispecific antibody MCLA-158 in patients with advanced solid tumors, specifically focusing on metastatic colorectal cancer (mCRC) and other solid tumor indications. The intervention includes MCLA-158 as a single agent and in combination with pembrolizumab, FOLFIRI, and FOLFOX, which is important for evaluating its safety and effectiveness in various treatment settings. Currently in Phase 1/2 and actively recruiting, this trial aims to determine the recommended Phase II dose and assess the drug's safety, tolerability, and anti-tumor activity. Eligibility criteria include specific biomarkers such as KRAS, NRAS, BRAF, HER2, and others, which may influence treatment response.

NCT: NCT03526835 · Enrollment: 523 · Sites: Boston, Cleveland, Dallas, Fort Myers, Houston

SIGNAL

PHASE3

All Stages

RESEARCH

• ACTIVE_NOT_RECRUITING

7

EPA for Metastasis Trial 2

The EPA for Metastasis Trial 2 is testing the efficacy of Icosapent Ethyl, a pure omega-3 fatty acid derived from fish oil, in patients with liver metastases from bowel cancer. The trial aims to determine if taking this supplement before and after liver surgery can improve progression-free survival compared to a placebo. Currently in Phase 3 and active but not recruiting, this study involves 418 participants and seeks to provide insights into the potential benefits of Icosapent Ethyl in preventing disease recurrence. Eligibility criteria include having liver metastases from bowel cancer, with a target enrollment of 448 individuals.

NCT: NCT03428477 · Enrollment: 418 · Sites: Aintree, United Kingdom, Basingstoke, United Kingdom, Birmingham, United Kingdom, Cambridge, United Kingdom, Cardiff, United Kingdom

SIGNAL

PHASE2, PHASE3

All Stages

RESEARCH

• ACTIVE_NOT_RECRUITING

7

Perioperative Systemic Therapy for Isolated Resectable Colorectal Peritoneal Metastases

This clinical trial is testing the effectiveness of perioperative systemic therapy combined with cytoreductive surgery and HIPEC in patients with isolated resectable colorectal peritoneal metastases. The intervention includes various combinations of chemotherapy regimens (CAPOX, FOLFOX, FOLFIRI) with bevacizumab, alongside the experimental procedure of cytoreductive surgery with HIPEC. Currently in Phase II-III and marked as active but not recruiting, this trial aims to determine if the combination therapy improves overall survival compared to surgery alone. Eligibility criteria include having isolated resectable colorectal peritoneal metastases, although specific biomarker requirements are not detailed.

NCT: NCT02758951 · Enrollment: 358 · Sites: Amsterdam, Netherlands, Eindhoven, Netherlands, Genk, Belgium, Groningen, Netherlands, Nieuwegein, Netherlands

MEDIUM

PHASE2

Stage IV

BRAF

TARGETED THERAPY

• NOT_YET_RECRUITING

6

Testing Ivonescimab in Combination With Chemotherapy in Patients With Metastatic Colorectal Cancers Without Liver Metastases

This clinical trial is testing the efficacy of ivonescimab in combination with the FOLFIRI chemotherapy regimen in patients with metastatic colorectal cancer (mCRC) who do not have liver metastases. The intervention involves administering ivonescimab alongside FOLFIRI, comparing its effectiveness to the standard treatment of FOLFIRI combined with bevacizumab, with a focus on improving progression-free survival (PFS). This is a Phase 2 trial that is not yet recruiting, meaning patients will need to wait until enrollment begins to participate. Eligibility criteria include specific biomarkers such as BRAF, MSI, MMR, and RAS status, which will determine patient inclusion in the study.

NCT: NCT07359456 · Enrollment: 130 · Sites: Auxerre, France, Beuvry, France, Caen, France, Calais, France, Compiègne, France

MEDIUM

PHASE2

Stage IV

MSI/MMR

TARGETED THERAPY

• RECRUITING

6

A Single-arm Phase II Clinical Study of Camrelizumab Combined With Long-course Chemoradiotherapy for Total Neoadjuvant Therapy in Locally Advanced Low pMMR/MSS Rectal Cancer

This clinical trial is testing the combination of camrelizumab with long-course chemoradiotherapy as total neoadjuvant therapy for patients with locally advanced low pMMR/MSS rectal cancer. The intervention involves the use of camrelizumab, a drug that may enhance the effectiveness of standard treatment by targeting the immune system. This is a Phase II trial currently recruiting participants, which means that eligible patients may have the opportunity to receive this treatment while contributing to the understanding of its efficacy. Eligibility criteria include specific biomarker assessments such as MSI, MMR, and RAS status.

NCT: NCT07527026 · Enrollment: 44 · Sites: Beijing, China

MEDIUM

PHASE2

Stage IV

MSI/MMR

TARGETED THERAPY

• NOT_YET_RECRUITING

6

Neoadjuvant Therapy for Locally Advanced Low Rectal Cancer (SMARTi-RC01)

The SMARTi-RC01 clinical trial is investigating the effectiveness of combining serplulimab, a PD-1 inhibitor, with bevacizumab, short-course radiotherapy, and chemotherapy in adults with locally advanced mid-to-low rectal cancer. This trial aims to determine if the addition of bevacizumab enhances the complete remission rate compared to serplulimab combined with chemotherapy and radiotherapy alone, while also assessing the safety of these treatments. Currently in Phase 2 and not yet recruiting, this trial will eventually enroll 138 participants, providing insights into treatment efficacy and safety for patients with this specific cancer type. Eligibility criteria include specific biomarkers such as MSI, MSI-H, MMR, DMMR, and microsatellite status.

NCT: NCT07134101 · Enrollment: 138 ·

MEDIUM

PHASE2

Stage IV

MSI/MMR

TARGETED THERAPY

• ACTIVE_NOT_RECRUITING

6

The Efficacy and Safety of Neoadjuvant Therapy With Iparomlimab and Tuvonralimab in Locally Advanced MSI-H/dMMR Colorectal Cancer: An Prospective, Single-Arm Study (Neo-IT)

This clinical trial is testing the efficacy and safety of neoadjuvant therapy with Iparomlimab and Tuvonralimab in patients with locally advanced colorectal cancer that is microsatellite instability-high (MSI-H) or mismatch repair deficient (dMMR). The intervention involves the use of these two immunotherapy drugs, which may provide significant benefits in the preoperative treatment of this specific cancer type. This is a Phase 2 trial that is currently active but not recruiting, indicating that patients are not being enrolled at this time, but results may inform future treatment options. Eligibility criteria include specific biomarkers such as MSI, MSI-H, MMR, dMMR, and RAS status.

NCT: NCT07160647 · Enrollment: 29 · Sites: Hangzhou, China

MEDIUM

PHASE1

Stage IV

MSI/MMR

TARGETED THERAPY

• RECRUITING

6

A First in Human Study of ALX2004 With Advanced or Metastatic Selected Solid Tumors

This clinical trial is testing ALX2004, an antibody drug conjugate targeting EGFR, in patients with advanced or metastatic selected solid tumors. The intervention involves administering ALX2004 to evaluate its safety and tolerability, specifically focusing on the incidence of dose limiting toxicities (DLTs). Currently in Phase 1 and actively recruiting, this trial aims to gather initial data on the drug's effects, which is crucial for determining its potential for further development. Eligibility criteria include specific biomarkers such as MSI, MSI-H, MMR, and DMMR.

NCT: NCT07085091 · Enrollment: 170 · Sites: Fairfax, Grand Rapids, Houston, Portland, Spokane

MEDIUM

N/A

Stage III

MSI/MMR

TARGETED THERAPY

• RECRUITING

6

Exploring the Treatment Duration of PD-1 Neoadjuvant Therapy in Stage II-III dMMR Rectal Cancer

This clinical trial is investigating the optimal treatment duration of PD-1 neoadjuvant therapy for patients with stage II-III dMMR rectal cancer. Participants will receive preoperative monotherapy with PD-1 antibodies, which is important for assessing its effectiveness in achieving a pathological complete response (pCR). The trial is currently recruiting 100 participants and aims to evaluate outcomes such as pCR rates, adverse reactions, and three-year event-free survival. Eligibility criteria include biomarkers such as MSI, MSI-H, MMR, DMMR, and RAS.

MEDIUM

PHASE1

Stage IV

BRAF

TARGETED THERAPY

• RECRUITING

6

A First-in-Human Escalation and Expansion Study of Patients With Advanced Solid Tumors

This clinical trial is testing the investigational drug JR8603 in patients with advanced solid tumors who have not responded to prior treatments. The study aims to evaluate the safety and effectiveness of JR8603, which is significant for patients seeking alternative options after standard therapies have failed. Currently in Phase 1 and actively recruiting participants, this trial provides an opportunity for patients to access a potential treatment option while contributing to research. Eligibility criteria include specific biomarker requirements such as BRAF, MSI, MSI-H, MMR, DMMR, HER2, and RAS.

NCT: NCT06783569 · Enrollment: 94 · Sites: Harbin, China, Shenyang, China

MEDIUM

PHASE2

Stage IV

MSI/MMR

TARGETED THERAPY

• RECRUITING

6

Phase 2 Study of SR-8541A in Combination With Botensilimab and Balstilimab in Subjects With Refractory Metastatic Microsatellite Stable Colorectal Cancer (MSS-CRC)

This clinical trial is testing the combination of SR-8541A, botensilimab, and balstilimab in patients with refractory metastatic microsatellite stable colorectal cancer (MSS-CRC). The intervention involves administering SR-8541A orally alongside the other two drugs to assess their safety, tolerability, and efficacy, as well as to determine the optimal dose for further studies. Currently, this is a Phase 2 trial that is actively recruiting participants, which means eligible patients may have the opportunity to receive this treatment while contributing to the research. Eligibility criteria include specific biomarkers related to microsatellite stability and mismatch repair status.

NCT: NCT06589440 · Enrollment: 70 · Sites: Austin, Dallas, Houston, Morristown, Seattle

MEDIUM

PHASE2

Stage IV

MSI/MMR

TARGETED THERAPY

• RECRUITING

6

Immune Checkpoint Inhibitors for Organ Preservation in Non-metastatic dMMR/MSI-H Gastric or Colon Cancers

This clinical trial is testing the effectiveness of immune checkpoint inhibitors for organ preservation in patients with non-metastatic dMMR/MSI-H gastric or colon cancers. The intervention includes PD1/PDL1 antibody monotherapy and a combination of intensive immunotherapy with anti-VEGF treatment, both administered for up to 24 weeks, alongside the option of radical surgery. Currently in Phase 2 and actively recruiting, this trial aims to assess the organ preservation rate, which is a critical outcome for patients considering treatment options. Eligibility criteria include specific biomarkers such as MSI, MSI-H, MMR, DMMR, and RAS, as well as NGS testing.

NCT: NCT06580574 · Enrollment: 38 · Sites: Beijing, China

SIGNAL

PHASE3

All Stages

RESEARCH

• RECRUITING

6

Reducing Inflammatory Syndrome in Surgery

The RISIS-CR trial is testing the effectiveness of alpha-ketoglutarate (AKG) in reducing postoperative inflammatory responses in patients undergoing colorectal cancer surgery. The intervention involves administering AKG as a dietary supplement to patients identified as high inflammatory responders, with the aim of minimizing complications and improving recovery outcomes. This is a Phase 3 trial currently recruiting 80 participants, which means eligible patients may have the opportunity to receive a potentially beneficial treatment while contributing to important research. Eligibility criteria include the identification of patients based on their inflammatory responsiveness prior to surgery.

NCT: NCT06646809 · Enrollment: 80 · Sites: Singapore, Singapore

MEDIUM

PHASE1, PHASE2

Stage IV

MSI/MMR

TARGETED THERAPY

• RECRUITING

6

Phase I/II Study of the Combination Immunotherapy Regimen: SX-682, TriAdeno Vaccine, Retifanlimab and IL-15 Agonist N-803 (STAR15) for Metastatic Colorectal Cancer (mCRC)

This clinical trial is testing a combination immunotherapy regimen for adults with metastatic colorectal cancer (mCRC). The intervention includes the drugs retifanlimab, SX-682, and N-803, along with the TriAdeno vaccine, aiming to evaluate their safety profiles in treating mCRC. Currently in Phase I/II and actively recruiting, this trial offers participants the opportunity to access experimental therapies that may not be available outside of a clinical trial setting. Eligible participants must be 18 years or older and diagnosed with mCRC, with specific biomarker criteria including MSI, MSI-H, MMR, DMMR, RAS, and microsatellite status.

MEDIUM

PHASE2

Stage IV

MSI/MMR

TARGETED THERAPY

• RECRUITING

6

Liposomal Irinotecan and Leucovorin/5-fluorouracil Plus Bevacizumab in Metastatic Colorectal Cancer

This clinical trial is testing the efficacy and safety of liposomal irinotecan combined with 5-fluorouracil, leucovorin, and bevacizumab as a second-line therapy for patients with metastatic colorectal cancer. The intervention involves a combination of these drugs, which is important for potentially improving treatment outcomes in this patient population. Currently, the trial is in Phase 2 and is actively recruiting participants, indicating that patients may have the opportunity to receive this treatment while the study is ongoing. Eligibility criteria include specific biomarkers such as MSI, MSI-H, MMR, DMMR, and RAS.

NCT: NCT06184698 · Enrollment: 173 · Sites: Shijiazhuang, China

MEDIUM

PHASE2

Stage IV

BRAF

LIQUID BIOPSY

• RECRUITING

6

Cetuximab Plus Irinotecan in Patients With NeoRAS Wild-type Metastatic Colorectal Cancer In Third-line Therapy

This clinical trial is testing the efficacy and safety of cetuximab plus irinotecan in patients with NeoRAS wild-type metastatic colorectal cancer (mCRC) who are receiving third-line therapy. The intervention involves the combination of cetuximab, a monoclonal antibody, and irinotecan, a chemotherapy drug, which may provide a targeted approach for this specific patient population. Currently in phase II and actively recruiting, this trial aims to assess the objective response rate (ORR) to the treatment. Eligibility criteria include specific biomarker requirements such as BRAF, MSI, MSI-H, MMR, DMMR, CTDNA, and RAS status.

NCT: NCT05962502 · Enrollment: 54 · Sites: Guangzhou, China

MEDIUM

NA

Stage III

MSI/MMR

TARGETED THERAPY

• RECRUITING

6

MRD-guided Deferred Adjuvant Therapy in Resectable Early-stage Colon Cancer

This clinical trial is testing the use of minimal residual disease (MRD) status, detected by circulating tumor DNA (ctDNA), to guide post-surgery adjuvant therapy in patients with resectable early-stage colon cancer. The intervention involves comparing standard CAPEOX adjuvant therapy, deferred CAPEOX adjuvant therapy, and mFOLFOXIRI intensive adjuvant therapy based on MRD status, which may help tailor treatment to individual patient needs. The trial is currently in the recruiting phase, meaning eligible patients can still enroll to participate in this study. Eligibility criteria include having high-risk stage II or low-risk stage III colon cancer and specific biomarker assessments such as MSI, MMR, and RAS status.

NCT: NCT06204484 · Enrollment: 349 · Sites: Guangzhou, China

MEDIUM

PHASE1, PHASE2

Stage IV

RAS

TARGETED THERAPY

• ACTIVE_NOT_RECRUITING

6

Study of JK08 in Patients with Unresectable Locally Advanced or Metastatic Cancer

This clinical trial is evaluating the safety and efficacy of JK08 in patients with unresectable locally advanced or metastatic cancer. The intervention involves administering JK08, along with Pembrolizumab and Lenvatinib, to assess its effects on cancer progression and patient outcomes. Currently in Phase 1/2 and marked as active but not recruiting, this trial indicates that patients may not be able to enroll at this time but can learn about the study's findings as they become available. Eligibility criteria include specific biomarkers such as HER2 and RAS.

NCT: NCT05620134 · Enrollment: 263 · Sites: Barcelona, Spain, Brussels, Belgium, Edegem, Belgium, Ghent, Belgium, Madrid, Spain

SIGNAL

PHASE3

All Stages

RESEARCH

• ACTIVE_NOT_RECRUITING

6

Chemotherapy Followed by Pelvic Reirradiation Versus Chemotherapy Alone as Pre-operative Treatment for Locally Recurrent Rectal Cancer

The GRECCAR 15 trial is testing the efficacy of neoadjuvant chemotherapy followed by pelvic reirradiation compared to neoadjuvant chemotherapy alone in patients with locally recurrent rectal cancer who have previously undergone pelvic radiotherapy. The intervention involves administering six cycles of the chemotherapy regimen FOLFIRINOX, followed by radiochemotherapy and subsequent surgery, which is significant due to the increased risk of non-curative resection in this

patient population. This is a Phase 3 trial that is currently active but not recruiting, indicating that it is in the advanced stages of evaluation and may provide insights into treatment options for patients with this specific condition. Eligibility criteria include patients with a history of pelvic radiotherapy for primary rectal cancer and locally recurrent disease.

NCT: NCT03879109 · Enrollment: 58 · Sites: Avignon, France, Bordeaux, France, Grenoble, France, Lille, France, Lyon, France

MEDIUM

PHASE2

Stage III

MSI/MMR

TARGETED THERAPY

• RECRUITING

6

Neoadjuvant Immune Checkpoint Inhibition and Novel IO Combinations in Early-stage Colon Cancer

This clinical trial is testing the effectiveness of neoadjuvant immune checkpoint inhibition and novel immunotherapy combinations in patients with early-stage colon cancer (stage 1-3 adenocarcinoma) who do not have distant metastases. The intervention includes a combination of drugs such as Nivolumab, Ipilimumab, Celecoxib, BMS-986253, and BMS-986016, which are being evaluated for their safety and potential to improve treatment outcomes before surgical resection. This is a Phase 2 trial that is currently recruiting participants, indicating that eligible patients may have the opportunity to receive this investigational treatment in the lead-up to surgery. Eligibility criteria include specific biomarker requirements related to MMR and DMMR status.

NCT: NCT03026140 · Enrollment: 353 · Sites: Amsterdam, Netherlands, Eindhoven, Netherlands, Haarlem, Netherlands, Hilversum, Netherlands, The Hague, Netherlands

MEDIUM

PHASE2

Stage IV

BRAF

TARGETED THERAPY

• NOT_YET_RECRUITING

5

Becotatug Vedotin Combined With Cetuximab in the Later-line Treatment of Metastatic RAS Wild-type Colorectal Cancer

This clinical trial is testing the combination of becotatug vedotin and cetuximab for patients with metastatic RAS wild-type colorectal cancer who have already received prior treatments. The intervention involves the use of becotatug vedotin, a drug that targets cancer cells, alongside cetuximab, which is designed to inhibit tumor growth. This is a Phase 2 trial that is currently not yet recruiting, meaning patients cannot enroll at this time. Eligibility criteria include specific biomarker assessments such as BRAF, MSI, MMR, and RAS status.

NCT: NCT07490119 · Enrollment: 31 ·

MEDIUM

PHASE2

All Stages

MSI/MMR

TARGETED THERAPY

• NOT_YET_RECRUITING

5

FULIRI Plus Targeted Therapy for First-line Conversion Therapy of Colorectal Cancer Liver Metastases.

This clinical trial is testing the safety and efficacy of the FULIRI chemotherapy regimen combined with targeted therapy (bevacizumab/cetuximab) as first-line conversion therapy for patients with colorectal cancer and liver metastases. The intervention involves administering the FULIRI combination to evaluate its effectiveness in achieving an objective response rate (ORR) and facilitating potential surgical options. This is a Phase II trial that is currently not yet recruiting, meaning patients will need to wait until enrollment begins to participate. Eligibility criteria include specific biomarkers such as MSI, MSI-H, MMR, DMMR, and others, which may be required for participation.

NCT: NCT07515963 · Enrollment: 24 · Sites: Wuhan, China

MEDIUM

PHASE1, PHASE2

Stage IV

MSI/MMR

TARGETED THERAPY

• NOT_YET_RECRUITING

5

LPM6690176 in Combination With Chemotherapy and Bevacizumab in Metastatic Colorectal Cancer Patients With RAS Mutation

This clinical trial is testing the combination of LPM6690176 with chemotherapy and Bevacizumab in patients with RAS mutant metastatic colorectal cancer (mCRC). The intervention involves LPM6690176, a drug being evaluated for its safety and tolerability alongside standard treatments, which is important for determining its potential role in managing this specific cancer type. The trial is currently in Phase 1b and Phase 2 and is not yet recruiting participants, meaning that patients will need to wait until recruitment begins to consider participation. Eligibility criteria include the presence of specific biomarkers such as RAS mutations and mismatch repair status.

NCT: NCT07391566 · Enrollment: 99 · Sites: Beijing, China

MEDIUM

NA

All Stages

TARGETED THERAPY

• RECRUITING

5

Effect of Meal Timing During Cancer Treatment in Patients in Alaska: A Randomized Clinical Trial

This clinical trial is testing the effect of meal timing during cancer treatment in patients with rectal or breast cancer receiving neoadjuvant therapy at the Alaska Native Medical Center. The intervention includes time-restricted eating, health coaching, and biospecimen collection, aiming to improve therapeutic response and metabolic health. The trial is currently in the recruiting phase, which means eligible patients can still enroll to participate in the study. Eligibility criteria include having HER2 biomarkers and being part of the specified patient population at the ANMC.

NCT: NCT06802172 · Enrollment: 100 · Sites: Anchorage

MEDIUM

PHASE1, PHASE2

Stage III

KRAS

TARGETED THERAPY

• RECRUITING

5

KRAS Neoantigen Nanovaccine as Adjuvant Therapy for Colorectal Cancer/Pancreatic Cancer

This clinical trial is testing the KRAS Neoantigen Nanovaccine as adjuvant therapy for patients with colorectal or pancreatic cancer who have KRAS mutations and are at high risk of recurrence. The intervention involves a nanovaccine constructed from the bacterial membranes of an engineered *Lactococcus lactis* strain, which expresses KRAS antigenic peptides, aiming to enhance the immune response against cancer cells. Currently in Phase 1 and Phase 2, the trial is recruiting participants, which means eligible patients can join to help evaluate the safety and effectiveness of this treatment. Eligibility criteria include the presence of specific biomarkers such as KRAS, BRAF, MSI, and MMR status.

NCT: NCT07353645 · Enrollment: 49 · Sites: Nanjing, China

MEDIUM

PHASE2

Stage IV

BRAF

TARGETED THERAPY

• RECRUITING

5

The Safety and Efficacy of Cetuximab Beta Plus Fruquintinib With or Without Immune Checkpoint Inhibitors in First-line Treatment of RAS/BRAF Wild Type Unresectable Metastatic Colorectal Cancer

This clinical trial is testing the safety and efficacy of Cetuximab Beta combined with Fruquintinib, with or without immune checkpoint inhibitors, in patients with RAS/BRAF wild type unresectable metastatic colorectal cancer. The intervention involves targeted therapies that aim to improve treatment outcomes while potentially avoiding the side effects associated with traditional chemotherapy. This is a Phase 2 trial currently recruiting 70 participants, which means eligible patients may have the opportunity to receive these investigational treatments as part of their first-line therapy. Eligibility criteria include specific biomarker requirements such as BRAF, MSI, MSI-H, MMR, DMMR, and RAS status.

NCT: NCT07257653 · Enrollment: 70 · Sites: Hangzhou, China

MEDIUM

PHASE2

Stage II

MSI/MMR

TARGETED THERAPY

• RECRUITING

5

Immunotherapy (Toripalimab) for Reducing Recurrence Risk After Surgery for Mismatch Repair Deficient Stage IIB, IIC, or III Colon Cancer

This phase II clinical trial is evaluating the effectiveness of immunotherapy with toripalimab in reducing the risk of cancer recurrence after surgery in patients with mismatch repair deficient stage IIB, IIC, or III colon cancer. The intervention involves several procedures, including biopsy, biospecimen collection, colonoscopy, and computed tomography, which are essential for assessing treatment response and monitoring disease status. Currently, the trial is in the recruiting phase, meaning eligible patients can still enroll to participate in the study. Eligibility criteria include specific biomarkers such as MSI, MSI-H, MMR, DMMR, RAS, and others related to mismatch repair.

NCT: NCT07140679 · Enrollment: 40 · Sites: Atlanta, Johns Creek

MEDIUM

PHASE1

Stage IV

MSI/MMR

TARGETED THERAPY

• RECRUITING

5

Clinical Trial of an Anti-cancer Drug, CA-4948 (Emavusertib), in Combination With Chemotherapy Treatment (FOLFOX Plus Bevacizumab) in Metastatic Colorectal Cancer

This clinical trial is testing the anti-cancer drug CA-4948 (Emavusertib) in combination with the chemotherapy regimen FOLFOX (fluorouracil, leucovorin, and oxaliplatin) plus bevacizumab in patients with metastatic colorectal cancer. The intervention aims to evaluate the safety and determine the optimal dose of CA-4948, which may inhibit tumor cell growth by blocking specific enzymes necessary for cell proliferation, while the chemotherapy and bevacizumab work to kill cancer cells and prevent tumor blood vessel formation. This is a Phase 1 trial currently recruiting participants, which means it is focused on assessing safety and dosage rather than efficacy, and patients may have the opportunity to access this investigational treatment. Eligibility criteria include specific biomarkers such as MSI, MSI-H, MMR, DMMR, RAS, and microsatellite status.

NCT: NCT06696768 · Enrollment: 24 · Sites: City of Saint Peters, Creve Coeur, Fairway, Gainesville, Kansas City

MEDIUM

EARLY_PHASE1

Stage IV

MSI/MMR

TARGETED THERAPY

• RECRUITING

5

A Dose-Expansion Trial of Intravenous HNF4 α srRNA for Unresectable or Metastatic Colorectal Cancer

This clinical trial is testing the safety and efficacy of intravenous HNF4 α srRNA (CD-GA-102) in patients with unresectable locally advanced or metastatic colorectal cancer. The intervention involves administering CD-GA-102 as a monotherapy or in combination with immunotherapy and other systemic treatments, which is important for understanding its potential role in managing this cancer type. Currently in the early phase 1 stage and actively recruiting, this trial aims to gather data on the objective response rate, which may inform future treatment options for patients. Eligibility criteria include specific biomarkers such as MSI, MSI-H, MMR, DMMR, RAS, and microsatellite status.

NCT: NCT07050394 · Enrollment: 20 · Sites: Shanghai, China

MEDIUM

PHASE1, PHASE2

Stage IV

BRAF

TARGETED THERAPY

• RECRUITING

5

Glumetinib Combined With Fruquintinib in the Treatment of MET Amplification or Protein Overexpression in Third-Line Unresectable Metastatic Colorectal Cancer

This clinical trial is testing the combination of Glumetinib and Fruquintinib for patients with third-line unresectable metastatic colorectal cancer who have MET amplification or protein overexpression. The intervention involves the use of these two drugs to evaluate their efficacy and safety in this specific patient population. Currently, the trial is in Phase 1 and Phase 2 and is actively recruiting participants, which means eligible patients may have the opportunity to receive this treatment while contributing to the research. Eligibility criteria include specific biomarkers such as BRAF, MSI, MSI-H, MMR, DMMR, and microsatellite status, as well as next-generation sequencing (NGS) results.

NCT: NCT06980532 · Enrollment: 42 · Sites: Wuhan, China

MEDIUM

PHASE2

Stage IV

MSI/MMR

TARGETED THERAPY

• RECRUITING

5

Y-90, Capecitabine, and Atezolizumab for Oligometastatic CRC

The BRUOG-430 trial is evaluating the combination of yttrium-90 radioembolization, capecitabine, and atezolizumab for patients with unresectable colorectal cancer liver metastases who have previously received two or more lines of systemic therapy. This intervention aims to assess its preliminary efficacy based on the intrahepatic disease control rate at 12 weeks. Currently in Phase 2 and actively recruiting, this trial offers patients the opportunity to participate in a study focused on a specific treatment approach for their condition. Eligibility criteria include specific biomarkers such as MSI, MSI-H, MMR, DMMR, TMB, RAS, and microsatellite instability.

NCT: NCT06555133 · Enrollment: 18 · Sites: Providence

MEDIUM

PHASE2

Stage IV

MSI/MMR

TARGETED THERAPY

• RECRUITING

5

SCRT Combined With Chemotherapy and Iparomlimab and Tuvonralimab in MSS or pMMR Patients With Locally Advanced Rectal Cancer

This clinical trial is testing a novel neoadjuvant regimen for patients with locally advanced rectal cancer (LARC) classified as T3-4/N+ without distant metastasis. The intervention involves short-course radiotherapy followed by four cycles of CAPEOX combined with the drugs Iparomlimab and Tuvonralimab, aiming to improve organ preservation and treatment efficacy. Currently in Phase 2 and actively recruiting, this trial offers patients the opportunity to participate in research that may enhance treatment outcomes. Eligibility criteria include patients with microsatellite stable (MSS) or mismatch repair proficient (pMMR) tumors, as well as those with specific biomarker profiles.

NCT: NCT06864013 · Enrollment: 116 · Sites: Hangzhou, China

MEDIUM

PHASE1, PHASE2

Stage IV

BRAF

TARGETED THERAPY

• NOT_YET_RECRUITING

5

Liposomal Irinotecan Plus Bevacizumab in Irinotecan-refractory Metastatic Colorectal Cancer

This clinical trial is testing the combination of liposomal irinotecan and bevacizumab in patients with irinotecan-refractory metastatic colorectal cancer. The intervention involves administering liposomal irinotecan, which is being evaluated for its efficacy and safety alongside bevacizumab, a drug that may enhance treatment outcomes. This study is in Phase 1 and Phase 2 and is currently not yet recruiting, meaning that patients will need to wait until recruitment begins to participate. Eligibility criteria include specific biomarker requirements such as BRAF, MSI, MMR, and RAS status.

NCT: NCT06434090 · Enrollment: 74 · Sites: Guangzhou, China

MEDIUM

N/A

All Stages

MSI/MMR

TARGETED THERAPY

• RECRUITING

5

Immune Checkpoint Inhibitor Response in Solid Tumors Using a Live Tumor Diagnostic Platform

This clinical trial is investigating the accuracy of the Elephas Score in predicting clinical response to various immunotherapies and chemoimmunotherapy in patients with solid tumors. The intervention involves the collection of biospecimens and tissue samples, which are essential for evaluating the effectiveness of the Elephas diagnostic platform. The trial is currently in the recruiting phase, meaning eligible patients can participate and contribute to the research. Eligibility criteria include specific biomarkers such as MSI, MSI-H, MMR, DMMR, TMB, and microsatellite status.

NCT: NCT06349642 · Enrollment: 324 · Sites: Jacksonville, Phoenix, Rochester

MEDIUM

NA

Stage IV

IMMUNOTHERAPY

• RECRUITING

5

A Randomized, Multicenter Phase II Basket Study of Hypofractionated Radiotherapy/Stereotactic Body Radiotherapy Followed by Immunotherapy-Based Systemic Therapy +/- L. Rhamnosus M9 for the First-Line Treatment of Advanced Digestive System Malignancies.

This clinical trial is testing the effectiveness of hypofractionated radiotherapy or stereotactic body radiotherapy followed by immunotherapy-based systemic therapy, with or without L. Rhamnosus M9, for patients with advanced digestive system malignancies. The intervention includes a combination of radiation therapy and various drugs, such as anti-PD-1 monoclonal antibodies and chemotherapy agents, which may enhance the immune response against tumors. Currently in the recruiting phase, this Phase II study aims to enroll 120 participants, indicating that eligible patients may have the opportunity to receive this integrated treatment approach. Specific eligibility criteria or biomarker requirements have not been detailed in the provided information.

NCT: NCT06349044 · Enrollment: 120 · Sites: Hangzhou, China

MEDIUM

PHASE1

Stage IV

MSI/MMR

TARGETED THERAPY

• RECRUITING

5

Dose Escalation and Expansion Clinical Trial of Irinotecan Liposome Combined With Oxaliplatin and 5-FU/LV Plus Bevacizumab as First-line Treatment of Metastatic Colorectal Cancer

This clinical trial is testing the combination of irinotecan liposome, oxaliplatin, 5-FU/LV, and bevacizumab as a first-line treatment for patients with advanced metastatic colorectal cancer. The intervention aims to determine the maximum tolerated dose and assess the safety and efficacy of this drug combination, which is significant for optimizing treatment regimens. Currently in Phase 1 and actively recruiting participants, this trial offers patients the opportunity to receive a potentially effective treatment while contributing to the understanding of dose-limiting toxicities. Eligibility criteria include specific biomarkers such as MSI, MSI-H, MMR, and DMMR.

NCT: NCT06225622 · Enrollment: 78 · Sites: Shanghai, China

MEDIUM

PHASE2

Stage IV

MSI/MMR

TARGETED THERAPY

• RECRUITING

5

Efficacy and Safety of Chemotherapy With XELOX (Oxaliplatin + Capecitabine) and Bevacizumab in Combination With Adebrelimab in First-line Treatment of Microsatellite Stable (MSS) Initially Unresectable Metastatic Colorectal Cancer

This clinical trial is testing the efficacy and safety of a combination treatment for patients with microsatellite stable (MSS) initially unresectable metastatic colorectal cancer. The intervention involves the use of Adebrelimab alongside chemotherapy agents Oxaliplatin and Capecitabine, as well as Bevacizumab, which may enhance treatment outcomes. Currently in Phase 2 and actively recruiting participants, this trial aims to assess progression-free survival as its primary outcome, providing insights into the effectiveness of this treatment regimen. Eligibility criteria include specific biomarker assessments such as MSI, MMR, and RAS status.

NCT: NCT06282445 · Enrollment: 36 · Sites: Jinhua, China

MEDIUM

PHASE2

All Stages

BRAF

LIQUID BIOPSY

• ACTIVE_NOT_RECRUITING

5

Clinical Trial of All-trans-retinoic Acid, Bevacizumab and Atezolizumab in Colorectal Cancer

This clinical trial is testing the combination of all-trans-retinoic acid (ATRA), bevacizumab, and atezolizumab for patients with advanced colorectal cancer. The intervention involves administering ATRA orally, while bevacizumab and atezolizumab are given intravenously, which may provide insights into their combined effects on treatment outcomes. Currently in Phase 2 and marked as active but not recruiting, this trial indicates that patients are not being enrolled at this

time, but results may still be forthcoming. Eligibility criteria include specific biomarkers such as BRAF, MSI, MSI-H, MMR, HER2, ctDNA, RAS, and microsatellite status.

NCT: NCT05999812 · Enrollment: 22 · Sites: Dallas

MEDIUM

PHASE4

Stage IV

MSI/MMR

TARGETED THERAPY

• RECRUITING

5

Survival Benefit of Compound Kushen Injection in the Treatment of Advanced Colorectal Cancer

This clinical trial is testing the effectiveness and safety of compound kushen injection in patients with advanced colorectal cancer. The intervention involves administering compound kushen injection alongside a first-line treatment scheme, which is significant for its potential impact on patient outcomes. Currently in Phase 4 and actively recruiting, this trial aims to provide insights into progression-free survival (PFS) for participants. Eligibility criteria include specific biomarkers such as MSI, MSI-H, MMR, DMMR, and RAS.

NCT: NCT05894694 · Enrollment: 320 · Sites: Beijing, China

MEDIUM

N/A

All Stages

MSI/MMR

TARGETED THERAPY

• RECRUITING

5

Observational Basket Trial to Collect Tissue to Develop and Train a Live Tumor Diagnostic Platform

This observational basket trial is designed for patients with colorectal cancer to collect tissue samples for the development and training of the Elephas live tumor diagnostic platform. The intervention involves various biopsy procedures, including core needle, forceps, and punch biopsies, which are essential for obtaining the necessary tissue for analysis. The trial is currently in the recruiting phase, indicating that eligible patients can participate and contribute to the advancement of diagnostic methods. Eligibility criteria include specific biomarker requirements such as MSI, MMR, and TMB, which are important for the study's focus on tumor characteristics.

NCT: NCT05520099 · Enrollment: 416 · Sites: Buffalo, Chapel Hill, Hackensack, Hagerstown, Little Rock

MEDIUM

PHASE2

All Stages

KRAS

TARGETED THERAPY

• NOT_YET_RECRUITING

5

A Combination Therapy Including Anti-PD-1 Immunotherapy in MSS Rectal Cancer With Resectable Distal Metastasis

This clinical trial is testing a combination therapy that includes the anti-PD-1 immunotherapy tislelizumab for patients with microsatellite-stable (MSS) rectal cancer who have resectable distal metastasis. The intervention aims to evaluate the effectiveness of preoperative short-course radiotherapy followed by neoadjuvant chemotherapy and immunotherapy in improving tumor response and reducing postoperative metastasis and recurrence. This is a Phase 2 trial that is currently not yet recruiting, meaning patients will need to wait until enrollment begins to participate. Eligibility criteria include specific biomarker assessments such as KRAS, NRAS, BRAF, MMR, DMMR, HER2, RAS, and microsatellite status.

NCT: NCT05359393 · Enrollment: 52 · Sites: Shanghai, China

MEDIUM

N/A

Stage IV

MSI/MMR

TARGETED THERAPY

• ACTIVE_NOT_RECRUITING

5

Retrospective Study on the Use of Immunotherapy in Patients With MSI-H Metastatic Colorectal Cancer

This clinical trial is a retrospective study examining the use of immunotherapy in patients with microsatellite instability-high (MSI-H) metastatic colorectal cancer (mCRC). The intervention involves the use of immune checkpoint inhibitors, specifically pembrolizumab and nivolumab, which are significant for their role in targeting the immune response against cancer cells. The trial is currently active but not recruiting, indicating that it is in the data collection phase, which may provide insights into treatment effectiveness for patients who have exhausted other options. Eligibility criteria include having MSI-H status, which is essential for determining the appropriateness of immunotherapy in this patient population.

NCT: NCT04612309 · Enrollment: 100 · Sites: Carpi, Italy

MEDIUM

PHASE2

Stage III

MSI/MMR

TARGETED THERAPY

• RECRUITING

5

Neo-adjuvant Pembrolizumab and Radiotherapy in Localised MSS Rectal Cancer

This clinical trial is testing the combination of pembrolizumab and short course radiotherapy in patients with localized microsatellite stable (MSS) rectal cancer. The intervention involves administering pembrolizumab, an immunotherapy drug, alongside radiotherapy to evaluate its effects on tumor regression. This is a Phase 2 trial that is currently recruiting

participants, which means eligible patients may have the opportunity to receive this treatment while contributing to the research. Eligibility criteria include having localized MSS rectal cancer and specific biomarker assessments related to microsatellite stability.

NCT: NCT04109755 · Enrollment: 25 · Sites: Geneva, Switzerland

MEDIUM PHASE2 Stage IV RAS IMMUNOTHERAPY • RECRUITING

5

Immunotherapy Using Tumor Infiltrating Lymphocytes for Patients With Metastatic Cancer

This clinical trial is testing an experimental immunotherapy using Tumor Infiltrating Lymphocytes (TIL) for patients with metastatic cancer, specifically targeting those with digestive tract, urothelial, breast, or ovarian/endometrial cancers. The intervention involves the administration of pembrolizumab, fludarabine, cyclophosphamide, aldesleukin, and young TIL, which are designed to enhance the body's immune response against tumors. Currently in Phase 2 and actively recruiting, this trial aims to evaluate the response rate of the treatment, which is crucial for determining its effectiveness and safety for patients. Eligible participants include adults aged 18-72 with specific cancer types and RAS biomarker requirements.

NCT: NCT01174121 · Enrollment: 332 · Sites: Bethesda

MEDIUM NA Stage IV MSI/MMR TARGETED THERAPY • RECRUITING

4

Clinical Efficacy of Transarterial Infusion Chemotherapy for Unresectable Colorectal Cancer

This clinical trial is testing the efficacy of transarterial infusion chemotherapy (TAIC) combined with either cetuximab or bevacizumab for patients with unresectable colorectal cancer (CRC). The intervention involves a procedure that delivers chemotherapy directly to the tumor site, which may enhance treatment effectiveness. The trial is currently in the recruiting phase, indicating that eligible patients can still enroll to participate in the study. Eligibility criteria include specific biomarkers such as MSI, MSI-H, MMR, DMMR, and RAS.

NCT: NCT07333053 · Enrollment: 30 · Sites: Xicheng, China

MEDIUM PHASE2 Stage IV BRAF TARGETED THERAPY • RECRUITING

4

A Phase II Study of Sintilimab Combined With Ipilimumab N01, Cetuximab and Dabrafenib in Patients With Microsatellite-Stable, BRAF V600E-Mutated Metastatic Colorectal Cancer

This clinical trial is testing the combination of sintilimab, ipilimumab N01, cetuximab, and dabrafenib in patients with microsatellite-stable, BRAF V600E-mutated metastatic colorectal cancer. The intervention involves a multi-drug regimen aimed at enhancing treatment efficacy for a patient population that typically has a poor prognosis with standard therapies. Currently in Phase II and actively recruiting participants, this trial seeks to assess the safety and efficacy of the treatment approach, which may offer an alternative for patients who have limited options. Eligibility criteria include the presence of BRAF mutations and microsatellite stability, which are essential for participation in this study.

NCT: NCT07506109 · Enrollment: 49 · Sites: Beijing, China, Chengdu, China, Tianjin, China, Wuhan, China

MEDIUM PHASE1 Stage IV RAS TARGETED THERAPY • RECRUITING

4

A Study of IDE034 in Adult Participants With Locally Advanced/Metastatic Solid Tumors Types

This clinical trial is testing the drug IDE034 in adult participants with locally advanced or metastatic solid tumors. The intervention, IDE034, is being evaluated for its safety, pharmacokinetics, immunogenicity, and preliminary efficacy, particularly in tumors that express the biomarkers B7-H3 and PTK7. Currently in Phase 1 and actively recruiting, this trial aims to assess the drug's safety and tolerability during the dose escalation phase, which is crucial for determining appropriate dosing for future studies. Eligibility criteria include the expression of specific biomarkers, such as HER2 and RAS, in the tumor.

NCT: NCT07503808 · Enrollment: 150 · Sites: Austin, Fairfax, Houston, Irving, San Antonio

MEDIUM PHASE2 Stage IV BRAF TARGETED THERAPY • NOT_YET_RECRUITING

4

A Single-arm, Single-center, Phase II Clinical Study of Aipalolitovorelizumab (QL1706) Combined With Bevacizumab and Standard Chemotherapy as First-line Treatment for MSS/pMMR Metastatic Colorectal Cancer With BRAF V600E Mutation

This clinical trial is testing the combination of Aipalolitovorelizumab (QL1706) with bevacizumab and standard chemotherapy as a first-line treatment for patients with MSS/pMMR metastatic colorectal cancer who have a BRAF

V600E mutation. The intervention involves administering Aipalolitovorelizumab alongside various chemotherapy regimens, which is significant for potentially improving treatment outcomes in this specific patient population. This is a Phase II study that is not yet recruiting, meaning patients will need to wait until enrollment begins to participate. Eligibility criteria include having the BRAF V600E mutation and meeting specific biomarker requirements related to MMR status.

NCT: NCT07229846 · Enrollment: 30 · Sites: Harbin, China

SIGNAL

NA

Early Detection

RESEARCH

• RECRUITING

⚡ 4

Randomized Trial Comparing Fecal Testing (FIT) to Colonoscopy for Post-polypectomy Surveillance

This clinical trial is testing the effectiveness of fecal occult blood testing (FIT) compared to colonoscopy for post-polypectomy surveillance in patients who have had colorectal polyps removed. The intervention involves using FIT as a less burdensome alternative to colonoscopy, which is important given the limited availability of colonoscopy and the fact that many surveillance procedures yield no findings. The trial is currently in the recruiting phase with an enrollment goal of 4,000 participants, which means eligible patients can still join the study. Eligibility criteria may include a history of polyp removal, but specific requirements are not detailed in the provided information.

NCT: NCT06983639 · Enrollment: 4000 · Sites: Arendal, Norway

MEDIUM

PHASE1, PHASE2

Stage IV

BRAF

TARGETED THERAPY

• NOT_YET_RECRUITING

⚡ 4

A Phase Ib/II Trial of HS-20110 Combination Therapies in Advanced Colorectal Cancer Patients.

This clinical trial is evaluating the safety and efficacy of HS-20110 combination therapies in patients with advanced colorectal cancer. The interventions include HS-20110 combined with Bevacizumab and various chemotherapy regimens, which is important for assessing potential treatment options for this patient population. The trial is currently in Phase Ib/II and is not yet recruiting, meaning patients will need to wait for enrollment to begin before participating. Eligibility criteria include specific biomarkers such as BRAF and MSI, which may be relevant for patient selection.

NCT: NCT07283367 · Enrollment: 502 ·

MEDIUM

PHASE1, PHASE2

Stage IV

BRAF

IMMUNOTHERAPY

• RECRUITING

⚡ 4

A FIH, Phase I/IIa, Trial Assessing Feasibility of Administrations of TIL-based Immunotherapy in Patients With Metastatic CRC and PC

This clinical trial is testing the feasibility of TIL-based immunotherapy in patients with metastatic colorectal cancer (CRC) and pancreatic cancer (PC). The intervention involves the administration of a novel autologous TIL product, CC-38, along with pembrolizumab, cyclophosphamide, interleukin-2, and uromitexan, which is important for evaluating safety and tolerability. Currently in Phase I/IIa and actively recruiting, this trial aims to assess the repeated administration of CC-38 in a small cohort of 12 participants. Eligibility criteria include the presence of the BRAF biomarker.

NCT: NCT07255664 · Enrollment: 12 · Sites: Frankfurt, Germany

MEDIUM

PHASE2

All Stages

MSI/MMR

LIQUID BIOPSY

• RECRUITING

⚡ 4

A Study of Botensilimab and Balstilimab for Colorectal Cancer With ctDNA+ After Surgery and Chemotherapy

This clinical trial is testing the combination of botensilimab and balstilimab for patients with microsatellite stable (MSS) colorectal cancer or colorectal liver metastases (CRLM) who have measurable residual disease (MRD) after surgery and chemotherapy. The intervention involves administering botensilimab and balstilimab, followed by balstilimab alone, to evaluate its effectiveness in achieving ctDNA clearance. This is a Phase 2 trial currently recruiting 284 participants, which means eligible patients may have the opportunity to receive this investigational treatment. Eligibility criteria include specific biomarkers such as MSI, MSI-H, MMR, ctDNA, RAS, and NGS.

NCT: NCT07227636 · Enrollment: 284 · Sites: Basking Ridge, Commack, Harrison, Middletown, Montvale

MEDIUM

PHASE2

Stage IV

BRAF

TARGETED THERAPY

• RECRUITING

⚡ 4

Amplitude-Modulated Radiofrequency Electromagnetic Fields (AM RF EMF) in Combination With Fruquintinib in Refractory Metastatic Colorectal Cancer

This clinical trial is testing the combination of amplitude-modulated radiofrequency electromagnetic fields (AM RF EMF) with Fruquintinib in patients with refractory metastatic colorectal cancer. The intervention involves using Fruquintinib, a drug, alongside the TheraBionic P1 device to assess its effectiveness in improving overall survival and its safety profile. Currently in Phase 2 and actively recruiting participants, this trial aims to provide insights into treatment options for patients whose cancer has not responded to standard therapies. Eligibility criteria include specific biomarkers such as BRAF, MSI, MMR, HER2, RAS, and microsatellite instability.

NCT: NCT07130903 · Enrollment: 102 · Sites: Bay City, Clarkston, Detroit, Flint, Lansing

MEDIUM

PHASE1

Stage IV

KRAS

IMMUNOTHERAPY

• RECRUITING

⚡ 4

AMG 410 Alone and in Combination With Other Agents in Participants With KRAS Altered Advanced or Metastatic Solid Tumors

This clinical trial is testing the safety and efficacy of AMG 410, both alone and in combination with pembrolizumab and panitumumab, in participants with advanced or metastatic solid tumors that have KRAS alterations. The intervention involves a dose-escalation study to identify the Maximum Tolerated Dose (MTD) and the Recommended Phase 2 Dose (RP2D) of AMG 410, which is important for determining the optimal dosing strategy for these patients. Currently in Phase 1 and actively recruiting, this trial aims to assess dose limiting toxicities (DLTs) and other pharmacological parameters, which will inform future treatment options. Eligibility criteria include having KRAS alterations, which is a specific biomarker requirement for participation.

NCT: NCT07094113 · Enrollment: 434 · Sites: Atlanta, Boston, Duarte, Durham, Fairfax

MEDIUM

PHASE2

Stage III

MSI/MMR

TARGETED THERAPY

• RECRUITING

⚡ 4

A Phase II Trial of Neoadjuvant PD-1 Vaccine PD1-Vaxx in Operable MSI High Colorectal Cancer

This clinical trial is testing the neoadjuvant PD-1 vaccine IMU-201 (PD1-Vaxx) in patients with operable microsatellite instability-high (MSI-H) colorectal cancer. The intervention involves administering three doses of the PD1-Vaxx prior to surgical resection, with the goal of achieving a Major Pathological Response (MPR) defined as less than 10% viable tumor cells. This is a Phase II trial that is currently recruiting participants, which means eligible patients may have the opportunity to receive this treatment while contributing to the study's findings. Eligibility criteria include having operable MSI-H colorectal cancer, with specific biomarker requirements such as MSI, MSI-H, MMR, and NGS.

NCT: NCT06692959 · Enrollment: 44 · Sites: Adelaide, Australia, Guildford, United Kingdom, Manchester, United Kingdom, Perth, Australia

SIGNAL

NA

Stage IV

RESEARCH

• RECRUITING

⚡ 4

Randomized Controlled Trial of the Effects of Combined Resistance and Aerobic Exercise on Health-related Quality of Life in Patients Undergoing First-line Chemotherapy for Metastatic Colorectal Cancer (REACH)

The REACH trial is testing the effects of combined resistance and aerobic exercise on health-related quality of life in patients undergoing first-line chemotherapy for metastatic colorectal cancer. The intervention involves 18 weeks of structured exercise training, which aims to assess its impact on the quality of life for these patients. This trial is currently in the recruiting phase, meaning eligible patients can still enroll to participate. Eligibility criteria include being diagnosed with unresectable metastatic colorectal cancer and undergoing first-line chemotherapy.

NCT: NCT06938971 · Enrollment: 150 · Sites: Copenhagen, Denmark, Herlev, Denmark, Hillerød, Denmark, Roskilde, Denmark

MEDIUM

PHASE2

Stage IV

MSI/MMR

TARGETED THERAPY

• NOT_YET_RECRUITING

⚡ 4

Short-course Radiotherapy Followed by CAPOX and Ivonescimab for Locally Advanced Rectal Cancer

This clinical trial is testing the efficacy and safety of preoperative short-course radiotherapy combined with ivonescimab and chemotherapy in patients with locally advanced rectal cancer. The intervention involves the drug ivonescimab, which is being evaluated for its potential to improve treatment outcomes when used alongside standard therapies. This is a Phase 2 trial that is not yet recruiting, meaning patients will need to wait until enrollment begins to participate. Eligibility criteria include specific biomarkers such as MSI, MSI-H, MMR, and DMMR.

NCT: NCT06760520 · Enrollment: 40 ·

SIGNAL **PHASE1** **Stage IV** **KRAS** **LIQUID BIOPSY** • **RECRUITING**

⚡ 4

A First-in-human Study of BGB-53038, a Pan-KRAS Inhibitor, Alone or in Combinations in Participants With Advanced or Metastatic Solid Tumors With KRAS Mutations or Amplification

This clinical trial is testing BGB-53038, a pan-KRAS inhibitor, in participants with advanced or metastatic solid tumors that have KRAS mutations or amplification. The intervention includes BGB-53038 alone and in combination with tislelizumab or cetuximab, which is significant for understanding its safety and potential effectiveness in treating these specific cancers. Currently in Phase 1 and actively recruiting, this trial aims to assess the safety, tolerability, and preliminary antitumor activity of the drug, which is crucial for determining the next steps in treatment options for eligible patients. Participants must have tumors with KRAS mutations or amplification, and the study includes a liquid biopsy component for biomarker assessment.

NCT: NCT06585488 · Enrollment: 514 · Sites: Baltimore, Houston, Kansas City, Los Angeles

MEDIUM **NA** **Stage IV** **MSI/MMR** **TARGETED THERAPY** • **RECRUITING**

⚡ 4

Study on the Safety and Efficacy of MR-Linac Technique in Patients With Unresectable Locally Advanced Colon Cancer

This clinical trial is testing the safety and efficacy of the MR-Linac technique in patients with unresectable locally advanced colon cancer. The intervention involves the use of MR-Linac for short course radiotherapy, followed by chemotherapy and surgical resection, which is significant for potentially improving treatment outcomes in this patient population. The study is currently in the recruiting phase and aims to enroll 20 participants, indicating that eligible patients may still join the trial. Eligibility criteria include specific biomarkers such as MSI, MSI-H, MMR, and dMMR.

NCT: NCT06244537 · Enrollment: 20 · Sites: Chengdu, China

MEDIUM **PHASE1, PHASE2** **Stage IV** **IMMUNOTHERAPY** • **RECRUITING**

⚡ 4

GM103 Intratumoral Injection in Patients With Locally Advanced, Unresectable, Refractory and/or Metastatic Solid Tumors

This clinical trial is testing the safety and efficacy of GM103, administered alone and in combination with pembrolizumab, in patients with locally advanced, unresectable, refractory, and/or metastatic solid tumors. The intervention involves the intratumoral injection of GM103, which is being evaluated for its potential to improve treatment outcomes in various cancer types, including colorectal cancer. Currently in Phase 1 and Phase 2, the trial is actively recruiting participants, which means eligible patients may have the opportunity to receive this investigational treatment. Eligibility criteria include having specific types of solid tumors that are advanced and difficult to treat.

NCT: NCT06265025 · Enrollment: 125 · Sites: Goyang-si, South Korea, Seoul, South Korea

MEDIUM **PHASE1, PHASE2** **Stage IV** **KRAS** **TARGETED THERAPY** • **RECRUITING**

⚡ 4

Study of Elironrasib and Daraxonrasib as Monotherapies and Combination Therapy in Participants With Advanced KRAS G12C Mutant Solid Tumors

This clinical trial is evaluating the safety and tolerability of Elironrasib and Daraxonrasib, both as monotherapies and in combination, for participants with advanced KRAS G12C mutant solid tumors. The intervention involves two drugs, Elironrasib and Daraxonrasib, which are being tested to understand their pharmacokinetics and potential effects on this specific mutation. Currently in Phase 1 and Phase 2, the trial is actively recruiting participants, which means eligible patients may have the opportunity to receive these treatments. Eligibility criteria include having tumors with KRAS G12C mutations, among other potential biomarker requirements.

NCT: NCT06128551 · Enrollment: 534 · Sites: Aurora, Boston, Dallas, Detroit, Duarte

MEDIUM **PHASE1, PHASE2** **Stage IV** **MSI/MMR** **TARGETED THERAPY** • **RECRUITING**

⚡ 4

Cadonilimab for PD-1/PD-L1 Blockade-refractory, MSI-H/dMMR, Advanced Colorectal Cancer

This clinical trial is testing the efficacy of cadonilimab in patients with advanced colorectal cancer who are refractory to PD-1/PD-L1 blockade and have microsatellite instability-high (MSI-H) or mismatch repair-deficient (dMMR) tumors. The intervention, cadonilimab, is a drug that targets the immune checkpoint pathway, which is significant for patients who have not responded to other therapies. This trial is currently in Phase 1 and Phase 2 and is actively recruiting participants, indicating that eligible patients may have the opportunity to access this treatment option. Eligibility criteria include specific biomarkers such as MSI, MSI-H, MMR, dMMR, RAS, and microsatellite status.

SIGNAL

N/A

Stage III

ctDNA

LIQUID BIOPSY

• RECRUITING

⚡ 4

ctDNA Methylation Application in Postoperative Relapse and Adjuvant Chemotherapy Efficacy Evaluation

This clinical trial is testing a multi-locus blood-based assay targeting circulating tumor DNA (ctDNA) methylation in patients with resected stage I and stage II colorectal cancer who are at low risk for recurrence. The intervention aims to monitor postoperative relapse and evaluate the efficacy of adjuvant chemotherapy, which is important for optimizing treatment strategies in this patient population. The trial is currently in the recruiting phase, meaning eligible patients can still enroll to participate. Eligibility criteria include having undergone radical resection for stage I or stage II colorectal cancer without high-risk features.

NCT: NCT05536089 · Enrollment: 2000 · Sites: Hangzhou, China

MEDIUM

PHASE1

All Stages

MSI/MMR

IMMUNOTHERAPY

• RECRUITING

⚡ 4

Study of Zanzalintinib in Combination With Immuno-Oncology Agents in Participants With Solid Tumors

This clinical trial is evaluating the safety and efficacy of zanzalintinib, both as a monotherapy and in combination with immuno-oncology agents, in participants with advanced solid tumors. The intervention involves zanzalintinib, nivolumab, and ipilimumab, which are being tested to assess their safety, tolerability, and preliminary antitumor activity. This is a Phase 1b trial currently recruiting participants, which means that it is in the early stages of testing to determine the best dosing and safety profile before larger studies can be conducted. Eligibility criteria include the presence of specific biomarkers, such as microsatellite instability.

NCT: NCT05176483 · Enrollment: 1314 · Sites: Austin, Baltimore, Boston, Celebration, Charlottesville

MEDIUM

PHASE2

Stage III

MSI/MMR

TARGETED THERAPY

• ACTIVE_NOT_RECRUITING

⚡ 4

Atezolizumab in Patients With MSI-h/MMR-D Stage II High Risk and Stage III Colorectal Cancer Ineligible for Oxaliplatin

This clinical trial is testing the efficacy of atezolizumab in patients with stage II high-risk and stage III colorectal cancer who are ineligible for oxaliplatin. The intervention involves administering atezolizumab, a drug that targets the immune system, alongside IMM-101, to potentially enhance disease-free survival in this specific patient population. Currently in Phase 2 and marked as active but not recruiting, this trial indicates that patients may not be able to enroll at this time but can still benefit from the findings once available. Eligibility criteria include biomarkers such as MSI, MSI-H, MMR, DMMR, RAS, and NGS.

NCT: NCT05118724 · Enrollment: 80 · Sites: Essen, Germany

MEDIUM

PHASE1, PHASE2

All Stages

IMMUNOTHERAPY

• RECRUITING

⚡ 4

A Beta-only IL-2 ImmunoTherapY Study

This clinical trial is testing the safety and efficacy of MDNA11, both as a monotherapy and in combination with Pembrolizumab (Keytruda®), for patients with advanced solid tumors, including colorectal cancer. The intervention involves a dose-escalation and expansion study to determine the recommended doses for both monotherapy and combination therapy, which is important for optimizing treatment strategies. Currently in Phase 1/2 and actively recruiting, this trial offers patients the opportunity to participate in early-stage research that may lead to improved treatment options. Eligibility criteria may include specific biomarkers, although details are not provided.

NCT: NCT05086692 · Enrollment: 115 · Sites: Atlanta, Boca Raton, Detroit, Houston, San Diego

MEDIUM

PHASE1

All Stages

HER2

TARGETED THERAPY

• ACTIVE_NOT_RECRUITING

⚡ 4

CAR-macrophages for the Treatment of HER2 Overexpressing Solid Tumors

This clinical trial is testing the safety and tolerability of CAR-macrophages in patients with HER2 overexpressing solid tumors. The intervention involves the use of CT-0508 and Pembrolizumab, which are biological agents aimed at targeting and treating these specific tumors. Currently in Phase 1 and marked as active but not recruiting, this trial is focused on evaluating adverse events in the enrolled participants. Eligibility criteria include the requirement for tumors to overexpress the HER2 biomarker.

MEDIUM

PHASE1, PHASE2

Stage IV

IMMUNOTHERAPY

• ACTIVE_NOT_RECRUITING

⚡ 4

Safety, PK and Efficacy of ONC-392 in Monotherapy and in Combination of Anti-PD-1 in Advanced Solid Tumors and NSCLC

This clinical trial is testing the safety, pharmacokinetics, and efficacy of ONC-392, a humanized anti-CTLA4 IgG1 monoclonal antibody, in patients with advanced or metastatic solid tumors and non-small cell lung cancer (NSCLC). The intervention involves administering ONC-392 both as a monotherapy and in combination with pembrolizumab and docetaxel, which is significant for understanding its potential effectiveness in these cancer types. Currently, the trial is in Phase 1 and Phase 2 and is active but not recruiting, indicating that patients cannot enroll at this time. Eligibility criteria may include specific biomarkers, although details on these requirements are not provided.

NCT: NCT04140526 · Enrollment: 733 · Sites: Ann Arbor, Atlanta, Atlantis, Aurora, Baltimore

MEDIUM

PHASE2

All Stages

HER2

TARGETED THERAPY

• ACTIVE_NOT_RECRUITING

⚡ 4

Tucatinib Plus Trastuzumab and Oxaliplatin-based Chemotherapy or Pembrolizumab-containing Combinations for HER2+ Gastrointestinal Cancers

This clinical trial is testing the safety and efficacy of tucatinib in combination with trastuzumab and various chemotherapy regimens for patients with HER2-positive gastrointestinal cancers. The intervention involves administering tucatinib alongside trastuzumab, oxaliplatin, leucovorin, and fluorouracil, with the aim of assessing potential side effects and therapeutic effectiveness. Currently in Phase 2 and marked as active but not recruiting, this status indicates that the trial is ongoing but not accepting new participants at this time. Eligible participants must have HER2-positive cancer affecting the gut, stomach, intestines, or gallbladder, and may also be evaluated for RAS biomarkers.

NCT: NCT04430738 · Enrollment: 40 · Sites: Albuquerque, Aurora, City of Saint Peters, Cleveland, Creve Coeur

SIGNAL

NA

Stage III

ctDNA

LIQUID BIOPSY

• ACTIVE_NOT_RECRUITING

⚡ 4

ctDNA-guided Surveillance for Stage III CRC, a Randomized Intervention Trial

The IMPROVE-IT2 trial is testing ctDNA-guided surveillance for patients with Stage III colorectal cancer (CRC) to evaluate its effectiveness compared to standard CT-scan surveillance. The intervention involves ctDNA analysis as a diagnostic test, which aims to enhance the detection of recurrent disease and potentially increase the number of patients eligible for curative treatment. This trial is currently active but not recruiting participants, indicating that it is ongoing and data is being collected from enrolled patients. Eligibility criteria include the presence of ctDNA as a biomarker for assessment.

NCT: NCT04084249 · Enrollment: 359 · Sites: Aalborg, Denmark, Aarhus, Denmark, Copenhagen, Denmark, Herlev, Denmark, Herning, Denmark

MEDIUM

PHASE1

Stage IV

RAS

TARGETED THERAPY

• ACTIVE_NOT_RECRUITING

⚡ 4

Naptumomab Estafenatox in Combination With Durvalumab in Subjects With Selected Advanced or Metastatic Solid Tumor, Including a Cohort Expansion in Esophageal Cancer.

This clinical trial is testing the combination of naptumomab estafenatox and durvalumab in patients with selected advanced or metastatic solid tumors, including a cohort expansion specifically for esophageal cancer. The intervention involves a combination of these two agents to evaluate their safety and tolerability, which is important for determining effective treatment options for this patient population. Currently, the trial is in Phase 1b and is active but not recruiting, meaning that while it is ongoing, no new participants are being enrolled at this time. Eligibility criteria include specific biomarkers such as HER2, RAS, and NGS, which may be relevant for patient selection.

NCT: NCT03983954 · Enrollment: 120 · Sites: Ahmedabad, India, Haifa, Israel, Hyderabad, India, Jhajjar, India, New Delhi, India

MEDIUM

PHASE2

Stage IV

PHASE II TRIAL

• ACTIVE_NOT_RECRUITING

⚡ 4

Microwave Ablation Versus Stereotactic Body Radiotherapy for Colorectal Liver Metastases in Oligometastatic Disease: a Prospective, Randomised, Phase 2 Trial

The LAVA-CRLM trial is testing the effectiveness of microwave ablation (MWA) compared to stereotactic body radiotherapy (SBRT) in patients with colorectal liver metastases and oligometastatic disease. The intervention involves using MWA, a device-based treatment, and SBRT, a form of radiation therapy, to assess their impact on local control and safety in this patient population. This is a Phase 2 trial that is currently active but not recruiting, indicating that while the

study is ongoing, no new participants are being enrolled at this time. Eligibility criteria for participation may include specific characteristics related to the patient's metastatic disease status.

NCT: NCT03654131 · Enrollment: 100 · Sites: Copenhagen, Denmark

MEDIUM PHASE2 All Stages RAS IMMUNOTHERAPY • ACTIVE_NOT_RECRUITING

⚡ 4

Nivolumab and Ipilimumab in Treating Patients With Rare Tumors

This phase II clinical trial is investigating the efficacy of nivolumab and ipilimumab in treating patients with rare tumors. The intervention involves the use of these monoclonal antibodies, which may enhance the immune system's ability to combat cancer and inhibit tumor growth and spread. Currently, the trial is active but not recruiting new participants, indicating that it is ongoing with enrolled patients being monitored for outcomes. Eligibility criteria include specific tumor types, such as epithelial tumors of the nasal cavity and salivary glands, with a focus on RAS biomarkers.

NCT: NCT02834013 · Enrollment: 818 · Sites: Aberdeen, Adrian, Aitkin, Albuquerque, Allentown

Appendix A — Patient Guide to Active Trials

For patients, caregivers, and family members. Always discuss with your oncologist before considering a clinical trial.

What is a clinical trial?

Clinical trials test new treatments, combinations, or approaches to care. Participating can give you access to treatments not yet widely available. Trials are listed by the biomarker your tumor must have — because many newer treatments only work for specific mutations. Ask your oncologist which biomarkers your tumor has been tested for, then look for matching trials below.

MSI/MMR

Botensilimab + Balstilimab vs Best Supportive Care as Therapy in Chemo-refractory, Unresectable, Colorectal Adenocarcinoma

Phase: PHASE3 · Status: RECRUITING · [NCT07152821](#)

This clinical trial is testing the combination of botensilimab and balstilimab in patients with chemo-refractory, unresectable colorectal adenocarcinoma. The intervention involves two immunotherapy drugs aimed at potentially improving overall survival and quality of life for participants. Currently in Phase 3 and actively recruiting, this trial may provide insights into the effectiveness and safety of these treatments for eligible patients. Participants may need to meet specific biomarker criteria, including MSI, MSI-H, MMR, DMMR, and microsatellite status.

BRAF

Symbiotic-GI-03: A Study to Learn About the Study Medicine Called PF-08634404 in Combination With Chemotherapy in Adult Participants With Metastatic Colorectal Cancer

Phase: PHASE3 · Status: RECRUITING · [NCT07222800](#)

The Symbiotic-GI-03 trial is investigating the efficacy of the study medicine PF-08634404 in combination with chemotherapy for adult participants with metastatic colorectal cancer. This trial aims to determine how well PF-08634404 works alongside standard chemotherapy and Bevacizumab, a biological therapy, in patients whose cancer has spread or returned after previous treatments. Currently in Phase 3 and actively recruiting, this trial offers eligible patients the opportunity to receive potentially beneficial treatment options while contributing to research on colorectal cancer therapies. To participate, individuals must be 18 years or older, have metastatic colorectal cancer, and meet specific health criteria, including not being pregnant prior to treatment.

KRAS

A Study of Amivantamab and FOLFIRI Versus Cetuximab/Bevacizumab and FOLFIRI in Participants With KRAS/NRAS and BRAF Wild-type Colorectal Cancer Who Have Previously Received Chemotherapy

Phase: PHASE3 · Status: RECRUITING · [NCT06750094](#)

This clinical trial is testing the efficacy of amivantamab combined with FOLFIRI chemotherapy in patients with KRAS/NRAS and BRAF wild-type colorectal cancer who have previously undergone chemotherapy. The intervention involves the biological agent amivantamab alongside standard chemotherapy agents, which is important for evaluating its potential benefits compared to existing treatment options. This is a Phase 3 trial currently recruiting 700 participants, which means that eligible patients may have the opportunity to receive a treatment that is being compared to standard therapies in a larger population. Eligibility criteria include specific biomarker requirements related to KRAS, NRAS, and BRAF status, as well as other factors such as microsatellite instability (MSI) and mismatch repair (MMR) status.

ctDNA

ctDNA-guided Selection of Adjuvant Chemotherapy Regimens for Elderly Colon Cancer Patients After Surgery: A Single-center, Randomized, Controlled Study

Phase: PHASE3 · Status: RECRUITING · [NCT06609551](#)

This clinical trial is testing the effectiveness of ctDNA-guided selection of adjuvant chemotherapy regimens in elderly patients with high-risk stage II and stage III colon cancer following surgery. Participants will receive either 6 months of 5-FU monotherapy or an intensive XELOX treatment regimen, with the aim of improving disease-free survival outcomes. Currently in Phase 3 and actively recruiting, this trial seeks to determine how ctDNA detection can inform treatment decisions and patient prognosis. Eligibility criteria include being an elderly patient with high-risk stage II or stage III colon cancer and undergoing ctDNA testing to assess disease status.

HER2

A Study of Tucatinib With Trastuzumab and mFOLFOX6 Versus Standard of Care Treatment in First-line HER2+ Metastatic Colorectal Cancer

Phase: PHASE3 · Status: RECRUITING · [NCT05253651](#)

This clinical trial is testing the efficacy of tucatinib in combination with trastuzumab and mFOLFOX6 compared to standard of care treatments for patients with HER2 positive metastatic colorectal cancer. The intervention involves the use of tucatinib, a

targeted therapy, alongside other established cancer drugs, to evaluate its effectiveness and side effects in this patient population. Currently in Phase 3 and actively recruiting, this trial aims to provide insights into progression-free survival outcomes for participants. Eligibility criteria include having HER2 positive colorectal cancer that is metastatic or unresectable, with specific biomarker assessments for HER2 and RAS.

RAS

Neoadjuvant FOLFOX Chemotherapy for Patients With Locally Advanced Colon Cancer

Phase: PHASE3 · Status: ACTIVE_NOT_RECRUITING · [NCT03426904](#)

This clinical trial is testing the effectiveness of neoadjuvant FOLFOX chemotherapy in patients with locally advanced colon cancer. The intervention involves administering FOLFOX chemotherapy before surgery, which may enhance treatment outcomes by targeting microscopic metastatic cancer cells earlier and reducing surgical complications. This is a Phase 3 trial that is currently active but not recruiting, indicating that it is in the final stages of testing and results may soon be available for clinical application. Eligibility criteria include the presence of RAS biomarkers, which may be relevant for patient selection.

NGS

Induction Chemotherapy for Locally Recurrent Rectal Cancer

Phase: PHASE3 · Status: RECRUITING · [NCT04389086](#)

This clinical trial is testing the effectiveness of induction chemotherapy followed by neoadjuvant chemoradiotherapy and surgery in patients with locally recurrent rectal cancer. The intervention involves a combination of drugs, radiation, and surgical procedures, which may improve surgical outcomes by increasing the likelihood of achieving clear resection margins. Currently in phase III and actively recruiting participants, this trial aims to enroll 364 patients, which may provide valuable insights into treatment options for this patient population. Eligibility criteria include the requirement for next-generation sequencing (NGS) biomarkers.

What this means for you

Ask your oncologist whether any active trials match your biomarker profile and disease stage. The most actively trialed biomarkers in colorectal cancer right now are **MSI/MMR**, **BRAF**, **KRAS**, **ctDNA**, and **HER2**. If your tumor has not been tested for these markers, ask your oncologist before your next treatment decision — biomarker results can determine which trials you qualify for.

Questions to ask your oncologist about trials

"Am I eligible for any active Phase III trials based on my biomarker profile?"

"Does my tumor need to be retested before I can qualify for a biomarker-directed trial?"

"What are the travel requirements and visit frequency for the trials I might qualify for?"

Source: ClinicalTrials.gov · 2026-05-02 10:17:00

This is not medical advice. Always speak with your oncologist before making any treatment decisions.

Appendix B — Clinical Trial Reference Guide

What the phases mean

Phase I — First testing in humans. Focus on safety and dosing. Small groups.

Phase II — Testing effectiveness. Larger groups. Determines if Phase III is warranted.

Phase III — Randomized comparison against standard treatment. Required for FDA approval.

Key biomarkers that determine trial eligibility

RAS (KRAS/NRAS) — mutation status affects anti-EGFR eligibility and KRAS G12C trial eligibility

BRAF V600E — required for encorafenib-based trials (~8–10% of CRC)

MSI-H/dMMR — required for immunotherapy trials; ~5% of mCRC, ~15% of Stage II/III

HER2 — required for HER2-directed trials (~2–3% of CRC)

NTRK fusion — required for TRK inhibitor trials (<1% of CRC)

KRAS G12C — required for sotorasib/adagrasib trials (~3–4% of CRC)

TMB-H — tumor mutational burden; may qualify for immunotherapy

ctDNA — liquid biopsy; used in MRD monitoring trials

How to find trials you may qualify for

1. Get complete biomarker testing: RAS, BRAF, MSI/MMR, HER2, NTRK, TMB
2. Search ClinicalTrials.gov with your biomarker and stage
3. Ask your oncologist about NCI-designated cancer centers with trial access
4. Contact the Colorectal Cancer Alliance: ccalliance.org

Key Resources

ClinicalTrials.gov · [NCI Cancer Information \(cancer.gov\)](http://cancer.gov) · [Colorectal Cancer Alliance \(ccalliance.org\)](http://ccalliance.org) · [Fight Colorectal Cancer \(fightcrc.org\)](http://fightcrc.org)

Colon & Colorectal Cancer — Clinical Trials Intelligence

About This Report

This report is generated automatically from publicly available medical databases including PubMed, ClinicalTrials.gov, and the FDA Drug@FDA database. Entries are classified using deterministic rule-based logic and scored by clinical significance. AI-generated summaries are grounded exclusively in the source data provided.

Limitations

This report does not represent a complete picture of all published research or approved treatments. Database coverage, feed availability, and relevance filtering affect what appears. New drug approvals and trial status changes may not be reflected immediately. Always verify current information at the original source before making clinical decisions.

Disclaimer

Trial data is sourced from ClinicalTrials.gov, a registry maintained by the U.S. National Library of Medicine. Entry summaries are AI-generated from publicly available trial records. Trial status and eligibility criteria may change — always verify at ClinicalTrials.gov before acting on this information. This report is made available under a free-use license for patient advocacy organizations, oncology practices, and individual patients. You may share or distribute it freely. You may not modify it, remove attribution, or republish it as your own work. Authorship and all intellectual property rights are retained by Randy Hinton. This report is provided for informational purposes only and does not constitute medical advice. Always consult your oncologist or a qualified healthcare provider before making any treatment decisions.

Copyright

© 2026 Randy Hinton. All rights reserved. This report may be shared freely for informational purposes. It may not be modified, republished, or presented as original work without attribution.

https://petroai1492.github.io/delta/timelines/coloncancer_clinicaltrials_timeline.pdf