

Colon & Colorectal Cancer “ Clinical Trial & Treatment Timeline

Alert Window: Last 30 Days · Generated: 2026-03-11 11:01:31

Clinical Summary

Recent developments in colorectal cancer research highlight several promising treatment options for patients, particularly those with advanced stages. Notably, the FDA has approved Pfizer's Braftovi combination for first-line treatment of Stage IV metastatic colorectal cancer with BRAF mutations, signaling a significant advancement for this patient population. Additionally, multiple clinical trials are underway exploring innovative combinations of therapies, such as dual-target CAR-NK cells and various chemotherapy regimens, which may improve outcomes for patients with unresectable or metastatic disease. Moving forward, it will be important to monitor the results of these trials and their implications for treatment protocols in colorectal cancer.

AI-generated from deterministic classification data · 2026-03-11 11:01:31

Situation Summary

Events Detected: 16
High Significance: 2
Top Category: New Trial
Primary Stage: Stage IV
Primary Target: General
Research Activity: Active Monitoring

Event Timeline

EVENT CLUSTER “ Dual-Target CAR-NK Cells for Biomarker-Selected Advanced Colorectal Cancer

MEDIUM

STAGE IV

PD-1/PD-L1

NEW TRIAL

CORROBORATED

PUBLISHED MAR 10

Mar 10 18:35 · immediate · [clinicaltrials.gov](#), [clinicaltrials.gov](#), [clinicaltrials.gov](#)

A new clinical trial is exploring the use of dual-target CAR-NK cells in patients with advanced colorectal cancer, potentially offering a promising treatment option for those whose tumors express specific biomarkers.

3 source(s) reporting · cluster 1 of 11

EVENT CLUSTER â€” A Clinical Study of Iparomlimab and Tuvonralimab Combined With Bevacizumab and Alternating Triweekly CAPOX/mCAPIRI Regimen as First-line Treatment for Unresectable Advanced Colorectal Cancer

MEDIUM

STAGE IV

VEGF

NEW TRIAL

CORROBORATED

PUBLISHED MAR 3

Mar 10 18:32 Â· immediate Â· [clinicaltrials.gov](#), [clinicaltrials.gov](#)

A new clinical trial is exploring the combination of Iparomlimab and Tuvonralimab with Bevacizumab and a specific chemotherapy regimen as a first-line treatment for patients with unresectable advanced colorectal cancer, potentially offering a new option for improved outcomes.

2 source(s) reporting Â· cluster 2 of 11

EVENT CLUSTER â€” FOLFOX Chemotherapy Combined With Fruquintinib and Serplulimab as First-Line Conversion Therapy for Initially Unresectable pMMR/MSS Colorectal Cancer

MEDIUM

STAGE IV

GENERAL

NEW TRIAL

CORROBORATED

PUBLISHED MAR 3

Mar 10 18:32 Â· immediate Â· [clinicaltrials.gov](#), [clinicaltrials.gov](#)

A new clinical trial is exploring the combination of FOLFOX chemotherapy with Fruquintinib and Serplulimab as a promising first-line treatment option for patients with initially unresectable stage IV colorectal cancer, potentially improving their chances for surgical conversion.

2 source(s) reporting Â· cluster 3 of 11

EVENT CLUSTER â€” FDA Approves Pfizerâ€™s Braftovi Combination for First-Line Metastatic Colorectal Cancer

HIGH

STAGE IV

BRAF

TARGETED THERAPY

CORROBORATED

PUBLISHED FEB 25

Mar 10 18:26 Â· immediate Â· [Contract Pharma](#), [Oncology News Central](#)

The FDA has approved Pfizer's Braftovi combination therapy for first-line treatment of Stage IV metastatic colorectal cancer with BRAF mutations, offering new hope for improved outcomes in this challenging stage of the disease.

2 source(s) reporting Â· cluster 4 of 11

EVENT CLUSTER â€” A Phase â€¦ Clinical Trial of RC148 Plus Chemotherapy as 1L Therapy for Unresectable or Metastatic Colorectal Cancer

MEDIUM

STAGE IV

GENERAL

NEW TRIAL

CORROBORATED

PUBLISHED MAR 10

Mar 10 18:26 Â· immediate Â· [clinicaltrials.gov](#)

A new clinical trial is testing the combination of RC148 and chemotherapy as a first-line treatment for patients with unresectable or metastatic colorectal cancer, potentially offering a new option for improving outcomes in this advanced stage of the disease.

1 source(s) reporting Â· cluster 5 of 11

EVENT CLUSTER “ Adding Encorafenib and Cetuximab to FOLFIRI Shows Benefit in BRAF-Mutant Colorectal Cancer

MEDIUM

ALL STAGES

BRAF

TRIAL RESULTS

VERIFIED

PUBLISHED FEB 25

Mar 9 21:21 · immediate · The ASCO Post

Combining Encorafenib and Cetuximab with the chemotherapy regimen FOLFIRI has been shown to improve treatment outcomes for patients with BRAF-mutant colorectal cancer.

1 source(s) reporting · cluster 6 of 11

EVENT CLUSTER “ Low-Dose Radiotherapy to Sensitize Pucotenlimab Plus CAPEOX for pMMR Locally Advanced Rectal Cancer

MEDIUM

STAGE III

GENERAL

NEW TRIAL

CORROBORATED

PUBLISHED MAR 4

Mar 9 21:21 · immediate · clinicaltrials.gov

A new clinical trial is exploring the use of low-dose radiotherapy to enhance the effectiveness of the drug pucotenlimab combined with CAPEOX chemotherapy for patients with locally advanced rectal cancer, potentially improving treatment outcomes.

1 source(s) reporting · cluster 7 of 11

EVENT CLUSTER “ Everolimus in CDK12-Deficient Metastatic Colorectal Cancer (EVER-RECODE)

MEDIUM

STAGE IV

GENERAL

NEW TRIAL

CORROBORATED

PUBLISHED FEB 27

Mar 9 21:21 · immediate · clinicaltrials.gov

The EVER-RECODE trial is exploring the use of Everolimus for treating Stage IV metastatic colorectal cancer with CDK12 deficiency, potentially offering a new targeted treatment option for patients with this specific genetic profile.

1 source(s) reporting · cluster 8 of 11

EVENT CLUSTER “ Effect of Perioperative Lidocaine or High-Dose Dexamethasone on Immune Response in Colon Cancer Surgery (PILDI Study)

MEDIUM

STAGE III

GENERAL

NEW TRIAL

CORROBORATED

PUBLISHED MAR 6

Mar 9 21:21 · immediate · clinicaltrials.gov

The PILDI Study is investigating whether administering perioperative lidocaine or high-dose dexamethasone during colon cancer surgery can enhance the immune response, potentially improving outcomes for patients with Stage III colorectal cancer.

1 source(s) reporting · cluster 9 of 11

EVENT CLUSTER " Neadjuvant Chemoradiotherapy Plus Tislelizumab With or Without Probio-M9 in pMMR/MSS Locally Advanced Rectal Cancer

MEDIUM

STAGE III

GENERAL

NEW TRIAL

CORROBORATED

PUBLISHED MAR 9

Mar 9 21:21 · immediate · clinicaltrials.gov

A new clinical trial is exploring the combination of neoadjuvant chemoradiotherapy with the immunotherapy drug Tislelizumab, potentially enhancing treatment options for patients with locally advanced rectal cancer.

1 source(s) reporting · cluster 10 of 11

EVENT CLUSTER " FOLFIRI and Bevacizumab With or Without Pelareorep for Second-Line Treatment of Metastatic RAS-Mutated, Microsatellite-Stable Colorectal Cancer

MEDIUM

STAGE IV

VEGF

NEW TRIAL

CORROBORATED

PUBLISHED MAR 4

Mar 9 21:21 · immediate · clinicaltrials.gov

A new clinical trial is exploring whether adding Pelareorep to the standard treatment of FOLFIRI and Bevacizumab can improve outcomes for patients with Stage IV metastatic RAS-mutated, microsatellite-stable colorectal cancer.

1 source(s) reporting · cluster 11 of 11

Appendix A – Patient-Friendly Summary

Plain-language overview for patients, caregivers, and family members

What's happening

Recently, there have been some exciting developments in colorectal cancer research. A new clinical trial is testing a treatment that uses special immune cells to target advanced colorectal cancer. Another trial is looking at a combination of different drugs to help patients whose cancer cannot be surgically removed. The FDA has also approved a new therapy for patients with a specific type of metastatic colorectal cancer. Other trials are exploring various combinations of chemotherapy and new drugs to improve outcomes for patients at different stages of the disease. These advancements offer new hope and options for many people facing colorectal cancer.

What this means for you

For patients, these developments could mean more personalized treatment options. If you have advanced colorectal cancer, ask your oncologist about the new trials and therapies that might be suitable for you. It's important to discuss your specific situation and any genetic factors that could influence your treatment. Keep an eye out for new clinical trials and ask about any potential benefits they could offer. Remember, you are not alone in this journey, and there are many people working hard to find better treatments for colorectal cancer.

AI-generated in plain language from clinical data · 2026-03-11 11:01:31

This is not medical advice. Always speak with your oncologist before making any treatment decisions.

Appendix B – Colon & Colorectal Cancer Treatment Reference

Incidence & Trends

~150,000 new US cases/year ~52,000 deaths/year 4th most common US cancer

▲ Early-Onset Alert: CRC rates in adults under 50 have risen ~2% per year since the 1990s Adults under 55 now account for ~20% of all new CRC diagnoses Cause under active investigation (diet, microbiome, obesity, sedentary lifestyle)

Current Screening Guidelines (2025)

USPSTF / ACS / ACG – Average Risk: Begin screening at age 45 (lowered from 50 in 2021)

Colonoscopy every 10 years FIT (fecal immunochemical test) annually Cologuard every 1-3 years CT colonography every 5 years

High Risk (family history / Lynch syndrome / IBD): Begin at age 40 or 10 years before youngest affected relative – whichever comes first Colonoscopy every 1-5 years depending on risk

Symptoms at any age: Rectal bleeding, change in bowel habits >4 weeks, unexplained weight loss, iron deficiency anemia – immediate evaluation regardless of age

Key Biomarkers – Test All mCRC at Diagnosis

RAS (KRAS/NRAS) BRAF V600E MSI/MMR status HER2 NTRK fusion TMB

Stage IV Targeted Options

KRAS G12C (~3-4%) – sotorasib + cetuximab adagrasib + cetuximab

BRAF V600E (~8-10%) – encorafenib + cetuximab ± binimetinib

MSI-H/dMMR (~5%) – pembrolizumab (1st line preferred) nivolumab ± ipilimumab

HER2 amplified (~2-3%) – trastuzumab + pertuzumab trastuzumab deruxtecan

RAS/BRAF WT – FOLFOX or FOLFIRI + cetuximab or panitumumab (left-sided preferred)

1st Line mCRC Standard of Care

FOLFOX or FOLFIRI ± bevacizumab (RAS mutant / right-sided)

FOLFOX or FOLFIRI + cetuximab or panitumumab (RAS/BRAF WT, left-sided)

Pembrolizumab monotherapy (MSI-H/dMMR)

Key Resources

ClinicalTrials.gov NCCN Guidelines (nccn.org) Colorectal Cancer Alliance (ccalliance.org) Fight Colorectal Cancer (fightcrc.org) ACS (cancer.org)

Colon & Colorectal Cancer – Clinical Trial & Treatment Timeline

Alert classification fields are generated using deterministic rule-based analysis. Alert summaries are AI-generated from publicly available news headlines. This report aggregates headline-level information for situational awareness only. No article text is reproduced. All original content remains the property of respective publishers. This report is provided for informational purposes only and does not constitute medical advice. Always consult your oncologist or a qualified healthcare provider before making

any treatment decisions.

© 2026 Randy Hinton. All rights reserved. · https://petroai1492.github.io/delta/coloncancer_timeline.pdf