Colon & Colorectal Cancer — Biomarker Testing Intelligence

Alert Window: Last 30 Days · Generated: 2026-11-12 07:30:01

Situation Summary

Stage IV: 9 of 38
Biomarker Count: 23 entries
Top Targets: ctDNA (10) · BRAF (4) · MSI/MMR (3)
Breakthrough: 18 entries scored ≥ 4

Research & Guidelines Timeline

Biomarker testing entries ranked by significance score. Higher scores indicate greater clinical relevance. Always discuss with your oncologist.

HIGH All Stages ctDNA VERIFIED LIQUID BIOPSY PUBLISHED NOV 11, 2026 10

Targeted liquid biopsy using CEUS to facilitate the release of ctDNA in colorectal liver metastasis

pubmed.ncbi.nlm.nih.gov

HIGH Stage III BROAD SCOPE VERIFIED LIQUID BIOPSY PUBLISHED OCT 30, 2026 6

Targeting minimal residual disease (MRD) in colorectal cancer: Therapeutic strategies, clinical trials, and the path toward molecular cure

pubmed.ncbi.nlm.nih.gov

HIGH All Stages ctDNA VERIFIED LIQUID BIOPSY PUBLISHED OCT 28, 2026 7

Risk stratification by circulating tumor DNA-based pretreatment genomic profile and post-operative molecular residual disease detection in resectable colorectal oligometastases: PRECISION, a prospective multicenter study

pubmed.ncbi.nlm.nih.gov

HIGH Stage IV ctDNA VERIFIED LIQUID BIOPSY PUBLISHED OCT 15, 2026 7

ctDNA retesting to guide anti-EGFR therapy in later-line metastatic colorectal cancer

pubmed.ncbi.nlm.nih.gov

HIGH All Stages ctDNA VERIFIED LIQUID BIOPSY PUBLISHED NOV 02, 2026 6

Circulating Tumor DNA and Cell-Free DNA as Candidate Biomarkers After Image-Guided Locoregional Therapy in Hepatocellular Carcinoma and Colorectal Liver Metastases: A Systematic Review

pubmed.ncbi.nlm.nih.gov

HIGH | All Stages | ctDNA | VERIFIED | LIQUID BIOPSY | PUBLISHED OCT 26, 2026 | 6

Circulating Tumor DNA (ctDNA) for Minimal Residual Disease (MRD) Detection in Colorectal Cancer (CRC): Current Evidence, Clinical Applications, and Future Directions

pubmed.ncbi.nlm.nih.gov

MEDIUM | Stage IV | BRAF | VERIFIED | GENOMIC PROFILING | PUBLISHED OCT 23, 2026 | 6

Prognostic Modeling Integrating RAS and SMAD4 Status in Patients with BRAF Wild-Type, Microsatellite-Stable Metastatic Colorectal Cancer

pubmed.ncbi.nlm.nih.gov

MEDIUM | Stage II | BROAD SCOPE | VERIFIED | RESEARCH | PUBLISHED OCT 31, 2026 | 4

The impact of PROgnostic SIgnatures in stage II colon cancer patients enrolled in the phase III clinical trial TOSCA: the PROSIT study

pubmed.ncbi.nlm.nih.gov

MEDIUM | Early Detection | MSI/MMR | VERIFIED | GENOMIC PROFILING | PUBLISHED OCT 30, 2026 | 3

A novel predictive model for DNA mismatch repair deficient colorectal carcinoma for cost-effective MMR testing in low-resource settings: HERALD-MMR (Histological Evaluation and Risk Assessment using Local Demographics for Mismatch Repair)

pubmed.ncbi.nlm.nih.gov

MEDIUM | Stage IV | BROAD SCOPE | VERIFIED | RESEARCH | PUBLISHED OCT 26, 2026 | 3

Prognostic Significance of CD47 and DHX9 in Metastatic Rectal Cancer

pubmed.ncbi.nlm.nih.gov

MEDIUM | Stage IV | ctDNA | VERIFIED | RESEARCH | PUBLISHED OCT 22, 2026 | 3

Irinotecan Plus Cetuximab Rechallenge Versus Trifluridine/Tipiracil Plus Bevacizumab as Third-Line Treatment in Circulating Tumor DNA-Selected Metastatic Colorectal Cancer (ROMANCE Trial)

pubmed.ncbi.nlm.nih.gov

MEDIUM | All Stages | NGS | VERIFIED | GENOMIC PROFILING | PUBLISHED NOV 05, 2026 | 2

Panel-based next-generation sequencing with copy number variant detection pipeline could increase the detection rate of hereditary colorectal cancer syndrome compared with Sanger sequencing

pubmed.ncbi.nlm.nih.gov

MEDIUM | All Stages | NGS | VERIFIED | GENOMIC PROFILING | PUBLISHED OCT 16, 2026 | 2

Laboratory-Developed NGS Panel for Characterization of Advanced Colorectal Cancer, Real-World Data from a Tertiary Referral Centre

pubmed.ncbi.nlm.nih.gov

MEDIUM All Stages RAS VERIFIED RESEARCH PUBLISHED OCT 17, 2026 1

RAS Mutations as Prognostic Biomarkers in Digestive Cancers: From Binary Classification to Variant-Specific and Context-Dependent Risk Stratification

pubmed.ncbi.nlm.nih.gov

Appendix A — Patient-Friendly Summary

Plain-language overview for patients, caregivers, and family members

What's happening

Biomarker testing — also called molecular profiling or genomic testing — analyzes your tumor's DNA to identify specific mutations. The results determine which FDA-approved treatments may be available to you. The entries below represent the highest-scoring study per biomarker target drawn from the full research archive — not limited to the current report window. Some entries may predate the timeline above; they are included because they remain the most clinically significant reference available for that biomarker.

ctDNA

Circulating tumor DNA tumor fraction as a biomarker in refractory metastatic colorectal cancer: findings from the CAVE-2 GOIM trial

In the CAVE-2 GOIM trial, circulating tumor DNA (ctDNA) was assessed using the FoundationOne Liquid assay to evaluate its role in guiding anti-EGFR rechallenge therapy with cetuximab ± avelumab in refractory metastatic colorectal cancer patients. The study highlights the need for biomarkers to stratify patients, particularly focusing on those with RAS/BRAF wild type mCRC.

MSI/MMR

Final results of nivolumab plus ipilimumab versus nivolumab or chemotherapy in microsatellite instability-high/mismatch repair-deficient metastatic colorectal cancer from CheckMate 8HW

KRAS

Circulating tumor DNA (ctDNA) clearance as an early indicator of sotorasib + panitumumab efficacy and prognosis in KRAS G12C–mutated colorectal cancer (mCRC): Results from phase 3 CodeBreaK 300.

In the phase 3 CodeBreaK 300 trial, ctDNA clearance was assessed as an early indicator of treatment efficacy and prognosis in KRAS G12C–mutated mCRC patients receiving sotorasib and panitumumab. The findings suggest that ctDNA clearance may serve as a valuable biomarker for monitoring therapeutic response and predicting outcomes in this patient population.

TP53

Clinical impact of TP53 mutation status on survival outcomes in metastatic colorectal cancer

TP53 mutations serve as an independent prognostic marker for prolonged progression-free survival in metastatic colorectal cancer. While overall survival differences were not statistically significant, the findings suggest the need for further validation in prospective studies to assess TP53's role in therapeutic stratification.

BRAF

Detection of KRAS, NRAS and BRAF Mutations in Liquid Biopsy from Patients with Colorectal Cancer

ddPCR analysis of circulating tumor DNA (ctDNA) in colorectal cancer patients reveals that it can provide complementary information for molecular diagnosis. The study highlights the utility of liquid biopsy in detecting KRAS, NRAS, and BRAF mutations, enhancing diagnostic accuracy in this patient population.

HER2

HER2-Positive Colorectal Cancer-from Diagnosis to Anti-HER2 Therapy

NGS

Distinct NGS mutational landscape and prognostic implications in early-onset colorectal cancer: A dual-cohort analysis of TCGA PanCancer Atlas and MSK-IMPACT 50K

EGFR

Second-line raltitrexed plus irinotecan (SALIRI) with bevacizumab, cetuximab, or alone for metastatic colorectal cancer: a prospective multicenter cohort-based secondary analysis of the SALLY trial

SALIRI, a combination of raltitrexed and irinotecan, demonstrated feasibility and tolerability as a second-line therapy for metastatic colorectal cancer when combined with bevacizumab or cetuximab in a nonrandomized secondary analysis of the SALLY trial. Favorable outcomes were noted in the SALIRI + cetuximab subgroup, though results should be interpreted with caution due to nonrandom treatment allocation. Further prospective, biomarker-driven randomized studies are needed to clarify the optimal use of SALIRI-based combinations.

RAS

The temporal evaluation of RAS mutation by liquid biopsy at progression after bevacizumab-based treatment in patients with metastatic colorectal cancer

Dynamic changes in plasma RAS mutation status during disease progression in metastatic colorectal cancer (mCRC) were observed, indicating tumor clonal evolution. Notably, RAS mutation clearance correlated with a survival advantage, suggesting prognostic relevance. Reassessing RAS status via liquid biopsy at progression may yield clinically relevant insights, though findings are exploratory and warrant further investigation.

Genomic Profiling

Integrated Molecular Profiling of Colorectal Cancer by Tumor Location: Evidence from a Real-World Cohort with Primary and Metastatic Samples

Colorectal cancer (CRC) tumor location correlates with distinct molecular profiles. Right-sided tumors exhibit dMMR, MSI-H, and elevated TMB, while left-sided tumors show increased ERBB2 alterations and class 5 APC mutations. These findings underscore the necessity of incorporating tumor location into personalized molecular diagnostics and treatment strategies for CRC patients.

Methylation

Lactate as a Master Regulator of Immune Suppression: From Metabolic Waste to Epigenetic Checkpoint in Colorectal Cancer

Lactate, a byproduct of aerobic glycolysis driven by KRAS and BRAF mutations in microsatellite-stable colorectal cancer, acts as a central immunosuppressive regulator in the tumor microenvironment. Its accumulation leads to immune suppression, contributing to resistance against immune checkpoint inhibitors. This highlights lactate's role beyond metabolic waste, suggesting potential therapeutic targets to enhance immune response in this cancer subtype.

TMB

Co-Mutation of CREBBP/EP300 and POLE/POLD1 Identifies a TMB-High Subset of MSS CRC

Co-mutation of CREBBP/EP300 and POLE/POLD1 identifies a subset of TMB-high microsatellite stable (MSS) colorectal cancer (CRC). Analysis of multiple public cohorts demonstrated that CREBBP/EP300 mutations are independently associated with TMB-high status ($p < 0.00001$). This finding suggests potential for these mutations as biomarkers for immunotherapy eligibility in MSS CRC.

miRNA

Diagnostic Performance of Circulating miRNA-92a and Related Circulating miRNA Panels for Colorectal Cancer: An Updated Systematic Review and Meta-Analysis

Circulating miRNA-92a shows superior diagnostic performance as a non-invasive biomarker for early colorectal cancer detection. The systematic review and meta-analysis suggest its potential utility, though further large-scale prospective studies are needed to standardize testing protocols and confirm clinical applicability.

Stool DNA

Enhancing non-invasive colorectal cancer screening with stool DNA methylation markers and light gradient-boosting machine learning

Stool DNA methylation markers CNRIP1, SFRP2, and VIM demonstrated high sensitivity for non-invasive colorectal cancer screening. The application of the LightGBM machine learning algorithm improved diagnostic accuracy, suggesting a viable method for early CRC detection.

IHC

Integration of deep learning and radiomic features from multiplex immunohistochemistry images for reproducible Multi-Outcome prediction in a Multi-Center study of colorectal cancer

Fused multiplex immunohistochemistry (mIHC)-derived radiomic and deep learning features provide accurate and interpretable predictions for multiple colorectal cancer outcomes. The study supports the integration of these features into precision oncology workflows, enhancing the potential for reproducible multi-outcome predictions across different centers.

What this means for you

If you have been diagnosed with colorectal cancer, ask your oncologist: *"Has my tumor been tested for all the biomarkers that have an FDA-approved treatment?"* This is especially important if you have been diagnosed with metastatic (Stage IV) colorectal cancer, where biomarker results directly determine which targeted therapies and immunotherapies are available to you. Based on the research in this report, the most actively studied biomarkers right now are **ctDNA, MSI/MMR, KRAS, TP53, and BRAF**. Testing should happen at diagnosis — before treatment begins — because results take 1–2 weeks and can shape your first treatment decision.

Generated from structured biomarker research data · 2026-11-12 07:30:01
This is not medical advice. Always speak with your oncologist before making any treatment decisions.

Appendix B — Colon & Colorectal Cancer: Biomarker Testing Reference

What Is Biomarker Testing?

Biomarker testing — also called molecular profiling or genomic testing — analyzes your tumor's DNA to identify specific mutations. A single next-generation sequencing (NGS) panel can check many gene mutations at once. Results typically take 1–2 weeks and should be obtained at diagnosis, before treatment begins.

Key Biomarkers — Test All mCRC at Diagnosis

RAS (KRAS/NRAS) · BRAF V600E · MSI/MMR status · HER2 · NTRK fusion · TMB

Testing Methods

NGS (next-generation sequencing) — checks many mutations at once from tumor tissue

Liquid biopsy / ctDNA — blood test that detects tumor DNA; useful when tissue is unavailable

IHC (immunohistochemistry) — used to test MMR status and HER2

PCR — used for specific mutation detection (e.g. MSI, KRAS)

Questions to Ask Your Oncologist

"Has my tumor been tested for all the biomarkers that have an FDA-approved treatment?"

"Was my sample taken from the primary tumor or a metastatic site — and does it need to be retested?"

"How long will results take, and will they affect my first treatment decision?"

"Is a liquid biopsy appropriate for my situation?"

Abbreviations

BRAF V600E — a specific gene mutation found in ~8–10% of colorectal cancers

ctDNA — circulating tumor DNA; tumor DNA fragments found in the bloodstream

dMMR — mismatch repair deficient; indicates the tumor may respond to immunotherapy

HER2 — a protein that drives growth in ~2–3% of colorectal cancers

IHC — immunohistochemistry; a lab staining method used to detect proteins in tissue

KRAS G12C — a specific gene mutation found in ~3–4% of colorectal cancers

mCRC — metastatic colorectal cancer (Stage IV)

MSI-H — microsatellite instability-high; indicates the tumor may respond to immunotherapy

NGS — next-generation sequencing; a method that checks many gene mutations at once

NTRK — a rare gene fusion found in <1% of colorectal cancers

RAS — a gene family (KRAS/NRAS) whose mutation status affects treatment selection

TMB — tumor mutational burden; a measure of how many mutations are in a tumor

Key Resources

ClinicalTrials.gov · NCCN Guidelines (nccn.org) · Colorectal Cancer Alliance (ccalliance.org) · Fight Colorectal Cancer (fightcrc.org) · ACS (cancer.org)

Colon & Colorectal Cancer — Biomarker Testing Intelligence

About This Report

This report is generated automatically from publicly available medical databases including PubMed, ASCO Journal of Clinical Oncology, Annals of Oncology, and NCI Cancer.gov. Entries are classified using deterministic rule-based logic and scored by clinical significance. AI-generated summaries are grounded exclusively in the source data provided.

Limitations

This report does not represent a complete picture of all published research. Database coverage, feed availability, and relevance filtering affect what appears. Research findings may not yet reflect current clinical guidelines. Always verify information at the original source before making clinical decisions.

Disclaimer

Alert classification fields are generated using deterministic rule-based analysis. Entry summaries are AI-generated from publicly available PubMed abstracts and society publications maintained by the U.S. National Library of Medicine and professional oncology societies. This report aggregates publicly available data for situational awareness only. No source content is reproduced verbatim. This report is made available under a free-use license for patient advocacy organizations, oncology practices, and individual patients. You may share or distribute it freely. You may not modify it, remove attribution, or republish it as your own work. Authorship and all intellectual property rights are retained by Randy Hinton. This report is provided for informational purposes only and does not constitute medical advice. Always consult your oncologist or a qualified healthcare provider before making any treatment decisions.

Copyright

© 2026 Randy Hinton. All rights reserved. This report may be shared freely for informational purposes. It may not be modified, republished, or presented as original work without attribution. ·
https://petroai1492.github.io/delta/timelines/coloncancer_biomarker_timeline.pdf