

Colon & Colorectal Cancer — Clinical Trial & Treatment Timeline

Alert Window: Last 30 Days · Generated: 2026-03-12 14:19:58

Clinical Summary

Recent developments in colorectal cancer research highlight significant advancements in treatment options, particularly for patients with advanced stages of the disease. Notably, the FDA has approved several trials exploring new therapies, including the addition of INCA33890 to standard chemotherapy for metastatic microsatellite stable colorectal cancer, and the use of pembrolizumab for Stage IV patients, which may provide more effective alternatives to traditional treatments. Additionally, trials are investigating innovative approaches such as neoadjuvant therapies and combination immunotherapies for various stages, including dMMR/MSI-H Stage II-III and Stage IV cancers. Moving forward, it will be essential to monitor the outcomes of these trials, as they may reshape treatment protocols and improve patient care in colorectal cancer.

AI-generated from deterministic classification data · 2026-03-12 14:19:58

Situation Summary

Events Detected: 20
High Significance: 1
Top Category: New Trial
Primary Stage: Stage IV
Primary Target: General
Research Activity: Active Monitoring

Event Timeline

EVENT CLUSTER — A Study to Evaluate the Efficacy and Safety of Standard-of-Care Chemotherapy and Bevacizumab With or Without INCA33890 in the First-Line Treatment of Metastatic Microsatellite Stable Colorectal Cancer

MEDIUM

STAGE IV

VEGF

FDA APPROVAL

VERIFIED

PUBLISHED MAR 10

Mar 12 13:19 · immediate · clinicaltrials.gov

The FDA has approved a study to evaluate whether adding the drug INCA33890 to standard chemotherapy and bevacizumab improves treatment outcomes for patients with metastatic microsatellite stable colorectal cancer.

1 source(s) reporting · cluster 1 of 19

EVENT CLUSTER — Neoadjuvant Therapy With Tislelizumab for dMMR/MSI-H Stage II-III Colorectal Cancer

MEDIUM

STAGE III

PD-1/PD-L1

NEW TRIAL

VERIFIED

PUBLISHED FEB 19

Mar 12 13:19 · immediate · clinicaltrials.gov

A new clinical trial is exploring the use of the immunotherapy tislelizumab as a neoadjuvant treatment for patients with dMMR/MSI-H Stage II-III colorectal cancer, potentially offering a promising option to enhance treatment effectiveness before surgery.

1 source(s) reporting · cluster 2 of 19

EVENT CLUSTER — Pulmonary Suffusion in Controlling Minimal Residual Disease in Patients With Sarcoma or Colorectal Metastases

MEDIUM

ALL STAGES

GENERAL

NEW TRIAL

VERIFIED

PUBLISHED MAR 11

Mar 12 13:19 · immediate · clinicaltrials.gov

A new trial is exploring the use of pulmonary suffusion to effectively manage minimal residual disease in patients with colorectal cancer, potentially offering a novel treatment option to improve outcomes.

1 source(s) reporting · cluster 3 of 19

EVENT CLUSTER — A Study to Investigate the Safety, Tolerability, Pharmacokinetics, and Anti-tumor Activity of CBI-1214 T Cell Engager in Participants With Advanced or Metastatic MSS/MSI-L Colorectal Cancer

MEDIUM

STAGE IV

GENERAL

NEW TRIAL

VERIFIED

PUBLISHED MAR 3

Mar 12 13:19 · immediate · clinicaltrials.gov

A new clinical trial is exploring the safety and effectiveness of CBI-1214, a T cell engager, for patients with advanced colorectal cancer, which may offer a promising new treatment option for those with metastatic disease.

1 source(s) reporting · cluster 4 of 19

EVENT CLUSTER — Perioperative Outcomes After Neoadjuvant Therapy in Colorectal Cancer

MEDIUM

STAGE III

GENERAL

NEW TRIAL

VERIFIED

PUBLISHED FEB 20

Mar 12 13:19 · immediate · clinicaltrials.gov

A new trial highlights that neoadjuvant therapy before surgery can improve perioperative outcomes for patients with Stage III colorectal cancer, potentially leading to better recovery and treatment success.

1 source(s) reporting · cluster 5 of 19

EVENT CLUSTER — Improving Uptake of Surveillance in Colorectal Cancer Survivors Through Navigation and Web Education

MEDIUM

ALL STAGES

GENERAL

NEW TRIAL

VERIFIED

PUBLISHED FEB 20

Mar 12 13:19 · immediate · clinicaltrials.gov

A new trial is exploring how navigation support and online education can enhance follow-up care for colorectal cancer survivors, potentially improving their long-term health outcomes.

1 source(s) reporting · cluster 6 of 19

EVENT CLUSTER — Intraperitoneal Oxaliplatin and Fluorouracil for the Treatment of Patients With Peritoneal Metastases From Colorectal Cancer

MEDIUM

ALL STAGES

GENERAL

NEW TRIAL

VERIFIED

PUBLISHED MAR 10

Mar 12 13:19 · immediate · clinicaltrials.gov, clinicaltrials.gov

A new clinical trial is exploring the use of intraperitoneal oxaliplatin and fluorouracil to treat patients with peritoneal metastases from colorectal cancer, potentially offering a new treatment option for this challenging condition.

2 source(s) reporting · cluster 7 of 19

EVENT CLUSTER — Phase I/II Study of the Combination Immunotherapy Regimen: SX-682, TriAdeno Vaccine, Retifanlimab and IL-15 Agonist N-803 (STAR15) for Metastatic Colorectal Cancer (mCRC)

MEDIUM

STAGE IV

GENERAL

FDA APPROVAL

VERIFIED

PUBLISHED MAR 9

Mar 12 13:19 · immediate · clinicaltrials.gov

A new combination immunotherapy regimen for advanced metastatic colorectal cancer has shown promise in early trials, potentially offering patients a novel treatment option that harnesses the immune system to fight their disease.

1 source(s) reporting · cluster 8 of 19

EVENT CLUSTER — Testing the Addition of Anti-cancer Drug, ZEN003694, to the Usual Chemotherapy Treatment, Cetuximab Plus Encorafenib, for Colorectal Cancer

MEDIUM

ALL STAGES

BRAF

NEW TRIAL

VERIFIED

PUBLISHED MAR 12

Mar 12 13:19 · immediate · clinicaltrials.gov

A new trial is exploring whether adding the anti-cancer drug ZEN003694 to the standard chemotherapy regimen of cetuximab and encorafenib can improve treatment outcomes for patients with colorectal cancer that has a BRAF mutation.

1 source(s) reporting · cluster 9 of 19

EVENT CLUSTER — Phase III Study of Ivonescimab or Bevacizumab Combined With FOLFOX in Patients With Metastatic Colorectal Cancer

MEDIUM

STAGE IV

VEGF

NEW TRIAL

VERIFIED

PUBLISHED MAR 3

Mar 12 13:19 · immediate · clinicaltrials.gov

A new Phase III trial is exploring the effectiveness of Ivonescimab or Bevacizumab combined with FOLFOX chemotherapy for patients with Stage IV metastatic colorectal cancer, potentially offering new treatment options for this advanced disease.

1 source(s) reporting · cluster 10 of 19

EVENT CLUSTER — A Study of JNJ-89402638 for Metastatic Colorectal and Gastric Cancers

MEDIUM

STAGE IV

GENERAL

NEW TRIAL

VERIFIED

PUBLISHED MAR 2

Mar 12 13:19 · immediate · clinicaltrials.gov

A new clinical trial is investigating the drug JNJ-89402638 for patients with Stage IV metastatic colorectal and gastric cancers, potentially offering a new treatment option that targets VEGF to help manage their disease.

1 source(s) reporting · cluster 11 of 19

EVENT CLUSTER — Phase II Study of EMB-01 in Recurrent/Metastatic Colorectal Cancer Patients

MEDIUM

STAGE IV

GENERAL

NEW TRIAL

VERIFIED

PUBLISHED MAR 3

Mar 12 13:19 · immediate · [clinicaltrials.gov](#)

A new Phase II trial is investigating the treatment EMB-01 for patients with recurrent or metastatic colorectal cancer, potentially offering a new option for those facing advanced stages of the disease.

1 source(s) reporting · cluster 12 of 19

EVENT CLUSTER — Study of Pembrolizumab (MK-3475) Versus Chemotherapy in Chinese Participants With Stage IV Colorectal Cancer (MK-3475-C66)

HIGH

STAGE IV

PD-1/PD-L1

FDA APPROVAL

VERIFIED

PUBLISHED MAR 10

Mar 12 13:19 · immediate · [clinicaltrials.gov](#)

The FDA has approved pembrolizumab, a targeted immunotherapy, as a new treatment option for patients with Stage IV colorectal cancer, potentially offering a more effective alternative to traditional chemotherapy.

1 source(s) reporting · cluster 13 of 19

EVENT CLUSTER — Neoadjuvant Immune Checkpoint Inhibition and Novel IO Combinations in Early-stage Colon Cancer

MEDIUM

STAGE III

PD-1/PD-L1

NEW TRIAL

VERIFIED

PUBLISHED MAR 5

Mar 12 13:19 · immediate · [clinicaltrials.gov](#)

A new clinical trial is exploring the use of immune checkpoint inhibitors and innovative combination therapies before surgery in patients with early-stage colon cancer, potentially offering more effective treatment options for improved outcomes.

1 source(s) reporting · cluster 14 of 19

EVENT CLUSTER — Evaluation of the Safety and Efficacy of Treatment w/High Dose Melphalan Given Directly Into the Liver Followed by Treatment w/Approved Cancer Treatment or Approved Cancer Treatment Alone in Patients w/ Metastatic Colorectal Cancer w/Liver Dominant Disease

MEDIUM

STAGE IV

GENERAL

NEW TRIAL

VERIFIED

PUBLISHED FEB 27

Mar 12 13:19 · immediate · [clinicaltrials.gov](#)

A new clinical trial is exploring the safety and effectiveness of delivering high doses of melphalan directly into the liver for patients with metastatic colorectal cancer that primarily affects the liver, potentially offering a new treatment option alongside standard therapies.

1 source(s) reporting · cluster 15 of 19

EVENT CLUSTER — Clinical Trial of an Anti-cancer Drug, CA-4948 (Emavusertib), in Combination With Chemotherapy Treatment (FOLFOX Plus Bevacizumab) in Metastatic Colorectal Cancer

MEDIUM

STAGE IV

VEGF

NEW TRIAL

VERIFIED

PUBLISHED MAR 11

Mar 12 13:19 · immediate · clinicaltrials.gov

A new clinical trial is exploring the combination of the anti-cancer drug CA-4948 with standard chemotherapy for patients with advanced metastatic colorectal cancer, potentially offering a new treatment option that targets tumor growth.

1 source(s) reporting · cluster 16 of 19

EVENT CLUSTER — Clinical Efficacy of Transarterial Infusion Chemotherapy for Unresectable Colorectal Cancer

MEDIUM

STAGE IV

GENERAL

NEW TRIAL

VERIFIED

PUBLISHED MAR 6

Mar 12 13:19 · immediate · clinicaltrials.gov

A new clinical trial shows that transarterial infusion chemotherapy may improve treatment outcomes for patients with unresectable Stage IV colorectal cancer targeting the EGFR pathway.

1 source(s) reporting · cluster 17 of 19

EVENT CLUSTER — Testing the Addition of Total Ablative Therapy to Usual Systemic Therapy Treatment for Limited Metastatic Colorectal Cancer, The ERASur Study

MEDIUM

STAGE IV

GENERAL

NEW TRIAL

VERIFIED

PUBLISHED MAR 4

Mar 12 13:19 · immediate · clinicaltrials.gov

The ERASur Study is exploring whether combining total ablative therapy with standard systemic treatment can improve outcomes for patients with limited metastatic colorectal cancer.

1 source(s) reporting · cluster 18 of 19

EVENT CLUSTER — Phase II Study of Neoadjuvant Dostarlimab in Patients With Untreated T3-4N0-2 or Stage III pMMR/MSS Resectable Colon Cancer

MEDIUM

STAGE III

GENERAL

NEW TRIAL

VERIFIED

PUBLISHED FEB 12

Mar 12 13:19 · immediate · clinicaltrials.gov

A new Phase II trial is investigating the use of the immune therapy dostarlimab before surgery in patients with advanced, resectable Stage III colon cancer, which may offer a promising new treatment option for improving outcomes.

1 source(s) reporting · cluster 19 of 19

Appendix A — Patient-Friendly Summary

Plain-language overview for patients, caregivers, and family members

What's happening

Recently, there have been some exciting developments in colon and colorectal cancer research. A new study has been approved to test if adding a drug called INCA33890 to standard chemotherapy can help patients with a specific type of metastatic colorectal cancer. Another trial is looking at an immunotherapy called tislelizumab as a treatment before surgery for patients with certain types of colorectal cancer. There are also trials exploring new combinations of existing treatments, like using a drug called pembrolizumab for Stage IV colorectal cancer, which could offer more effective options for patients. Overall, these studies aim to improve treatment outcomes and provide new hope for those facing advanced stages of the disease.

What this means for you

For patients, these developments could mean new options and better outcomes. If you or a loved one is dealing with colon cancer, it might be a good idea to ask your oncologist about these new trials and whether they could be relevant. Questions like, "Are there new treatments available?" or "How do these studies compare to my current treatment?" can help guide the conversation. Stay informed and hopeful, as research is continually evolving, and new options may be on the horizon that could make a difference in your treatment journey.

AI-generated in plain language from clinical data · 2026-03-12 14:19:58

This is not medical advice. Always speak with your oncologist before making any treatment decisions.

Appendix B — Colon & Colorectal Cancer Treatment Reference

Incidence & Trends

~150,000 new US cases/year · ~52,000 deaths/year · 4th most common US cancer

▲ Early-Onset Alert: CRC rates in adults under 50 have risen ~2% per year since the 1990s · Adults under 55 now account for ~20% of all new CRC diagnoses · Cause under active investigation (diet, microbiome, obesity, sedentary lifestyle)

Current Screening Guidelines (2025)

USPSTF / ACS / ACG — Average Risk: Begin screening at age 45 (lowered from 50 in 2021)

Colonoscopy every 10 years · FIT (fecal immunochemical test) annually · Cologuard every 1-3 years · CT colonography every 5 years

High Risk (family history / Lynch syndrome / IBD): Begin at age 40 or 10 years before youngest affected relative — whichever comes first · Colonoscopy every 1-5 years depending on risk

Symptoms at any age: Rectal bleeding, change in bowel habits >4 weeks, unexplained weight loss, iron deficiency anemia → immediate evaluation regardless of age

Key Biomarkers — Test All mCRC at Diagnosis

RAS (KRAS/NRAS) · BRAF V600E · MSI/MMR status · HER2 · NTRK fusion · TMB

Stage IV Targeted Options

KRAS G12C (~3-4%) → sotorasib + cetuximab · adagrasib + cetuximab

BRAF V600E (~8-10%) → encorafenib + cetuximab ± binimetinib

MSI-H/dMMR (~5%) → pembrolizumab (1st line preferred) · nivolumab ± ipilimumab

HER2 amplified (~2-3%) → trastuzumab + pertuzumab · trastuzumab deruxtecan

RAS/BRAF WT → FOLFOX or FOLFIRI + cetuximab or panitumumab (left-sided preferred)

1st Line mCRC Standard of Care

FOLFOX or FOLFIRI ± bevacizumab (RAS mutant / right-sided)

FOLFOX or FOLFIRI + cetuximab or panitumumab (RAS/BRAF WT, left-sided)

Pembrolizumab monotherapy (MSI-H/dMMR)

Key Resources

ClinicalTrials.gov · NCCN Guidelines (nccn.org) · Colorectal Cancer Alliance (ccalliance.org) · Fight Colorectal Cancer (fightcrc.org) · ACS (cancer.org)

Colon & Colorectal Cancer — Clinical Trial & Treatment Timeline

Alert classification fields are generated using deterministic rule-based analysis. Alert summaries are AI-generated from ClinicalTrials.gov, a registry and database maintained by the U.S. National Library of Medicine. This report aggregates publicly available trial registry data from U.S. government databases for situational awareness only. No registry content is reproduced verbatim. This report is made available under a free-use license for patient advocacy organizations, oncology practices, and individual patients. You may share or distribute it freely. You may not modify it, remove attribution, or republish it as your own work. Authorship and all intellectual property rights are retained by Randy Hinton. This report is provided for informational purposes only and does not constitute medical advice. Always consult your oncologist or a qualified healthcare provider before making any treatment decisions.

© 2026 Randy Hinton. All rights reserved. · https://petroai1492.github.io/delta/timelines/coloncancer_timeline.pdf