

Colon Cancer — Clinical Trial & Treatment Timeline

Alert Window: Last 30 Days · Generated: 2026-03-09 20:25:59

Situation Brief

Recent developments in colorectal cancer research highlight several promising treatment options for patients, particularly those with Stage IV disease and specific biomarkers. Notably, the FDA has approved the combination of encorafenib for patients with BRAF mutations, while new trials are investigating combinations of therapies such as Iparomlimab and Tuvonralimab for unresectable advanced cases, and low-dose radiotherapy with pucotenlimab for locally advanced rectal cancer. These advancements may influence treatment decisions by providing new first-line and combination therapy options. Moving forward, continued monitoring of trial results and FDA approvals will be critical for understanding the evolving landscape of colorectal cancer treatment.

AI-generated from deterministic classification data · 2026-03-09 20:25:59

Situation Summary

Events Detected: 12
High Significance: 3
Top Category: New Trial
Primary Stage: Stage IV
Primary Target: General
Research Activity: Active Monitoring

Event Timeline

EVENT CLUSTER — Low-Dose Radiotherapy to Sensitize Pucotenlimab Plus CAPEOX for pMMR Locally Advanced Rectal Cancer

MEDIUM

STAGE IV

GENERAL

NEW TRIAL

CORROBORATED

Mar 9 19:24 · immediate · clinicaltrials.gov

A new trial is exploring the use of low-dose radiotherapy to enhance the effectiveness of the drug pucotenlimab combined with CAPEOX chemotherapy for patients with locally advanced rectal cancer, potentially offering a more effective treatment option for those with this condition.

1 source(s) reporting · cluster 1 of 10

EVENT CLUSTER — A Clinical Study of Iparomlimab and Tuvonralimab Combined With Bevacizumab and Alternating Triweekly CAPOX/mCAPIRI Regimen as First-line Treatment for Unresectable Advanced Colorectal Cancer

MEDIUM STAGE IV VEGF NEW TRIAL CORROBORATED

Mar 9 19:24 · immediate · [clinicaltrials.gov](#)

A new clinical trial is exploring a combination of Iparomlimab, Tuvonralimab, and Bevacizumab with a chemotherapy regimen for patients with advanced colorectal cancer, potentially offering a new first-line treatment option for those with unresectable disease.

1 source(s) reporting · cluster 2 of 10

EVENT CLUSTER — Everolimus in CDK12-Deficient Metastatic Colorectal Cancer (EVER-RECODE)

MEDIUM STAGE IV GENERAL NEW TRIAL CORROBORATED

Mar 9 19:24 · immediate · [clinicaltrials.gov](#)

A new clinical trial is exploring the use of everolimus for patients with Stage IV metastatic colorectal cancer that has CDK12 deficiencies, potentially offering a new treatment option for this specific group.

1 source(s) reporting · cluster 3 of 10

EVENT CLUSTER — Effect of Perioperative Lidocaine or High-Dose Dexamethasone on Immune Response in Colon Cancer Surgery (PILDI Study)

MEDIUM ALL STAGES GENERAL NEW TRIAL CORROBORATED

Mar 9 19:24 · immediate · [clinicaltrials.gov](#)

The PILDI Study is exploring whether perioperative lidocaine or high-dose dexamethasone can enhance the immune response in patients undergoing colon cancer surgery, potentially leading to improved recovery and outcomes.

1 source(s) reporting · cluster 4 of 10

EVENT CLUSTER — Neoadjuvant Chemoradiotherapy Plus Tislelizumab With or Without Probio-M9 in pMMR/MSS Locally Advanced Rectal Cancer

MEDIUM STAGE IV GENERAL NEW TRIAL CORROBORATED

Mar 9 19:24 · immediate · [clinicaltrials.gov](#)

A new clinical trial is exploring the combination of neoadjuvant chemoradiotherapy and the immunotherapy drug Tislelizumab, with or without the addition of Probio-M9, for patients with locally advanced rectal cancer, potentially offering a new treatment option for those with this challenging disease.

1 source(s) reporting · cluster 5 of 10

EVENT CLUSTER — FOLFIRI and Bevacizumab With or Without Pelareorep for Second-Line Treatment of Metastatic RAS-Mutated, Microsatellite-Stable Colorectal Cancer

MEDIUM STAGE IV VEGF NEW TRIAL CORROBORATED

Mar 9 19:24 · immediate · clinicaltrials.gov

A new clinical trial is exploring whether adding Pelareorep to the standard treatment of FOLFIRI and Bevacizumab can improve outcomes for patients with advanced colorectal cancer that has specific genetic characteristics.

1 source(s) reporting · cluster 6 of 10

EVENT CLUSTER — FDA Approves Encorafenib for Metastatic BRAF V600E–Mutated Colorectal Cancer

HIGH STAGE IV BRAF TRIAL RESULTS VERIFIED

Mar 9 13:57 · immediate · [Oncology News Central](#), [The ASCO Post](#), [The ASCO Post](#)

The FDA's approval of encorafenib offers a new targeted treatment option for patients with metastatic colorectal cancer harboring the BRAF V600E mutation, potentially improving outcomes for this challenging group.

3 source(s) reporting · cluster 7 of 10

EVENT CLUSTER — Final Phase 3 CONVERT Trial Results: Neoadjuvant CAPOX vs Capecitabine-Based ChemoRT in Locally Advanced Rectal Cancer

HIGH STAGE IV GENERAL TRIAL RESULTS UNVERIFIED

Mar 9 13:40 · immediate · [Oncodaily](#)

The final results of the Phase 3 CONVERT trial indicate that neoadjuvant CAPOX may offer improved treatment outcomes for patients with locally advanced rectal cancer compared to traditional capecitabine-based chemotherapy and radiation, potentially expanding options for effective pre-surgical therapy.

1 source(s) reporting · cluster 8 of 10

EVENT CLUSTER — Adding Encorafenib and Cetuximab to FOLFIRI Shows Benefit in BRAF-Mutant Colorectal Cancer

MEDIUM ALL STAGES BRAF TRIAL RESULTS VERIFIED

Mar 9 13:40 · immediate · [The ASCO Post](#)

Combining Encorafenib and Cetuximab with the chemotherapy regimen FOLFIRI has been shown to improve treatment outcomes for patients with BRAF-mutant colorectal cancer.

1 source(s) reporting · cluster 9 of 10

EVENT CLUSTER — Oncolytics Biotech (NASDAQ: ONCY) launches Phase 2 pelareorep colorectal cancer study

MEDIUM

ALL STAGES

GENERAL

NEW TRIAL

UNVERIFIED

Mar 8 9:15 · immediate · Stock Titan

Oncolytics Biotech has initiated a Phase 2 study of pelareorep for colorectal cancer, which could provide new treatment options for patients at all stages of the disease.

1 source(s) reporting · cluster 10 of 10

Colon Cancer Treatment Reference

Incidence & Trends

~150,000 new US cases/year · ~52,000 deaths/year · 4th most common US cancer

▲ Early-Onset Alert: CRC rates in adults under 50 have risen ~2% per year since the 1990s · Adults under 55 now account for ~20% of all new CRC diagnoses · Cause under active investigation (diet, microbiome, obesity, sedentary lifestyle)

Current Screening Guidelines (2025)

USPSTF / ACS / ACG — Average Risk: Begin screening at age 45 (lowered from 50 in 2021)

Colonoscopy every 10 years · FIT (fecal immunochemical test) annually · Cologuard every 1-3 years · CT colonography every 5 years

High Risk (family history / Lynch syndrome / IBD): Begin at age 40 or 10 years before youngest affected relative — whichever comes first · Colonoscopy every 1-5 years depending on risk

Symptoms at any age: Rectal bleeding, change in bowel habits >4 weeks, unexplained weight loss, iron deficiency anemia → immediate evaluation regardless of age

Key Biomarkers — Test All mCRC at Diagnosis

RAS (KRAS/NRAS) · BRAF V600E · MSI/MMR status · HER2 · NTRK fusion · TMB

Stage IV Targeted Options

KRAS G12C (~3-4%) → sotorasib + cetuximab · adagrasib + cetuximab

BRAF V600E (~8-10%) → encorafenib + cetuximab ± binimetinib

MSI-H/dMMR (~5%) → pembrolizumab (1st line preferred) · nivolumab ± ipilimumab

HER2 amplified (~2-3%) → trastuzumab + pertuzumab · trastuzumab deruxtecan

RAS/BRAF WT → FOLFOX or FOLFIRI + cetuximab or panitumumab (left-sided preferred)

1st Line mCRC Standard of Care

FOLFOX or FOLFIRI ± bevacizumab (RAS mutant / right-sided)

FOLFOX or FOLFIRI + cetuximab or panitumumab (RAS/BRAF WT, left-sided)

Pembrolizumab monotherapy (MSI-H/dMMR)

Key Resources

ClinicalTrials.gov · NCCN Guidelines (nccn.org) · Colorectal Cancer Alliance (ccalliance.org) · Fight Colorectal Cancer (fightcrc.org) · ACS (cancer.org)

Colon Cancer — Clinical Trial & Treatment Timeline

Alert classification fields are generated using deterministic rule-based analysis. Alert summaries are AI-generated from publicly available news headlines. This report aggregates headline-level information for situational awareness only. No article text is reproduced. All original content remains the property of respective publishers. This report is provided for informational purposes only and does not constitute medical advice. Always consult your oncologist or a qualified healthcare provider before making any treatment decisions.

